

② 19809/B

Charles Barker
1840

THE
COLLECTED WORKS
OF
SIR HUMPHRY DAVY, BART.

77349

THE
COLLECTED WORKS

OF

SIR HUMPHRY DAVY, BART.

LL.D. F.R.S.

FOREIGN ASSOCIATE OF THE INSTITUTE OF FRANCE, ETC.

EDITED BY HIS BROTHER,

JOHN DAVY, M.D. F.R.S.

VOL. IV.

ELEMENTS OF CHEMICAL PHILOSOPHY.

LONDON:

SMITH, ELDER AND CO. CORNHILL.

1840.

ELEMENTS

OF

CHEMICAL PHILOSOPHY:

AS REGARDS

THE LAWS OF CHEMICAL CHANGES: UNDECOMPOUNDED
BODIES AND THEIR PRIMARY COMBINATIONS.

BY

SIR HUMPHRY DAVY, BART.

LONDON:

SMITH, ELDER AND CO. CORNHILL.

1840.

298

8

ALFRED

CHEMICAL PHOTOGRAPHY

THEORY AND PRACTICE OF CHEMICAL PHOTOGRAPHY
BY ALFRED M. MURRAY, F.R.S.

THE HURST AND BLACKET CO. LTD.

LONDON:

PRINTED BY STEWART AND MURRAY, OLD BAILEY.

[AFTER some consideration, it has appeared most advisable that the Elements of Chemical Philosophy, in the order of appearance of the Collected Works, should precede rather than follow the Author's most important Researches, contained in his Bakerian Lectures and other Papers communicated to the Royal Society in rapid succession, after 1806.

According to this order, the Elements of Chemical Philosophy will be a preparation for these Researches: the Student of Chemistry may derive from the one that degree of general elementary information which may enable him, with ease and profit, to apply himself to the other.

Another reason for adopting this course is, that the Author himself contemplated such a plan, as has been already mentioned in the Memoir of his Life, with expressions of regret that he did not carry his intention into effect, and as will be seen in his own Advertisement to the present Volume.

This Volume was published in 1812, when he was about to retire from the Royal Institution and to enter into a new course of life. The circumstances under which, and the manner in which the Work was composed, have been already alluded to. At its first appearance it excited much attention. In an article in the Quarterly Review, No. 15, written by a very able judge, the late Dr. Thomas Young, the researches on which it was based were pronounced to be "more splendidly successful than any which have ever before illustrated the physical sciences in any of their departments," not even excepting the optics of Newton.

Almost immediately after its publication, the Author commenced preparations for another edition: the Advertisement which follows was then written, and some few additions and alterations, which will be introduced into the text, were then made. In this Advertisement it will be perceived he states his views relative to Chemical Nomenclature; and mentions an intention in relation to it, viz., the inserting in the form of notes at the bottom of the page, the appellations which he wished to suggest for use; but which, as he did not give, I have not thought it right to introduce: any Reader who chooses, having a knowledge of the principle, can supply the deficiency.]

TO
LADY DAVY.

THERE is no individual to whom I can with so much propriety or so much pleasure dedicate this Work as to you. The interest you have taken in the progress of it, has been a constant motive for my exertions ; and it was begun and finished in a period of my life which, owing to you, has been the happiest. Regard it as a pledge that I shall continue to pursue Science with unabated ardour. Receive it as a proof of my ardent affection, which must be unalterable, for it is founded upon the admiration of your moral and intellectual qualities.

H. DAVY.

ADVERTISEMENT.

IN this work I have found great difficulties in adapting the nomenclature of chemistry to the progress of discovery. Many of the names which have been long current now imply erroneous ideas; it is absolutely necessary that they should be changed; but it is difficult to discover unexceptionable principles on which these alterations may be founded.

In the Bakerian lecture for 1810 I proposed the name chlorine for oxymuriatic gas, because no oxygen has as yet been discovered in this substance; and I ventured to suggest that the combinations of this substance with metals and inflammable bodies might be signified by the name of the base with a termination in *ane*, *ana*, and *anée*; *ane* signifying that the substance contained one proportion of chlorine; *ana*, two proportions; and *anée*, three.

In the first edition of these elements, I employed these forms principally with the view of guarding students against the prejudice which the use of the common

names must leave in the mind, and to prevent them from assuming the existence of oxygen and muriatic acid in bodies from which they have not been obtained. I do not, however, mean to press the adoption of them : the subject is open to discussion. Some persons think if oxymuriatic gas be called chlorine, that the compounds of it should be named chlorides, and others conceive it more in conformity to the existing nomenclature to call oxymuriatic gas murine or murigen, and its compounds murides. If the old nomenclature be preserved entire in its other parts, either of the two last plans will be more in harmony with the arrangement belonging to it ; but if it should be changed so as to be made less theoretical by adopting terminations to signify classes and not names to denote the chemical nature of compounds, which I conceive would be for the advantage of science, then the plan that I have proposed under certain modifications may be made part of an arrangement founded on the same general principles.

The fluates as well as the muriates appear to form a peculiar class of bodies ; I have not been able to detect any oxygen in them, and it is most probable that they consist of a peculiar principle analogous to oxygen and chlorine united to metals and inflammable bases.

The earths and alkalies, which are oxidated bodies, are at present called by names differing from the names of their bases only by their terminations ; thus potassium and potassa are names which express the relations

of the two substances without implying any theoretical views.

The same plan might be applied to the other oxidated compounds, and the deutoxides containing 2 proportions of oxygen signified by terminations in *e*, those containing 3 proportions by terminations in *i*, and so on. Thus in a scientific nomenclature the term *ferrum* may be adopted for iron; *ferré* for the deutoxide of iron, and *ferrî* for the tritoxide, or red oxide.

If the terminations in *ane*, *ana*, *ani*, &c., or *ana*, *ane*, *ani*, be applied to the combinations of oxymuriatic gas, those of *ile*, *ila*, *ili*, or of *ita*, *ite*, *iti*, might be made to signify that a compound contained 1, 2, or 3 proportions of the fluoric principle; thus calcile or calcila would signify fluor spar; potassile or potassila would signify the substance improperly called fluuate of potassa.

According to the theoretical forms of the old nomenclature, if chloride or muride be adopted, the combinations of the fluoric principle may be named fluorids; thus fluor spar may be called fluorid of calcium.

In this edition I have given in the text the names commonly used, and have added in notes the new names which may be applied to them according to the two different principles of nomenclature. For the sake of uniformity, I have made the first combination in the three classes of oxides, muriates, and fluates, to terminate in *a*; and in the second plan I have adopted Dr. Thomson's method of signifying different compounds

of the same principle, by prefixing to them syllables derived from the Greek; thus, deutoxide, tritoxide, tetroxide, signify that the bodies contain 2, 3 or 4 proportions of oxygen.

The true salts, whether they have alkalies, earths, or metallic oxides for their bases, are known to be substances of the same kind, and the nomenclature for them ought to be made analogous; thus, if the name sulphate of potassa be continued, the substance called sulphate of lead should be named sulphate of deutoxide of lead, or sulphate of plumbi.

Whatever pains be taken, it will not be possible to make the existing nomenclature conformable to the idiom of our language. At no very distant period it is to be hoped that the enlightened chemists of Europe will agree to adopt some forms from the dead languages, which shall be common to all countries, and which from their nature may be permanent. Till this is done, the language of chemistry must be very imperfect, without unity, and composed of shreds and patches; innovation will be censured even when it is absolutely necessary, and neology a constant reproach.

I have in a few instances only detailed the experiments, from the results of which the numbers furnished by myself representing the undecomposed bodies were calculated.

To have given accurate histories of those experiments would have been incompatible with the object of an

elementary book devoted to the general truths and methods of the science. I shall, however, shortly present them to the public in a work containing an account of the labours that I have carried on during the last twelve years in analytical chemistry.

I have usually given whole numbers, taking away or adding fractional parts, that they may be more easily retained in the memory. When the number was gained from experiments in which a loss might be supposed, I have added fractional parts, so as to make a whole number. Thus the number representing Barium is nearer 129 than 130; but it is given as 130, because it was deduced from an indirect experiment, in which a loss of weight was more probable than an increase from any foreign source.

The numbers gained by calculations on the results of other chemists are likewise given. Where many different results are found to agree very nearly, the mean may be taken as the correct approximation to the true number; where the results of accurate experiments differ considerably, it may be concluded that some circumstance respecting the nature of the body remains to be discovered.*

I cannot conclude without acknowledging my obligations to my brother, Mr. John Davy, for the able

* [This advertisement for a new edition was dated, Paris, Oct. 31, 1813. The following concluding paragraphs belonged to the advertisement to the first, with the date of Berkeley Square, June 1, 1812.]

assistance he afforded me in the progress of the researches which form the foundation of this treatise.

I have likewise received much useful experimental aid from Mr. E. Davy, and Mr. W. Moore.

The greater number of the experiments were made in the laboratory of the Royal Institution; and all that were fitted for demonstration have been exhibited in the Theatre of that useful public establishment in my annual courses of lectures; and have been received by the members in a manner which I shall always remember with gratitude.

CONTENTS.

INTRODUCTION.

HISTORICAL VIEW OF THE PROGRESS OF CHEMISTRY.

ON THE LAWS OF CHEMICAL CHANGES; ON UNDECOMPOUNDED BODIES, AND THEIR PRIMARY COMBINATIONS.

DIVISION I.

ON THE POWERS AND PROPERTIES OF MATTER, AND THE GENERAL LAWS OF CHEMICAL CHANGES.

	Page
1. Preliminary Observations	43
2. Of the Forms of Matter	44
3. Gravitation	46
4. Cohesion	47
5. Of Heat, or Calorific Repulsion	48
6. On Chemical Attraction, and the Laws of Combination and Decomposition	69
7. Of Electrical Attraction and Repulsion, and their Relations to Chemical Changes	91
8. On Analysis and Synthesis; on the Circumstances to be attended to in these Operations, and on the Arrangement of Undecomposed Bodies	131

DIVISION II.

OF RADIANT OR ETHEREAL MATTER.

1. Of the Effects of Radiant Matter, in producing the Phenomena of Vision	141
---	-----

	Page
2. Of the Operation of Radiant Matter in producing Heat	146
3. Of the Effects of Radiant Matter in producing Chemical Changes	154
4. Of the Nature of the Motions or Affections of Radiant Matter	157

DIVISION III.

OF EMPYREAL UNDECOMPOUNDED SUBSTANCES, OR UNDECOMPOUNDED SUBSTANCES THAT SUPPORT COMBUSTION, AND THEIR COMBINATION WITH EACH OTHER.

1. General Observations	165
2. Of Oxygen Gas	167
3. Chlorine, or Oxymuriatic Gas	173

DIVISION IV.

OF UNDECOMPOUNDED INFLAMMABLE OR ACIDIFEROUS SUBSTANCES NOT METALLIC, AND THEIR BINARY COMBINATIONS WITH OXYGEN AND CHLORINE, OR WITH EACH OTHER.

1. Preliminary Observations	180
2. Hydrogen Gas, or Inflammable Air	181
3. Of Azote, or Nitrogen Gas	188
4. Of Sulphur	200
5. Of Phosphorus	210
6. Of Carbon or Charcoal, and the Diamond	221
7. Of Boron, or the Boracic basis	232

DIVISION V.

OF METALS ; THEIR PRIMARY COMBINATIONS WITH OTHER UNDECOMPOUNDED BODIES, AND WITH EACH OTHER.

1. General Observations	235
2. Of Potassium	237
3. Sodium	244
4. Barium	249
5. Strontium	253
6. Calcium	255
7. Magnesium	258
8. Aluminum	262

CONTENTS.

xix

	Page
9. Glucinum	264
10. Zirconum	266
11. Silicum	268
12. Ittrium	270
13. Manganesum	271
14. Zinc, or Zincum	276
15. Tin, or Stannum	281
16. Iron, or Ferrum	285
17. Lead, or Plumbum	292
18. Antimony, or Antimonium	296
19. Bismuth, or Bismuthium	300
20. Tellurium	303
21. Cobalt, or Cobaltum	305
22. Copper, Cuprum	308
23. Nickel, or Nickolum	312
24. Uranium	315
25. Osmium	316
26. Tungsten, or Tungstenum	317
27. Titanium	319
28. Columbium	321
29. Cerium	322
30. Palladium	323
31. Iridium	324
32. Rhodium	326
33. Mercury, or Mercurium	327
34. Silver, or Argentum	330
35. Gold, or Aurum	333
36. Platinum	334
37. Arsenic, or Arsenicum	338
38. Molybdenum	342
39. Chromium	345

DIVISION VI.

OF SOME SUBSTANCES, THE NATURE OF WHICH IS NOT YET
CERTAINLY KNOWN.

1. Preliminary Observations	347
2. Of the Fluoric Principle	327
3. Of the Amalgam produced from Ammoniacal Compounds	353

DIVISION VII.

ON THE ANALOGIES BETWEEN THE UNDECOMPOUNDED SUBSTANCES ; SPECULATIONS RESPECTING THEIR NATURE ; ON THE MODES OF SEPARATING THEM, AND ON THE RELATIONS OF THEIR COMPOUNDS.

	Page
1. Of the Analogies between the Undecomposed Substances : ideas respecting their Nature . . .	356
2. Of the Analogies between the Properties of the Primary Compounds, and on their Chemical Relations .	365
3. On the Relative Attractions of the Undecomposed Substances for each other	370
4. On the Methods of separating the Undecomposed Bodies from each other	372
5. General Observations, and Conclusion	374

ELEMENTS

OF

CHEMICAL PHILOSOPHY.

INTRODUCTION.

HISTORICAL VIEW OF THE PROGRESS OF CHEMISTRY.

MOST of the substances belonging to our globe are constantly undergoing alterations in sensible qualities, and one variety of matter becomes as it were transmuted into another.

Such changes, whether natural or artificial, whether slowly or rapidly performed, are called chemical; thus the gradual and almost imperceptible decay of the leaves and branches of a fallen tree exposed to the atmosphere, and the rapid combustion of wood in our fires, are both chemical operations.

The object of Chemical Philosophy is to ascertain the causes of all phenomena of this kind, and to discover the laws by which they are governed.

The ends of this branch of knowledge are the applications of natural substances to new uses, for increasing the comforts and enjoyments of man, and the demon-

stration of the order, harmony, and intelligent design of the system of the earth.

The foundations of chemical philosophy are, observation, experiment, and analogy. By observation, facts are distinctly and minutely impressed on the mind. By analogy, similar facts are connected. By experiment, new facts are discovered; and, in the progression of knowledge, observation, guided by analogy, leads to experiment, and analogy confirmed by experiment, becomes scientific truth.

To give an instance.—Whoever will consider with attention the slender green vegetable filaments (*Conferva rivularis*) which in the summer exist in almost all streams, lakes, or pools, under the different circumstances of shade and sunshine, will discover globules of air upon the filaments exposed under water to the sun, but no air on the filaments that are shaded. He will find that the effect is owing to the presence of light. This is an *observation*; but it gives no information respecting the nature of the air. Let a wine glass filled with water be inverted over the conferva, the air will collect in the upper part of the glass, and when the glass is filled with air, it may be closed by the hand, placed in its usual position, and an inflamed taper introduced into it; the taper will burn with more brilliancy than in the atmosphere. This is an *experiment*. If the phenomena are reasoned upon, and the question is put, whether all vegetables of this kind, in fresh or in salt water, do not produce such air under like circumstances, the inquirer is guided by *analogy*: and when this is determined to be the case by new trials, a *general scientific truth* is established—That all confervæ in the sunshine produce a species of air that supports flame in a superior degree; which has been shown to be the case by various minute investigations.

These principles of research, and combinations of methods, have been little applied, except in late times. A transient view of the progress of chemical philosophy will prove that the most brilliant discoveries, and the happiest theoretical arrangements belonging to it are of very recent origin; and a few historical details and general observations upon the progress and effects of the science will form, perhaps, no improper introduction to the elements of this branch of knowledge.

The only processes which can be called chemical, known to the civilized nations of antiquity, belonged to certain arts, such as metallurgy, dyeing, and the manufacture of glass or porcelain; but these processes appear to have been independent of each other, pursued in the workshop alone, and unconnected with general knowledge.

In the early mythological systems of the Egyptian priests, and the Brahmins of Hindostan, some views respecting the chemical changes of the elements seem to have been developed, which passed, under new modifications, into the theories of the Greeks; but as the most refined doctrines of this enlightened people, concerning natural causes, in their best times, were little more than a collection of vague speculations, rather poetical than philosophical, it cannot well be supposed that in earlier ages, and amongst nations less advanced in cultivation, there were any traces of genuine science.

The inhabitants of Lower Egypt, where the overflowing of the Nile covered a sandy desert with vegetation and life, might easily adopt the notion, that water, in different modifications, produced all the varieties of inanimate and organized matter; and this dogma characterized the earliest school of Greece.

To generalize upon the great forms or powers of nature, as elements, requires only very superficial observation; and hence the theories seem to have originated, which have been attributed to Anaximander, and others of the early Greek philosophers, concerning air, earth, water, and fire.

As geometry and the mathematical sciences became improved, mechanical solutions of the changes of bodies were natural consequences, such as the atomic philosophy of the Ionian sect, and the five regular solids assumed by the Pythagoreans as the materials of the universe.

In the beginning of the Macedonian dynasty, the school of Aristotle gave a transient attention to the objects of natural science, but the great founder attempted too many subjects to be able to offer correct views of any one series. And his erroneous practice, that of advancing general principles, and applying them to particular instances, so fatal to truth in all sciences, more particularly opposed itself to the progress of one founded upon a minute examination of obscure and hidden properties of natural bodies.*

Theophrastus, the successor of Aristotle, did not, it appears, adopt the sublime, though purely speculative doctrine of his master, the identity of matter, and its

* [These remarks are to be considered as applicable to the ancient dialectic method, opposed to the experimental and inductive method of the moderns,—or that of the school of Aristotle in contradistinction to that of Bacon,—not the science which is effective and fruitful of works,—the knowledge which is power; but that which deals more in words than in facts, and gives skill chiefly in disputation,—“*Sapientiam istam, quam à Græcis potissimum hausimus, pueritiam quandam scientiæ videri, atque habere quod proprium est puerorum; ut ad garriendum prompta, ad generandum invalida et immatura sit. Controversiarum enim ferax, operum effæta est.*”—*Bacon's Preface to his Instauratio Magna.*]

diversity of form ;*—for he says, in the beginning of his book concerning fossils, ‘stones are produced from earth, metals from water.’† How such a notion as the last could have been formed, it is difficult to discover ; yet, Theophrastus is perhaps the best observer amongst the ancients, whose works are in our possession, and the theories of this distinguished teacher, who is said to have had a class of 2000 pupils, cannot be considered as an unfavourable specimen of the theoretical physics of the age.

In all pursuits which required only the native powers of the intellect, or the refinements of taste, the Greeks were pre-eminent ; their literature, their works of art, offer models which have never been excelled. They possessed, as if instinctively, the perception of every thing beautiful, grand, and decorous. As philosophers, they failed not from a want of genius, or even of application, but merely because they pursued a false path, —because they reasoned more upon an imaginary system of nature, than upon the visible and tangible universe.

It will be in vain to look in the annals of Rome for science, that did not exist in Greece. The conquerors became the pupils of the conquered ; and the Romans did little more than clothe the systems of their masters in a new dress, and adapt them to a new people.

The grand, but unequal poem of Lucretius, contains the abstract of the opinions of Epicurus, compared with those of other celebrated teachers. The Natural History of Pliny, is a collection from all sources, but principally from Theophrastus and Aristotle. The details

* Επειδὲ δὲ ἡ φύσις διχῶς, τὸ τε εἶδος καὶ ἡ ὕλη. Aristotelis Natural. Auscult. Lib. ii. 495, fol. Par. 1654.

† Ὑδατος μὲν τὰ μελλευνόμενα καθάπερ ἄργυρος καὶ χρυσοῦς, καὶ τὰ λαλα. γῆς δὲ λίθος τε καὶ ὅσα λίθων περιπτώτερα. Theophrasti de Lapidibus. Lug. Bat. 1613.

from his own observation are more interesting when they relate to artificial, than when they refer to natural operations; the speculative notions are of the rudest kind. The earlier philosophical work of the Romans, as if indicative of the youth of the people, is marked by power and genius, by boldness and incorrectness; the later, as if it belonged to their old age, by garrulity, copious and amusing anecdote, superstitious notions, and vulgar prejudices.

Some of the historians of this science,* in their zeal for the honour of its antiquity, have indeed endeavoured to find instances of an acquaintance with some doctrines of practical chemistry, at least, amongst the ancients.—Thus Democritus is quoted by Laertius as having employed himself in processes for imitating gems, and for softening and working ivory. Caligula is said to have made experiments with the view of extracting gold from orpiment.—Dioscorides, who is supposed to have been physician to the celebrated Cleopatra, has described the process of subliming mercury from its ores.—Even Cleopatra herself, on the evidence of such circumstances, might be considered as an experimenter, because in the madness of profusion, she dissolved a pearl in vinegar, and made a nauseous draught of a

* Many of the alchemical writers derive alchemy from Tubal Cain; others from Hermes Trismegistus, the Mercury of the Greeks. The first writing specifically on a chemical subject, is a manuscript supposed to be of the fifth century, by Zosimus, on the art of making gold and silver; which was in the king's library at Paris. Suidas, who wrote in the ninth or tenth century, mentions Diocletian as having burnt the books of the Egyptians concerning the chemistry of silver and gold: “περι χυμείας αργυρου και χρυσου.” *Lexicon*, Tom. i. p. 595.

For a minute investigation of the claims of the ancients to chemical knowledge, the reader may consult Borrichius de Ortu et Progress. Chæm. Bergman. *Opuscula*, vol. iv. de primordiis Chæm. and Lenglet Dufrenoy, *Histoire de la Philosophie hermetique*.

costly and beautiful substance; but it is idle to relate such circumstances as indications of *science*. If chemical operations had been known to any extent, beyond their mere relations to the arts, some mention of them might have been expected in the medical writings of those times; but not even distillation is noticed in the works of Hippocrates or Galen; and the same Dioscorides who has just been alluded to, and who probably possessed whatever knowledge was at that time extant in Egypt, recommends the use of a fleece of wool or a sponge, for collecting the products from boiling or burning substances.*

The origin of chemistry, as a science of experiment, cannot be dated farther back than the seventh or eighth century of the Christian era, and it seems to have been coeval with the short period in which cultivation and improvements were promoted by the Arabians.

The early Mahometans endeavoured to destroy all the records of the former progress of the human mind; and, as if to make compensation for this barbarian spirit, the same people were destined, in a more advanced period, to rekindle the light of letters, and to become the inventors and cultivators of a new science.

The early nomenclature of chemistry demonstrates how much it owes to the Arabians.—The words alcohol, alkahest, aludel, alembic, alkali, require no comment.

The first Arabian systematic works on chemistry are said to have been composed by Geber in the reigns of the caliphs Almainon and Almanzor. The preparation of medicines seems to have been the primary object in this study; and Rhases, Avicenna, and Avenzoar, who

* Discoridis, liber 1. de picino oleo, p. 52.

have described various chemical operations in their works, were the celebrated physicians of the age.

Amongst a people of conquerors, disposed to sensuality and luxury even from the spirit of their religion, and romantic and magnificent in their views of power, it was not to be expected that any new knowledge should be followed in a rational and philosophical manner; and the early chemical discoveries led to the pursuit of alchemy, the objects of which were to produce a substance capable of converting all other metals into gold: and an universal remedy calculated indefinitely to prolong the period of human life.

Reasonings upon the nature of the metals, and the composition of the philosopher's stone, form a principal part of the treatises ascribed to Geber;* and the disciples of the school of Bagdat seem to have been the first professed alchemists.

It required strong motives to induce men to pursue the tedious and disgusting processes of the furnace; but labourers could hardly be wanting, when prospects so brilliant and magnificent were offered to them; the means of procuring unbounded wealth; of forming a paradise on earth; and of enjoying an immortality depending on their own powers.

* The library of the British Museum contains several works bearing the name of Geber: amongst them are *De Alchemia argentea*, *Speculum Alchemiæ*, et de *Inventione perfectionis*: but they appear to be compilations formed by alchemists of the 15th and 16th centuries. Arsenic, mercury, and sulphur, are considered in them as elements of the metals; distillation is distinctly described. Alcohol, corrosive sublimate, and different saline combinations of iron, tin, copper, and lead are mentioned in them; but they abound in obscure descriptions of mysterious processes, and contain an account of some impracticable experiments.—The *Liber Fornacum* is the most intelligible part of the works ascribed to Geber; it contains a description of several metallurgical operations, and of the common apparatus of the assayer.

The processes supposed to relate to the transmutation of metals, and the elixir of life, were probably first made known to the Europeans during the time of the crusades; and many of the warriors who, animated with visionary plans of conquest, fought the battles of their religion on the plains of Palestine, seemed to have returned to their native countries under the influence of a new delusion.

The public spirit in the West, was calculated to assist the progress of all pursuits that carried with them an air of mysticism. Warm with the ardour of an extending and exalted religion, men were much more disposed to believe than to reason;—the love of knowledge and power is instinctive in the human mind; in darkness it desires light, and follows it with enthusiasm even when appearing merely in delusive glimmerings.

The records of the middle ages contain a great variety of anecdotes relating to the transmutation of metals, and the views or pretensions of persons considered as adepts in alchemy; these early periods constitute what may be regarded as the heroic or fabulous ages of chemistry. Some of the alchemists were low impostors, whose object was to delude the credulous and the ignorant; others seemed to have deceived themselves with vain hopes; but all followed the pursuit as a secret and mysterious study. The processes were communicated only to chosen disciples, and being veiled in the most enigmatic and obscure language, their importance was enhanced by the concealment. In all times men are governed more by what they desire or fear, than by what they know; and in this age it was peculiarly easy to deceive, but difficult to enlighten, the public mind; truths were discovered, but they were blended with the false and marvellous; and another era was required to

separate them from absurdities, and to demonstrate their importance and uses.

Arnald of Villa Nova, who is said to have died in 1250, was one of the earliest European inquirers who attended to chemical operations. In the edition of the works ascribed to him, published at Leyden in 1509,* there are several treatises on alchemical subjects, which shew that he firmly believed in the transmutation of metals; the same opinions are attributed to him and to Geber; and he seems to have followed the study with no other views than those of preparing medicines, and attempting the composition of the philosopher's stone.

Raymund Lully of Majorca is said to have been a disciple of Arnald, and applied himself much more than his instructor to philosophy; but the works on general science, ascribed to him, are more abundant in abstract metaphysical propositions, than in facts; he followed, in his physical views, the plan of Aristotle, and our opinion of his chemical talents cannot be very exalted, if the alchemical treatises bearing his name be regarded as genuine documents.

Arnald and Lully are both celebrated by the vindicators of alchemy, as having been certainly possessed of the secret of transmutation. Arnald is said to have converted iron into gold at Rome; and it is pretended that Lully performed a similar operation before Edward I. in London, of which gold nobles were said to have been made.†

That the delusions of alchemy were ardently pursued at this time may be learned from a reference to the public acts of these periods. Pope John XXII. who was raised to the pontificate in the year 1316,

* Opera Arnaldi de Villa Nova, fol. 1509.

† Bergman. Opuscula, tom. iv. p. 126.

openly condemned the alchemists as impostors, and the bull begins by stating, that “they promise what they do not perform;” and in England an act of Parliament was passed in the fifth year of the reign of Henry IV. prohibiting the attempts at transmutation, and making them felonious.*

Even in these times, however, there were some few efforts to form scientific views. In the beginning of the thirteenth century, Roger Bacon of Oxford applied himself to experiment, and his works offer proofs of talents, industry, and sagacity. He was a man of a truly philosophical turn, desirous of investigating nature, and of extending the resources of art, and his inquiries offered some very extraordinary combinations; but neither his labours, nor those of Albert of Cologne, his contemporary, who appears to have been a genius of a kindred character, had any considerable influence on the improvement of their age. The wonders performed by the experimental art were attributed by the vulgar to magic; and at a time when knowledge belonged only to the cloister, any new philosophy was of course regarded even by the learned with a jealous eye.

It would be a labour of little profit to dwell upon the works of the professed alchemists of the fourteenth and fifteenth centuries, of Richard and Ripley in England, Isac in Holland, Pico of Mirandula and Koffsky, in Poland. The works attributed to these persons are of similar stamp,† and contain nothing which can either instruct

* Lord Coke calls this act the shortest he ever met with, 5 H. IV. Statutes at Large, vol. i. p. 457. “None from henceforth shall use to multiply gold or silver, or use the craft of multiplication, and if any the same do, he shall incur the pain of felony.”

† Amongst them are Ricardi Angli Libellus, *περι χημειας*; Opus Saturni Johan. Isac; Compounde of Alchemy by George Ripley.

or amuse an intelligent reader. Basil Valentine of Erfurt deserves to be separated from the rest of the inquirers of this age, on account of the novelty and variety of his experiments on metallic preparations, particularly antimony: in his *Currus triumphalis Antimonii* he has described a number of the combinations of this metal. He used the mineral acids for solutions, and seems to have been one of the first persons who observed the production of ether from alcohol. He flourished about the year 1413.

Cornelius Agrippa, who was born at Cologne in 1486, openly professed magic, and endeavoured to connect together judicial astrology, the hermetic art, and metaphysical philosophy; and he was followed by Paracelsus, in Switzerland, and Digby, Kelly, and Dee, in England.

The first Arabian alchemists seem to have adopted the idea, that the elements were under the dominion of spiritual beings, who might be submitted to human power; and the notions of fairies and of genii, which have been depicted with so much vividness of fancy and liveliness of description in the *Thousand and One Nights*, seem to have been connected with the pursuit of the science of transmutation, and the production of the elixir of life. The speculative ideas of the Arabians were more or less adopted by their European disciples. The Rosicrucian philosophy, in which gnomes, sylphs, salamanders, and nymphs were the spiritual agents, supposed capable of being governed or enslaved by man, seems to have originated with the alchemists of this period; and Agrippa, Paracelsus, and their followers, above-mentioned, all professed to believe in supernatural powers, in an art above experiment, in a system of knowledge not derived from the senses. It

would be a tedious and useless task, to describe all the absurdities in the opinions and practices of this school. Paracelsus alone deserves particular notice, from the circumstance of his being the first public lecturer on chemistry in Europe, and from the more important circumstance of his application of mercurial preparations to the cure of diseases. The magistrates of Basle established a professor's chair for their countryman, but he soon quitted an occupation in which regularity was necessary, and spent his days in wandering from place to place, searching for, and revealing secrets. He pretended to confer immortality, by his medicines, and yet died at the age of 49, at Saltsburg, in the year 1541.*

The enthusiasm of this man almost supplied his want of genius. He formed a number of new preparations of the metals, which were studied and applied by his disciples; his exaggerated censure of the methods of the ancients, and of the systems of his day, had an effect in diminishing their popularity; one error was expelled by another; and it is a great step towards improvement, that men should know they have been in delusion.

Van Helmont, of Brussels, born in 1588,† was formed in the school of Alchemy, and his mind was tinctured with its prejudices: but his views concerning nature and the elements were distinguished by much more philosophical acuteness, and more sagacity, than those of any former writer. He is the first person who seems to have had any idea respecting elastic fluids, different from the air of the atmosphere; and he has distinctly mentioned three of these substances, to which he applied the term gases: namely, aqueous gas or steam, unctu-

* Dictionnaire Historique, par Moréri, tom. viii. p. 64.

† Ibid. tom. v. p. 570.

ous or inflammable gas, and gas from wood or carbonic acid gas. Van Helmont developed some accurate views respecting the permanent elasticity of air, and the operation of heat upon it; and a sketch of a curious instrument very similar to the differential thermometer, is to be found in his works.*

Van Helmont has used a term not so applicable or intelligible as gas, namely, Blas; which he supposed to be an influence derived from the heavenly bodies, of a most subtile and etherial nature; and on the idea of its operations in our terrestrial system, he has endeavoured to found the vindication of astrology.†

At this period there was no taste in the public mind to restrain vain imaginations. There were no severe critics to correct the wanderings of genius. The systems of logic, adopted in the schools, were founded rather upon the analogies of words than upon the relations of things; and they were more calculated to conceal error, than to discover truth. Till the revival of literature in Europe, there was no attempt at philosophical discussion in any of the sciences; the diffusion of letters gradually brought the opinions of men to the standard of nature and truth; failures in the experimental arts produced caution, and the detection of imposture created rational scepticism.

The delusions of Alchemy were exposed by Guibert, Gassendi, and Kepler. Libavius answered Guibert in a tone which demonstrated the weakness of his cause. This person, who died in 1616, was the last active experimentalist who believed that transmutation had actually been performed; and in the beginning of the

* Johan. Baptist. Van Helmont, *Opera Omnia*, 4to. p. 61. article Aer. [Vide pl. i. fig. 3.]

† Ibid. page 114.

17th century the processes of rational chemistry were pursued by a number of enlightened persons in different parts of Europe.

A metallurgical school had before this time been founded in Germany. George Agricola published, in 1542, his twelve books, *de Re Metallica*, or, on the methods of extracting and purifying the useful metals; and he was followed by Lazarus Erckern, Assay-Master General of the Empire of Germany, whose works, brought forward in 1574, contain a number of useful practices detailed in a simple and perspicuous manner.

Lord Bacon happily described the Alchemists as similar to those husbandmen who in searching for a treasure supposed to be hidden in their land, by turning up and pulverising the soil, rendered it fertile; in seeking for brilliant impossibilities, they sometimes discovered useful realities; and in speaking of the chemistry of his time, he says, a new philosophy has arisen from the furnaces, which has confounded all the reasonings of the ancients. This illustrious man himself pointed out many important objects of chemical inquiry; but he was a still greater benefactor to the science, by his development of the general system for improving natural knowledge. Till his time there had been no distinct views concerning the art of experiment and observation. Lord Bacon demonstrated how little could be effected by the unassisted human powers, and the weakness of the strongest intellect even without artificial resources. He directed the attention of inquirers to instruments for assisting the senses, and for examining bodies under new relations. He taught that Man was but the servant and interpreter of Nature; capable of discovering truth in no other way but by observing and imitating her operations; that facts were to be col-

lected and not speculations formed; and that the materials for the foundations of true systems of knowledge were to be discovered, not in the books of the ancients, not in metaphysical theories, not in the fancies of men, but in the visible and tangible external world.

Though Van Helmont had formed some just notions respecting the properties of air, yet his views were blended with obscure and vague speculations; and it is to the disciples of Galileo that the true knowledge of the mechanical qualities and agencies of elastic fluids is owing. After Torricelli and Pascal had shewn the pressure and weight of the atmosphere, the investigation of its effects in chemical operations became an obvious problem.

John Rey is generally quoted as the first person who shewed by experiments that air is fixed in bodies during calcination; but it appears from the work of this acute and learned man that he reasoned upon the processes of others, rather than upon his own observations.

He quotes Fachsius, Libavius, Cesalpin, and Cardan, as having ascertained the increase of weight of lead during its conversion into a calx,* and he mentions an experiment of Hammerus Poppius, who found that antimony calcined by a burning-glass, notwithstanding the loss of vapours, yet was heavier after the process.

Rey ridicules the various notions of the Alchemists on the cause of this phenomenon; and ascribes it to the union of air with the metal; he supposes that air is miscible with other bodies besides metals, and states distinctly that it may be expelled from water.

The observations of John Rey seem to have excited

* Sur la Recherche de la cause par laquelle Estain et le Plomb augmentent de poids quand on les calcine. A Bazas, 1630.

no attention amongst his contemporaries. The philosophical spirit was only beginning to animate chemistry, and the labourers in this science, occupied by their own peculiar processes, were little disposed to listen to the reasonings of an inquirer in general science; yet, though the most active of the forms of matter were neglected in the processes of the operative chemists of this day, and consequently no just views formed by them, still they discovered a number of important facts respecting the combinations and agencies of solid and fluid bodies. Glauber at Amsterdam, about 1640, made known several neutral salts, and several compounds of metallic and vegetable substances. Kunckel in Saxony and Sweden, pursued technical chemistry with very great success, and was the first person who made any philosophical experiments upon phosphorus, which was accidentally discovered by Brandt in 1669.* Barner in Poland, and Glaser in France, published elementary books on the science, and Borrichius in Denmark, Bohn at Leipzic, and Hoffman at Halle, pursued specific scientific investigations with much zeal and success; and Hoffman was the first person who attempted the philosophical analysis of mineral waters.

About the middle of this century likewise mathematical and physical investigations were pursued in every part of the civilized world with an enthusiasm before unknown. The new mode of improving knowledge by collecting facts, associated together a number of labourers in the same pursuit. It was felt that the whole of nature was yet to be investigated, that there were distinct subjects connected with utility and glory, sufficient to employ all inquirers, yet tending to the common end of promoting the progress of the human mind.

* Homberg, *Mem. Acad. Paris*, tom. x. page 58.

Learned bodies were formed in Italy, England, and France, for the purpose of the interchange of opinions, the combination of labour and division of expense in performing new experiments, and the accumulation and diffusion of knowledge.

The Academy del Cimento was established in 1651, under the patronage of the Duke of Tuscany; the Royal Society of London, in 1660; the Royal Academy of Sciences of Paris, in 1666. And a number of celebrated men, who have been the great luminaries of the different departments of science, were brought together or formed in these noble establishments. The ardour of scientific investigation was excited and kept alive by sympathy: taste was improved by discussion, and by a comparison of opinions. The conviction that useful discoveries would be appreciated and rewarded, was a constant stimulus to industry, and every field of inquiry was open for the free and unbiassed exercise of the powers of genius.

Boyle, Hooke, and Slare, were the principal early chemical investigators attached to the Royal Society of London. Homberg, Geoffroy, and the two Lemerys, a few years later, distinguished themselves in France.

Otto de Guericke of Magdeburgh invented the air-pump; and this instrument, improved by Boyle and Hooke, was made an important apparatus for investigating the properties of air. Boyle* and Hooke,† from their experiments, concluded that air was absolutely necessary to combustion and respiration, and that one part of it only was employed in these processes. And Hooke formed the sagacious conclusion, that this principle is the same as the substance fixed in nitre, and

* Boyle's Works, vol. iv. page 90.

† Hooke's Micrographia, page 45, 104, 105.

that combustion is a chemical process, the solution of the burning body in elastic fluid, or its union with this matter.

Mayow of Oxford, in 1674, published his treatises on the nitro-aerial spirit, in which he advanced opinions similar to those of Boyle and Hooke, and supported them by a number of original and curious experiments;* but his work, though marked by strong ingenuity, abounds in vague hypotheses. He attempted to apply the imperfect chemistry of his day to physiology; his failure was complete, but it was the failure of a man of genius.

Boyle was one of the most active experimenters, and certainly the greatest chemist of his age. He introduced the use of tests or re-agents, active substances for detecting the presence of other bodies: he overturned the ideas which at that time were prevalent, that the results of operations by fire were the real elements of things, and he ascertained a number of important facts respecting inflammable bodies, acids, alkalies, and the phenomena of combination; but neither he nor any of his contemporaries endeavoured to account for the changes of bodies by any fixed principles. The solutions of the phenomena were attempted either on rude mechanical notions, or by occult qualities, or peculiar subtile spirits or ethers supposed to exist in the different bodies.— And it is to the same great genius who developed the laws that regulate the motions of the heavenly bodies, that chemistry owes the first distinct philosophical elucidations of the powers which produce the changes and

* Tract. p. 28. He has particularly assigned the cause of the calcination of metals, "*Quippe vix concipi potest unde augmentum illud antimonii nisi a particulis nitro aereis ignisque inter calcinandum fixis procedat.*"

apparent transmutations of the substances belonging to the earth.

Sugar dissolves in water, alkalies unite with acids, metals dissolve in acids. Is not this, says Newton, on account of an attraction between their particles? Copper dissolved in aquafortis is thrown down by iron. Is not this because the particles of the iron have a stronger attraction for the particles of the acid, than those of copper: and do not different bodies attract each other with different degrees of force?*

A few years after Newton had brought forward these sagacious views, the elder Geoffroy endeavoured to ascertain the relative attractive powers of bodies for each other, and to arrange them in an order in which these forces, which he named affinities, were expressed.†

Chemistry had scarcely begun to assume the form of a science, when the attention of the most powerful minds were directed to other objects of research;—the same great man who bestowed on it its first accurate principles, in some measure impeded its immediate progress, by his more important discoveries in optics, mechanics, and astronomy.

These objects of the Newtonian philosophy were calculated by their grandeur, their simplicity, and their importance, to become the study of the men of most distinguished talents; the effect that they occasioned on the scientific mind may be compared to that which the new sensations of vision produce on the blind receiving sight;—they awakened the highest interest, the most enthusiastic admiration, and for nearly half a century, absorbed the attention of the most eminent philosophers of Britain and France.

* Newton's Works, 4to, tom. iv. page 242.

† Mémoires de l'Académie, 1718, page 256.

Germany still continued the great school of practical chemistry, and at this period it gained an ascendancy of no mean character over the rest of Europe in the philosophy of the science. Beccher, who was born at Spire in 1645, after having studied with minute attention, the operations of metallurgy, and the phenomena of the mineral kingdom, formed the bold idea of explaining the whole system of the earth by the mutual agency and changes of a few elements. And by supposing the existence of a vitrifiable, a metallic, and an inflammable earth, he attempted to account for the various productions of rocks, crystalline bodies, and metallic veins, assuming a continued interchange of principles between the atmosphere, the ocean, and the solid surface of the globe, and considering the operations of nature as all capable of being imitated by art.

The *Physica subterranea*, and the *Oedipus chemicus* of this author, are very extraordinary productions. They display the efforts of a vigorous mind, the conceptions of a most fertile imagination, but the conclusions are too rapidly formed; there is a want of logical precision in his reasonings; the objects he attempted were grand, but his means of execution comparatively feeble. He endeavoured to raise a perfect and lasting edifice upon foundations too weak, from materials too scanty, and not sufficiently solid; and the work, though magnificent in design, was rude, unfinished, and feeble, and rapidly fell into decay.

Beccher added very little to the collection of chemical experiments, but he improved the instruments of research, simplified the manipulations, and by the novelty and boldness of his speculations, excited inquiry amongst his disciples.

His most distinguished follower was George Ernest

Stahl, born in 1660, who soon attained a reputation superior to that of his master, and developed doctrines which for nearly a century constituted the theory of chemistry of the whole of Europe.

Albertus Magnus had advanced the idea that the metals were earthy substances impregnated with a certain inflammable principle. Beccher supported the idea of this principle, not only as the cause of metalization, but likewise of combustibility; and Stahl endeavoured, by a number of ingenious and elaborate experiments, to prove the existence of phlogiston, as it was called, and to explain its agencies in the phenomena of nature and art.

Glauber, about fifty years before Stahl began his labours, had discovered the combination of fossil alkali and sulphuric acid, which still bears his name. And Stahl, in operating upon this body, thought he had discovered the proof, that the inflammability not only of metals, but likewise of all other substances, was owing to the same principle. Charcoal is entirely dissipated or consumed in combustion, therefore, says this philosopher, it must be phlogiston nearly pure; by heating charcoal with metallic earths, they become metals; therefore they are compounds of metallic earths and phlogiston: by heating Glauber's salt, which consists of sulphuric acid and fossil alkali, with charcoal, a compound of sulphur and alkali is obtained; therefore sulphur is an acid combined with phlogiston. Stahl entirely neglected the chemical influence of air on these phenomena; and though Boyle had proved that phosphorus and sulphur would not burn without air, and had stated that sulphur was contained in sulphuric acid, and not the acid in sulphur, yet the ideas of the Prussian school were received without controversy.

Similar opinions were adopted in France by Homberg and Geoffroy, who assumed them without reference to the views of the Prussian philosopher, and opposed them to the more correct and sagacious views of the English school of chemistry.

Though misled in his general notions, few men have done more than Stahl for the progress of chemical science. His processes were, many of them, of the most beautiful and satisfactory kind: he discovered a number of properties of the caustic alkalies and metallic calces, and the nature of sulphureous acid; he reasoned upon all the operations of chemistry in which gaseous bodies were not concerned, with admirable precision. He gave an axiomatic form to the science, banishing from it vague details, circumlocutions and enigmatic descriptions, in which even Beccher had too much indulged; he laboured in the spirit of the Baconian school, multiplying instances, and cautiously making inductions, and appealing in all cases to experiments which, though not of the most refined kind, were more perfect than any which preceded them.

Dr. Hales, about 1724, resumed the investigations commenced with so much success by Boyle, Hooke, and Mayow; and endeavoured to ascertain the chemical relations of air to other substances, and to ascertain by statical experiments the cases in nature, in which it is absorbed or emitted. He obtained a number of important and curious results; but, misled by the notion of one elementary principle constituting elastic matter, and modified in its properties by the effluvia of solid or fluid bodies, he formed few inferences connected with the refined philosophy of the subject: he disengaged, however, elastic fluids from a number of substances, and drew the conclusion that air was a

chemical element in many compound bodies, and that flame resulted from the action and re-action of aerial and sulphurous particles.*

In 1756 Dr. Black published his admirable researches on calcareous, magnesian, and alkaline substances, by which he proved the existence of a gaseous body, perfectly distinct from the air of the atmosphere. He shewed that quick-lime differed from marble and chalk by containing this substance, and that it was a weak acid, capable of being expelled from alkaline and earthy substances by strong acids.†

Ideas so new and important as those of the British philosopher, were not received without opposition; several German inquirers endeavoured to controvert them. Meyer attempted to shew that limestones became caustic, not by the emission of elastic matter, but by combining with a peculiar substance in the fire; but the loss of weight was perfectly inconsistent with this view: and Bergman at Upsal, Macbride in Ireland, Keir at Birmingham, and Cavendish in London, demonstrated the correctness of the opinions of Black; and a few years were sufficient to establish his theory upon immutable foundations.

The knowledge of one elastic fluid different from air, immediately led to the inquiry whether there might not be others. The processes of fermentation which had been observed by the ancient chemists, and those by which Hales had disengaged and collected elastic substances, were now regarded under a novel point of view; and the consequence was, that a number of new bodies, possessed of very extraordinary properties, were discovered.

* Hales' Statical Essays, 2d edit. 8vo. vol. i. page 315.

† Essays and Observations, Physical and Literary, vol. ii. page 159.

Mr. Cavendish, about 1765, invented an apparatus for examining elastic fluids confined by water, which has been since called the hydro-pneumatic apparatus. He discovered inflammable air, and described its properties; he ascertained the relative weights of fixed air, inflammable air, and common air, and made a number of beautiful and accurate experiments on the properties of these elastic substances.

Dr. Priestley, in 1771, entered the same interesting path of inquiry; and principally by repeating the processes of Hales, added a number of most important facts to this department of chemical philosophy. He discovered nitrous air, nitrous oxide, and dephlogisticated air; and by substituting mercury for water in the pneumatic apparatus, ascertained the existence of several aëriform substances, which are rapidly absorbable by water, muriatic acid air, sulphurous acid air, and ammonia.

Whilst a new branch of the science was making this rapid progress in Britain, the chemistry of solid and fluid substances was pursued with considerable zeal and success in France and Germany; and Macquer, Rouelle, Margraff, and Pott, added considerably to the knowledge of fossile bodies, and the properties of the metals. Bergman, in Sweden, developed refined ideas on the powers of chemical attraction, and reasoned in a happy spirit of generalization on many of the new phenomena of the science; and in the same country, Scheele, independently of Priestley, discovered several of the same aëriform substances: he ascertained the composition of the atmosphere; he brought to light fluoric acid, prussic acid, and the substance which has been improperly called oximuriatic gas.

Black, Cavendish, Priestley, and Scheele, were un-

doubtedly the greatest chemical discoverers of the eighteenth century; and their merits are distinct, peculiar, and of the most exalted kind. Black made a smaller number of original experiments than either of the other philosophers; but being the first labourer in this new department of the science, he had greater difficulties to overcome. His methods are distinguished for their simplicity; his reasonings are admirable for their precision; and his modest, clear, and unaffected manner, is well calculated to impress upon the mind a conviction of the accuracy of his processes, and the truth and candour of his narrations.

Cavendish was possessed of a minute knowledge of most of the departments of Natural Philosophy: he carried into his chemical researches a delicacy and precision, which have never been exceeded: possessing depth and extent of mathematical knowledge, he reasoned with the caution of a geometer upon the results of his experiments: and it may be said of him, what, perhaps, can scarcely be said of any other person, that whatever he accomplished, was perfect at the moment of its production. His processes were all of a finished nature; executed by the hand of a master, they required no correction; the accuracy and beauty of his earliest labours even, have remained unimpaired amidst the progress of discovery, and their merits have been illustrated by discussion, and exalted by time.

Dr. Priestley began his career of discovery without any general knowledge of chemistry, and with a very imperfect apparatus. His characteristics were ardent zeal and the most unwearied industry. He exposed all the substances he could procure to chemical agencies, and brought forward his results as they occurred, without attempting logical method or scientific arrangement. His hypotheses were usually founded upon a few loose

analogies; but he changed them with facility; and being framed without much effort, they were relinquished with little regret. He possessed in the highest degree ingenuousness and the love of truth. His manipulations, though never very refined, were always simple, and often ingenious. Chemistry owes to him some of her most important instruments of research, and many of her most useful combinations; and no single person ever discovered so many new and curious substances.

Scheele possessed in the highest degree the faculty of invention; all his labours were instituted with an object in view, and after happy or bold analogies. He owed little to fortune or accidental circumstances: born in an obscure situation, occupied in the duties of an irksome employment, nothing could damp the ardour of his mind or chill the fire of his genius: with very small means he accomplished very great things. No difficulties deterred him from submitting his ideas to the test of experiment. Occasionally misled in his views, in consequence of the imperfection of his apparatus, or the infant state of the inquiry, he never hesitated to give up his opinions the moment they were contradicted by facts. He was eminently endowed with that candour which is characteristic of great minds, and which induces them to rejoice as well in the detection of their own errors, as in the discovery of truth. His papers are admirable models of the manner in which experimental research ought to be pursued; and they contain details on some of the most important and brilliant phenomena of chemical philosophy.

The discovery of the gases, of a new class of bodies, more active than any others in most of the phenomena of nature and art, could not fail to modify the whole

theory of chemistry. The ancient doctrines were revised; new modifications of them were formed by some philosophers; whilst others discarded entirely all the former hypotheses, and endeavoured to establish new generalizations.

The idea of a peculiar principle of inflammability was so firmly established in the chemical schools, that even the knowledge of the composition of the atmosphere for a long while was not supposed to interfere with it; and the part of the atmosphere which is absorbed by bodies in burning, was conceived to owe its powers to its attraction for phlogiston.

All the modern chemists who made experiments upon combustion, found that bodies increased in weight by burning, and that there was no loss of ponderable matter. It was necessary therefore to suppose, contrary to the ideas of Stahl, that phlogiston was not emitted in combustion, but that it remained in the inflammable body after absorbing gaseous matter from the air. But what is phlogiston? was a question constantly agitated. Inflammable air had been obtained during the dissolution of certain metals, and during the distillation of a number of combustible bodies. This light and subtile matter, therefore, was fixed upon as the principle of inflammability; and Cavendish, Kirwan, Priestley, and Fontana, were the illustrious advocates of this very ingenious hypothesis.

In 1774, Bayen* shewed that mercury converted into a calx or earth, by the absorption of air, could be revived without the addition of any inflammable substance; and hence he concluded, that there was no necessity for supposing the existence of any peculiar principle of inflammability, in accounting for the calci-

* Journal de Physique, 1774, p. 288.

nation of metals. The subject, nearly about the same time, was taken up by Lavoisier, who had been for some time engaged in repeating the experiments of the British philosophers. Bayen formed no opinion respecting the nature of the air produced from the calx of mercury. Lavoisier, in 1775, shewed that it was an air which supported flame and respiration better than common air, which he afterwards named oxygen; the same substance that Priestley and Scheele had procured from other metallic substances the year before, and had particularly described.*

Lavoisier discovered that the same air is produced during the revivification of metallic calces by charcoal, as that which is emitted during the calcination of limestone; hence he concluded, that this elastic fluid is composed of oxygen and charcoal; and from his experiments on nitrous acid and oil of vitriol, he concluded that this gas entered into the composition of these substances.

Dr. Black had demonstrated by a series of beautiful experiments, that when gases are condensed, or when fluids are converted into solids, heat is produced. In combustion gaseous matter usually assumes the solid or the fluid form. Oxygen gas, said Lavoisier, seems to be compounded of the matter of heat, and a basis. In the act of burning, this basis is united to the combustibile body, and the heat is evolved. There is no necessity, said this acute philosopher, to suppose any phlogiston, any peculiar principle of inflammability; for all the phenomena may be accounted for without this imaginary existence.

* In the *Journal de Physique* for 1789, Preliminary Discourse, De la Metherie has given an admirable view of the progress of the investigations concerning the gases. See p. 24, &c.

Lavoisier must be regarded as one of the most sagacious of the chemical philosophers of the last century ; indeed, except Cavendish, there is no other inquirer who can be compared to him for precision of logic, extent of view, and sagacity of induction. His discoveries were few, but he reasoned with extraordinary correctness upon the labours of others. He introduced weight and measure, and strict accuracy of manipulation into all chemical processes. His mind was unbiassed by prejudice ; his combinations were of the most philosophical nature : and in his investigations upon ponderable substances, he has entered the true path of experiment with cautious steps, following just analogies, and measuring hypotheses by their simple relations to facts.

The doctrine of Lavoisier, soon after it was framed, received some important confirmations from the two grand discoveries of Mr. Cavendish, respecting the composition of water, and nitric acid ; and the elaborate and beautiful investigations of Berthollet respecting the nature of ammonia ; in which phenomena, before anomalous, were shewn to depend upon combinations of aëriform matter.

The notion of phlogiston was, however, defended for nearly twenty years, by some philosophers in Germany, Sweden, Britain, and Ireland. Mr. Cavendish in 1784, drew a parallel between the hypothesis, that all inflammable bodies contain inflammable air, and the doctrine in which they are considered as simple substances, in a paper equally remarkable for the precision of the views displayed in it, and for the accuracy and minuteness of the experiments it contains. To this great man, the assumption of M. Lavoisier, of the matter of heat, appeared more hypo-

thetical than that of a principle of inflammability. He states, that the phenomena may be explained on either doctrine; but he prefers the earlier view, as accounting, in a happier manner, for some of the operations of nature.

De Morveau, Berthollet, and Fourcroy, in France, and William Higgins and Dr. Hope, in Britain, were the first advocates for the antiphlogistic chemistry. Sooner or later, that doctrine which is an expression of facts, must prevail over that which is an expression of opinion. The most important part of the theory of Lavoisier was merely an arrangement of the facts relating to the combinations of oxygen: the principle of reasoning which the French school professed to adopt was, that every body which was not yet decomposed, should be considered as simple; and though mistakes were made with respect to the results of experiments on the nature of bodies, yet this logical and truly philosophical principle was not violated; and the systematic manner in which it was enforced, was of the greatest use in promoting the progress of the science.

Till 1786, there had been no attempt to reform the nomenclature of chemistry; the names applied by discoverers to the substances which they made known, were still employed. Some of these names, which originated amongst the alchemists, were of the most barbarous kind; few of them were sufficiently definite or precise, and most of them were founded upon loose analogies, or upon false theoretical views.

It was felt by many philosophers, particularly by the illustrious Bergman, that an improvement in chemical nomenclature was necessary, and in 1787, Messrs. Lavoisier, Morveau, Berthollet, and Fourcroy, presented to the world a plan for an almost entire change in the

denomination of chemical substances, founded upon the idea of calling simple bodies by some names characteristic of their most striking qualities, and of naming compound bodies from the elements which composed them.

The new nomenclature was speedily adopted in France; under some modifications it was received in Germany; and after much discussion and opposition, it became the language of a new and rising generation of chemists in England. It materially assisted the diffusion of the antiphlogistic doctrine, and even facilitated the general acquisition of the science; and many of its details were contrived with much address, and were worthy of its celebrated authors: but a very slight reference to the philosophical principles of language will evince that its foundations were imperfect, and that the plan adopted was not calculated for a progressive branch of knowledge.

Simplicity and precision ought to be the characteristics of a scientific nomenclature: words should signify things, or the analogies of things, and not opinions. If all the elements were certainly known, the principle adopted by Lavoisier would have possessed an admirable application; but a substance in one age supposed to be simple, in another is proved to be compound; and *vice versa*. A theoretical nomenclature is liable to continued alterations; *oxigenated muriatic acid* is as improper a name as *dephlogisticated marine acid*. Every school believes itself in the right; and if every school assumes to itself the liberty of altering the names of chemical substances, in consequence of new ideas of their composition, or decomposition, there can be no permanency in the language of the science, it must always be confused and uncertain. Bodies which are similar to each other should always be classed together;

and there is a presumption that their composition is analogous. Metals, earths, alkalies, are appropriate names for the bodies they represent, and independent of all speculative views; whereas oxides, sulphurets, and muriates, are terms founded upon opinions of the composition of bodies, some of which have been already found erroneous. The least dangerous mode of giving a systematic form to a language, seems to be, to signify the analogies of substances by some common sign affixed to the beginning or the termination of the word. Thus, as the metals have been distinguished by a termination in *um*, as *aurum*, so their calciform or oxidated state, might have been denoted by a termination in *a*, as *aura*; and no progress, however great, in the science, could render it necessary that such a mode of appellation should be changed. Moreover, the principle of a composite nomenclature must always be very limited. It is scarcely possible to represent bodies consisting of five or six elements in this way, and yet it is in such difficult cases that a name implying a chemical truth would be most useful.

The new doctrines of chemistry, before 1795, were embraced by almost all the active experimental inquirers in Europe; and the adoption of a precise mode of reasoning, and more refined forms of experiment, led not only to the discovery of new substances, but likewise to a more accurate acquaintance with the properties and composition of bodies that had long been known.

New investigations were instituted with respect to all the productions of nature, and the immense variety of substances in the mineral, vegetable, and animal kingdom, submitted to chemical experiments.

The analysis of mineral bodies first attempted by

Pott in experiments principally on their igneous fusion, and afterwards refined by the application of acid and alkaline menstrea, by Margraff, Bergman, Bayen, and Achard, received still greater improvements from the labours of Klaproth, Vauquelin, and Hatchett. Hoffman, in the beginning of the 18th century, pointed out magnesia as a peculiar substance.* Margraaf, about fifty years later,† distinguished accurately between the silicious, calcareous, and aluminous earths. Scheele, in 1774, discovered barytes. Klaproth,‡ in 1788, made known zircon. Dr. Hope,§ strontites in 1791. Gadolin, ittria|| in 1794; and Vauquelin, glucine in 1798.

Seven metals only had been accurately known to the ancients, gold, silver, mercury, copper, lead, tin, and iron. Zinc, bismuth, arsenic, and antimony, though mentioned by the Greek and Roman authors, yet were employed only in certain combinations, and the production of them in the form of reguli or pure metals, was owing to the Alchemists.

Cobalt had been used to tinge glass in Saxony in the sixteenth century; but the metal was unknown till the time of Brandt, and this celebrated Swedish chemist discovered it in 1733. Nickel¶ was procured by Cronstedt in 1751. The properties of manganese, which was announced as a peculiar metal by Kaim** in 1770, were minutely investigated by Scheele and Bergman a few years after. Molybdic acid was discovered by

* Hoffman. Opera, tom. iv. p. 479.

† Opuscles, tom. ii. p. 137.

‡ Annales de Chimie, tom. i. p. 183.

§ Edinburgh Trans. vol. iv. p. 44.

|| Crell's Annals, 1796.

¶ Bergman Opuscles, tom. ii. p. 22.

** De Metallis dubiis, p. 48.

Scheele in 1778, and a metal procured from it by Hielm in 1782, the same year that tellurium was made known by Müller. Scheele discovered tungstic acid in 1781; and soon after a metal was extracted from it by Messrs. D'Elhuyars. Klaproth discovered uranium in 1789.* The first description of the properties of the oxide of titanium was given by Gregor in 1791.† Vauquelin made known chromium in 1797;‡ Hatchett, columbium in 1801;§ and shortly after, the same substance was noticed by Ekeberg, and named by him tantalum. Cerium was discovered in 1804, by Hisinger and Berzelius. Platina had been brought into Europe and examined by Lewis in 1749; and in 1803, Descotils, Fourcroy, and Vauquelin announced a new metallic substance in it; but the complete investigation of the properties of this extraordinary body was reserved for Messrs. Tennant and Wollaston, who in 1803 and 1804 discovered in it no less than four new metallic substances, besides the body which exists in it in the largest proportion, namely, iridium, osmium, palladium, and rhodium.

The attempts made to analyze vegetable substances previous to 1720, merely produced their resolution into the supposed elements of the chemists of those days, namely, salts, earths, phlegm, and sulphur. Boerhaave and Newmann attempted an examination by fluid menstrua, which was pursued with some success by Rouelle, Macquer and Lewis. Scheele, between 1770 and 1780, pointed out several new vegetable acids. Fourcroy, Vauquelin, Deyeux, Seguin, Proust, Jacquin, and

* Journal de Physique, 1789, p. 39.

† Annales de Chimie, xii. p. 147.

‡ Ibid. xxv. 21.

§ Phil. Trans. 1802.

Hermbstadt, between 1780 and 1790, in various interesting series of experiments, distinguished between different secondary elements of vegetable matter, particularly extract, tannin, gums, and resinous substances; and investigations of this kind have been pursued with great success by Hatchett, Pearson, Shroeder, Chenevix, Gehlen, Thomson, Thenard, Chevreul, Kind, Brande, Bostock, and Duncan. The chemistry of animal substances has received great elucidations from several of the same inquirers; and Berzelius has examined most of their results, and has added several new ones, in a comprehensive work expressly devoted to the subject, published in 1808.

That solid masses fell from above, connected with the appearance of meteors, had been advanced as early as 500 years before the Christian æra, by Anaxagoras; and the same idea had been brought forward in a vague manner by other inquirers amongst the Greeks and Romans, and was revived in modern times; but till 1802 it was regarded by the greater number of philosophers as a mere vulgar error, when Mr. Howard, by an accurate examination of the testimonies connected with events of this kind, and by a minute analysis of the substances said to have fallen in different parts of the globe, proved the authenticity of the circumstance, and shewed that these meteoric productions differed from any substances belonging to our earth; and since that period a number of these phenomena have occurred, and have been minutely recorded.

The philosophy of heat, the foundations of which were laid between 1757 and 1785, by Black, Wilcke, Crawford, Irvine, and Lavoisier, since that period has received some new and very important additions, from the inquiries of Pictet, Rumford, Herschel, Leslie, Dal-

ton, and Gay Lussac. The circumstances under which bodies absorb and communicate heat, have been minutely investigated; and the important discoveries of the different physical and chemical powers of the different solar rays, and of a property analogous to polarity in light, bear immediate relation to the most refined doctrines of corpuscular science, and promise to connect, by close analogies, the chemical and mechanical laws of matter.

A general view of the philosophy of chemistry was published under the name of Chemical Statics, in 1803, by the celebrated Berthollet. It is a work remarkable for the new views that it contains on the doctrines of attraction; views which are still objects of discussion, and which bear an immediate relation to some of the conclusions depending upon very recent discoveries.

At the time when the antiphlogistic theory was established, electricity had little or no relation to chemistry. The grand results of Franklin, respecting the cause of lightning, had led many philosophers to conjecture, that certain chemical changes in the atmosphere might be connected with electrical phenomena;—and electrical discharges had been employed by Cavendish, Priestley, and Vanmarum, for decomposing and igniting bodies; but it was not till the era of the wonderful discovery of Volta, in 1800, of a new electrical apparatus, that any great progress was made in chemical investigation by means of electrical combinations.

Nothing tends so much to the advancement of knowledge as the application of a new instrument. The native intellectual powers of men in different times, are not so much the causes of the different success of their labours, as the peculiar nature of the means and arti-

ficial resources in their possession. Independent of vessels of glass, there could have been no accurate manipulations in common chemistry; the air-pump was necessary for the investigation of the properties of gaseous matter; and without the Voltaic apparatus, there was no possibility of examining the relations of electrical polarities to chemical attractions.

By researches, the commencement of which is owing to Messrs. Nicholson and Carlisle, in 1800, which were continued by Cruickshank, Henry, Wollaston, Children, Pepys, Pfaff, Desormes, Biot, Thenard, Hissinger, and Berzelius, it appeared that various compound bodies were capable of decomposition by electricity; and experiments, which it was my good fortune to institute, proved that several substances which had never been separated into any other forms of matter in the common processes of experiment, were susceptible of analysis by electrical powers: in consequence of these circumstances, the fixed alkalies and several of the earths have been shewn to be metals combined with oxygen; various new agents have been furnished to chemistry, and many novel results obtained by their application, which at the same time that they have strengthened some of the doctrines of the school of Lavoisier, have overturned others, and have proved that the generalizations of the antiphlogistic philosophers were far from having anticipated the whole progress of discovery.

Certain bodies which attract each other chemically, and combine when their particles have freedom of motion, when brought into contact still preserving their aggregation, exhibit what may be called electrical polarities; and by certain combinations these polarities may be highly exalted; and in this case they become subservient to chemical decompositions; and by means

of electrical arrangements the constituent parts of bodies are separated in an uniform order, and in definite proportions.

Bodies combine with a force, which in many cases is correspondent to their power of exhibiting electrical polarity by contact; and heat, or heat and light, are produced in proportion to the energy of their combination. Vivid inflammation occurs in a number of cases in which gaseous matter is not fixed; and this phenomenon happens in various instances without the interference of free or combined oxygen.

Experiments made by Richter and Morveau had shewn that, when there is an interchange of elements between two neutral salts, there is never an excess of acid or basis; and the same law seems to apply generally to double decompositions. When one body combines with another in more than one proportion, the second proportion appears to be some multiple or divisor of the first; and this circumstance, observed and ingeniously illustrated by Mr. Dalton, led him to adopt the atomic hypothesis of chemical changes, which had been ably defended by Mr. Higgins in 1789, namely, that the chemical elements consist of certain indestructible particles which unite one and one, or one and two, or in some definite numbers.

Whether matter consists of indivisible corpuscles, or physical points endowed with attraction and repulsion, still the same conclusions may be formed concerning the powers by which they act, and the quantities in which they combine; and the powers seem capable of being measured by their electrical relations, and the quantities on which they act of being expressed by numbers.

In combination certain bodies form regular solids;

and all the varieties of crystalline aggregates have been resolved by the genius of Haiüy into a few primary forms. The laws of crystallization, of definite proportions, and of the electrical polarities of bodies, seem to be intimately related; and the complete illustration of their connection probably will constitute the mature age of chemistry.

To dwell more minutely upon the particular merits of the chemical philosophers of the present age, will be a grateful labour for some future historian of chemistry; but for a contemporary writer, it would be indelicate to assume the right of arbitrator, even where praise only can be bestowed. The just fame of those who have enlightened the science by new and accurate experiments, cannot fail to be universally acknowledged; and concerning the publication of novel facts there can be but one judgment; for facts are independent of fashion, taste, and caprice, and are subject to no code of criticism; they are more useful perhaps even when they contradict, than when they support received doctrines, for our theories are only imperfect approximations to the real knowledge of things; and in physical research, doubt is usually of excellent effect, for it is a principal motive for new labours, and tends continually to the development of truth.

The slight sketch that has been given of the progress of chemistry, has necessarily been limited to the philosophical details of discovery. To point out in historical order the manner in which the truths of the science have been applied to the arts of life, or the benefits derived by society from them, would occupy many volumes. From the first discovery of the production of metals from rude ores, to the knowledge of the bleaching liquor, chemistry has been continually subservient

to cultivation and improvement. In the manufacture of porcelain and glass, in the arts of dyeing and tanning, it has added to the elegancies, refinement, and comforts of life; in its application to medicine it has removed the most formidable of diseases; and in leading to the discovery of gunpowder, it has changed the institutions of society, and rendered war more independent of brutal strength, less personal, and less barbarous.

It is indeed a double source of interest in this science, that whilst it is connected with the grand operations of nature, it is likewise subservient to the common processes as well as the most refined arts of life. New laws cannot be discovered in it, without increasing our admiration of the beauty and order of the system of the universe; and no new substances can be made known which are not sooner or later subservient to some purpose of utility.

When the great progress made in chemistry within the last few years is considered, and the number of able labourers who are at present actively employed in cultivating the science, it is impossible not to augur well concerning its rapid advancement and future applications. The most important truths belonging to it are capable of extremely simple numerical expressions, which may be acquired with facility by students; and the apparatus for pursuing original researches is daily improved, the use of it rendered more easy, and the acquisition less expensive.

Complexity almost always belongs to the early epochs of every science; and the grandest results are usually obtained by the most simple means. A great part of the phenomena of chemistry may be already submitted to calculation; and there is great reason to believe, that at no very distant period the whole science will be cap-

able of elucidation by mathematical principles. The relations of the common metals to the bases of the alkalis and earths, and the gradations of resemblance between the bases of the earths and acids, point out as probable a similarity in the constitution of all inflammable bodies; and there are not wanting experiments, which render their possible decomposition far from a chimerical idea. It is contrary to the usual order of things, that events so harmonious as those of the system of the earth, should depend on such diversified agents, as are supposed to exist in our artificial arrangements; and there is reason to anticipate a great reduction in the number of the undecomposed bodies, and to expect that the analogies of nature will be found conformable to the refined operations of art. The more the phenomena of the universe are studied, the more distinct their connection appears, the more simple their causes, the more magnificent their design, and the more wonderful the wisdom and power of their Author.

PART I.

ON THE LAWS OF CHEMICAL CHANGES; ON UNDECOM-
POUNDED BODIES AND THEIR PRIMARY COMBINA-
TIONS.

DIVISION I.

ON THE POWERS AND PROPERTIES OF MATTER, AND THE
GENERAL LAWS OF CHEMICAL CHANGES.

I. *Preliminary Observations.*

1. THE forms and appearances of the beings and substances of the external world are almost infinitely various, and they are in a state of continued alteration: the whole surface of the earth even undergoes modifications: acted on by moisture and air, it affords the food of plants; an immense number of vegetable productions arise from apparently the same materials; these become the substance of animals; one species of animal matter is converted into another; the most perfect and beautiful of the forms of organised life ultimately decay, and are resolved into inorganic aggregates; and the same elementary substances, differently arranged, are contained in the inert soil, or bloom and emit fragrance in the flower, or become in animals the active organs of mind and intelligence. In artificial operations, changes of the same order occur; substances having the characters of earths are converted into metals; clays and sands are united so as to become por-

celain; earths and alkalies are combined into glass; acrid and corrosive matters are formed from tasteless substances;—colours are fixed upon stuffs, or changed, or made to disappear; and the productions of the mineral, vegetable, and animal kingdoms are converted into new forms, and made subservient to the purposes of civilized life.

2. To trace in detail these diversified and complicated phenomena, to arrange them and deduce general laws from their analogies, would be a labour to which even the longest life of the most industrious and sagacious individual might be devoted in vain. The student who has the advantage of referring to the knowledge accumulated by many philosophers in different times, may adopt much more simple methods of acquiring the science;—those of recurring to its general principles, so as to ascertain the powers and properties of matter, which are the causes of the phenomena of chemistry; and of applying these principles to the actions taking place between the various substances existing in nature, or produced by art; proceeding gradually, referring to observations, experiments, and distinct analogies, from the more simple to the more complicated changes, so as to understand the laws by which they are governed.

II. *Of the Forms of Matter.*

1. In the general views that may be taken of the properties of natural substances, certain relations appear which afford the means of arranging them in distinct classes, each of which is distinguished by certain sensible and obvious qualities.

2. The first class consists of *solids*, which compose the great known part of the globe. Solid bodies,

when in small masses, retain whatever mechanical form is given to them : their parts are separated with difficulty, and cannot readily be made to unite after separation ; some solid bodies yield to pressure, and do not recover their former figure when the compressing force is removed, and they are called non-elastic solids ; others that regain this form, are called elastic bodies. Solids differ in degrees of hardness, in colour, in degrees of opacity or transparency, in density, or in the weight afforded by equal volumes ; and when their forms are regular or crystallized, in the nature of these forms.

3. The second class consists of *fluids*, of which there are much fewer varieties. Fluids when in small masses assume the spherical form ; their parts possess freedom of motion ; they differ in degrees of density and tenacity, in colour and degrees of opacity or transparency. They are usually regarded as incompressible, at least a very great mechanical force is required to make them occupy a space perceptibly smaller.

4. *Elastic fluids* or *gases*, the third class, exist free in the atmosphere ; but they may be confined by solids, or by solids and fluids, and their properties examined. Their parts are highly moveable ; they are compressible and expansible, and their volumes are inversely as the weights compressing them. All known elastic fluids are transparent, and present only two or three varieties of colour ; they differ materially in density.

5. Besides these forms of matter which are easily submitted to experiment, and the parts of which may be considered as in a state of apparent rest, there are other forms of matter which are known to us only in their states of motion when acting upon our organs of sense, or upon other matter, and which are not susceptible of being confined. They have been sometimes

called *etherial substances*, which appears a more unexceptionable name than *imponderable substances*. It cannot be doubted that there is matter in motion in space, between the sun and the stars and our globe, though it is a subject of discussion whether successions of particles be emitted from these heavenly bodies, or motions communicated by them, to particles in their vicinity, and transmitted by successive impulses to other particles. *Etherial* matter differs either in its nature, or in its affections by motion; for it produces different effects; for instance, as radiant heat, and as different kinds of light.

6. The various forms of matter, and the changes of these forms, depend upon active powers, such as gravitation, cohesion, calorific repulsion or heat, chemical attraction, and electrical attraction, the laws of which it is necessary to study with attention.

III. *Gravitation.*

1. When a stone is thrown into the atmosphere it rapidly descends towards the surface of the earth. This is owing to *gravitation*. All the great bodies in the universe are urged towards each other by a similar force. A cannon ball sent from a piece of artillery describes a curve, and at last falls to the ground; were the impulse given to it by the gunpowder increased to a certain extent, and exerted in free space, it would continuously revolve round the earth, in consequence of the equilibrium between the two forces. The moon and the planets as Newton has demonstrated, are retained in their orbits by similar laws, and their harmonious and constant revolutions produced.

2. Bodies mutually gravitate towards each other;

but the smaller body proportionally more than the larger one: hence the power of gravity is said to be directly as the mass; it is in fact the measure of the mass or quantity of matter.

3. Gravitation acts inversely, as the square of the distance.

IV. *Cohesion.*

1. When two particles of quicksilver are brought into apparent contact they may be made to unite and form one globule: when a glass tube, having a very fine bore, is introduced into a vessel containing water, the water rises in the tube to a higher level than it occupied in the vessel: both these effects are said to be owing to *cohesion*, or *cohesive attraction*. It is the same force which preserves the forms of solids, and gives globularity to fluids, and is thus a prime cause of the permanency of the arrangements which compose the surface of the globe. It is usually said to act only at the surfaces of bodies, or by their immediate contact; but this does not seem to be the case. It certainly acts with much greater energy at small distances; but the spherical form of minute portions of fluid matter can only be produced by the attractions of all the parts of which they are composed, for each other; and most of these attractions must be exerted at sensible distances, so that for any thing we know to the contrary, gravitation and cohesion may be mere modifications of the same general power of attraction, in the one case acting at distances that can be easily measured, and in the other case, operating at distances which it is difficult to estimate.

2. Some philosophers have attempted to account for

attraction in general by supposing that there is a certain unknown matter always moving through the universe in right lines, by which bodies are impelled towards each other; but though the phenomena may be explained by such a supposition, it is as yet without proofs.

V. *Of Heat, or Calorific Repulsion.*

1. When a body which occasions the sensation of heat on our organs, is brought into contact with another body which has no such effect, the result of their mutual action is that the hot body contracts, and loses to a certain extent its power of communicating heat, and the other body expands, and in a degree acquires this power.

This law may be exemplified with respect to every form of ponderable matter. If a polished cylinder of tin, which accurately fits a ring, be heated so as to make water boil, it will no longer pass through the ring, and will be found enlarged in all its dimensions. If spirit of wine be heated in a glass vessel having a narrow tubulated neck, as it becomes capable of communicating the sensation of heat, it will be found to expand and to rise in the narrow neck; and if the body of the same vessel be filled with air, and it be inverted in water, its neck containing water, the air will rapidly expand, on the application of a heated body, and will cause the water to descend in the neck of the vessel.*

2. Different solids and fluids expand very differently when heated by the same means.

Glass is less expansible than any of the metals; a rod, the length of which at 32° is 1.000000, raised to the temperature of boiling water, expands so as to

* Plate I. fig. 1.

become 1·000861 parts; 1·000000 of platinum under similar circumstances expands so as to become 1·000884; gold, antimony, cast-iron, steel, iron, bismuth, copper, cast-brass, silver, tin, lead and zinc, expand in the following order:—Gold, 1·001466; antimony, 1·001083; cast-iron, 1·001111; steel, 1·001189; iron, 1·001220; bismuth, 1·001392; copper, 1·001718; cast-brass, 1·001875; silver, 1·001817; tin, 1·002483; lead, 1·002867; zinc, 1·002942; hammered zinc, 1·003011. The expansive power of liquids in general is greater than that of solids; alcohol appears to be more expansible than oils, and oils in general more expansible than water. Mercury by being heated from 32° to 212° expands in volume 0·01810. All the elastic fluids, or the different species of air that have been examined, as has been demonstrated by Messrs. Dalton and Gay Lussac, expand alike when heated to the same degree; 100 parts of each at the freezing point of water becoming about 137·5 at the boiling point.

It is evident that the density of bodies must be diminished by expansion; and in the case of fluids and gases, the parts of which are mobile, many important phenomena depend upon this circumstance. If heat be applied to fluids or to gases, the heated parts change their places and rise; and the colder parts descend and occupy their places. Currents are constantly produced in the ocean and in great bodies of water, in consequence of this effect. The heated water rises to the surface in the tropical climates, and flows towards colder ones, thus the warmth of the Gulf stream is felt a thousand miles from its source; and deep currents pass from the colder to the warmer parts of the sea; and the general tendency of these changes is to equalize the temperature of the globe.

In the atmosphere, heated air is constantly rising, and colder air rushes in to supply its place; and this event is the principal cause of winds; the air that flows from the poles towards the equator, in consequence of the rotation of the earth, has less motion than the atmosphere into which it passes, and occasions an easterly current; the air passing from the equator towards the poles having more motion, occasions a westerly current; and by these changes, the different parts of the atmosphere are mixed together; cold is subdued by heat, moist air from the sea is mixed with dry air from the land, and the great mass of elastic fluid surrounding the globe, preserved in a state fitted for the purposes of vegetable and animal life.

3. There are very few exceptions to the law of the *expansion* of bodies, at the time they become capable of communicating the sensation of heat; and these exceptions seem entirely to depend upon some chemical change in the constitution of bodies or on their crystalline arrangements. Thus clay contracts considerably in dimensions by a very intense heat, and on the measure of its contractions the pyrometer of Wedgwood is founded; but in this case the clay first gives off water, which was united to its parts, and afterwards these parts cohere together with more force, and from being in a state of loose aggregation become strongly united. Water expands a little before it congeals, and expands considerably during its conversion into ice; but in this case it assumes the crystalline form; and its parts, whilst they are arranging themselves to form regular solids, probably leave greater interstices than they occupied when at uniform distances in the fluid. Thus the same weight of matter will occupy much more space when arranged in a certain number of octahedrons, than when arranged in a similar number of cubes, or hexagonal

prisms. Certain saline solutions likewise that shoot into prismatic crystals, expand at the moment they become solid; and the case is the same with cast-iron, bismuth, and antimony.

The expansion of water during its conversion into ice, is shewn by the circumstance of ice swimming upon water; and if water in a deep vessel be examined at the time ice is forming, it will be found a little warmer at the bottom than at the top; and these circumstances are of great importance in the economy of nature. Water congeals only at the surface, where it is liable to be acted upon by the sun, and by warm currents of air which tend to restore it to the fluid state; and when water approaches near the point of freezing it begins to descend, so that no ice can form till the whole of the water has been cooled to the point where it possesses the greatest density; and in the deep parts of the sea and lakes, even in some of the northern latitudes, the duration of the long winter is insufficient to cool the water to the degree at which ice forms.

4. When equal quantities of the same matter differently heated are mixed together, as much as the one contracts, so much the other seems to expand. It is easy to prove this by shaking together 100 parts of mercury so hot as not to be touched without pain, and 100 parts in its common state, having previously measured the space they occupy; if the mixture is made in the tube that contained the hot mercury, there will be no sensible change of volume.

It is on the idea, that when heat or the power of repulsion is communicated from body to body, as much is gained by one body as is lost by the other, that thermometers have been framed, and the doctrines of temperature and capacity for heat founded.

5. The most common thermometer is a glass bulb, containing mercury, terminated by a glass tube, having a very narrow bore. The mercury is boiled to expel any air or moisture that might be attached to it; and at the moment it is in ebullition, the extremity of the tube being drawn to a fine point, is hermetically sealed by a spirit lamp. For the purpose of acquiring a scale, the bulb is first plunged into melting ice, and the place where the mercury stands is marked; the bulb is afterwards plunged into boiling water and the same operation repeated. On Fahrenheit's scale, this space is divided into 180 equal parts, and similar parts are taken above and below for extending the scale, and the freezing point of water is placed at 32° , and the boiling point at 212° . 1·8 degrees of Fahrenheit are equal to one degree of the centigrade thermometer, and 2·25 to one degree of Reaumur.

Other fluids besides mercury, such as alcohol, are sometimes used in thermometers, particularly for measuring low degrees when mercury freezes.

Air is employed in the differential thermometer, which consists of two bulbs filled with air, and connected by a capillary tube containing oil of vitriol: the heated body is brought in contact with one bulb, the air of which expands and drives the fluid towards the other bulb.*

6. *Temperature* is the power bodies possess of communicating or receiving heat, or the energy of repulsion; and the temperature of a body is said to be high or low with respect to another, in proportion as it occasions an expansion or contraction of its parts; and the thermometer is the common measure of temperature.

* Plate I. fig. 2. represents Mr. Leslie's differential thermometer. Fig. 3. is copied from Van Helmont. [Vide Introduction, p. 14.]

7. When equal volumes of different bodies of different temperatures are suffered to remain in contact till they are possessed of the same temperature, it is found that this temperature is not a mean one, as it would be in the case of equal volumes of the same body. Thus, if a pint of quicksilver at 100° , be mixed with a pint of water at 50° , the resulting temperature is not 75° , but about 70° : the mercury has lost 30° , whereas the water has gained only 20° . In the common language of chemical philosophers this difference is said to depend upon the different *capacities* of bodies for heat; and the *capacity* of a body is said to be greater or less, in proportion as its temperature is less or more raised by the addition, or diminished by the subtraction of equal quantities of the power of repulsion, or heat. Thus mercury is said to have a much less capacity for heat than water; and taking the facts above stated as data, and comparing the weights of the two bodies, which are as 13.3 to 1, their capacities will be to each other as about 19 to 1.

Tables of the relative capacities of bodies are given in the works of different authors. In referring to the various bodies which are the subjects of chemical changes, this property will be described amongst other properties. In general it appears that the substances most expansible by heat are those which have the greatest capacities; thus gases in general have greater capacities than fluids, and fluids than solids; but the exact ratio has not been yet determined.

8. Different bodies, it appears, have their temperatures differently raised by the addition, or diminished by the subtraction of equal quantities of heat, or the power of repulsion, and they are likewise affected by heat, or expanded with very different degrees of celerity.

If slender cylinders of silver, of glass, and of charcoal, of equal length and size, be held in the central part of the flame of a candle, the silver rapidly becomes heated throughout, and cannot be held in the hand; the heat is more slowly communicated through the glass, but the charcoal becomes red hot at the one extremity long before any heat is felt at the other extremity. These differences are said to depend upon the different powers of these bodies *for conducting* heat; thus the silver is said to be a better conductor than the glass, and the glass than the charcoal. In general those bodies that are the densest, and that have the least capacity for heat, are the best conductors; thus the metals conduct better than any other solids; gases are worse conductors than fluids, and fluids than solids: but there are exceptions with respect to this correspondence between conducting power and density, and a remarkable one, in the densest known body in nature, platina, which is the worst conductor amongst the metals.

Animal and vegetable substances in general, are very bad conductors; thus the hair and wool of animals, and the feathers of birds, are admirably fitted to protect them from the cold, and they inclose and retain air, which being a still worse conductor, enhances the effect. It was supposed by Count Rumford, that fluids and gases are perfect *nonconductors* of heat, and that their particles can be heated in no other way, except by coming in succession to the source of heat; but some conclusive experiments seem to render this opinion in its full extent untenable. In general, however, fluids and gases alter their places, from a change of specific gravity much more rapidly than they communicate or receive heat. This is illustrated by a very simple experiment; let an air thermometer be inverted in a vessel

of water, so that the extremity of the bulb is barely beneath the surface; let a little ether be poured upon the water so as to form a stratum about $\frac{1}{8}$ of an inch above the thermometer, and let the ether be inflamed;* however delicate the thermometer, the air in it will not expand; the ether boils violently, but a very long process of this kind is required to communicate any sensible heat to the water.

Unless the particles of gases and fluids had been capable of communicating heat to a certain extent, the upper strata of liquids would be almost the only permanently heated parts; and heat would be constantly accumulating on the surface of extensive seas. Our lower atmosphere likewise would be intensely cold during the absence of the sun:—but by the relations between the conducting power and the mobility of fluids and gases; the changes of temperature of air and water are made progressive and equable, and adapted to a habitable globe. As heat is propagated very slowly through gaseous bodies, so they communicate it very slowly to other bodies, a circumstance that might be expected from the small quantity of matter they contain, when compared to other substances. The heat of metals at the temperature of 120° is scarcely supportable; water scalds at 150° ; but air may be heated to 240° without being painful to our organs of sensation, and a temperature near this was experienced for some minutes, by Sir Joseph Banks, Sir Charles Blagden, and Dr. Fordyce, in a room artificially heated.

The power in air of abstracting heat is likewise comparatively very small; in the high northern latitudes a cold has been experienced without injury, in which mercury froze; and if in this state of the atmosphere, metallic substances, of the same temperature, were

* See Plate I. fig. 4.

touched, a sensation like that of burning was experienced, and the part blistered.

9. Heat, or the power of repulsion may be considered as the *antagonist* power to the attraction of cohesion, the one tending to separate, the other to unite the parts of bodies; and the forms of bodies depend upon their respective agencies. In solids the attractive force predominates over the repulsive; in fluids, and in elastic fluids they may be regarded as in different states of equilibrium; and in ethereal substances the repulsive must be considered as predominating over, and destroying the attractive force.

All the different substances in nature, under certain circumstances, are probably capable of assuming all these forms; thus solids, by a certain increase of temperature, become fluids, and fluids gases; and *vice versa*, by a diminution of temperature gases become fluids, and fluids solids.

Instances of the fusion of solids by heat are too familiar to require any particular notice; when water becomes steam by boiling, it is merely the conversion of a fluid into an elastic fluid; and a simple instance of this circumstance may be given in the ebullition of ether. Let a little ether be introduced into a small glass retort filled with water, and inverted in water; the ether will swim above the water, in the upper part of the retort; let a heated bar of metal* be held near the part of the retort containing the ether, as the heat is communicated, globules will be seen to rise, and in a very short time elastic fluid will be formed, in such quantities, as to expel the water from the vessel: on suffering the glass to cool, the elastic matter will be condensed, and will become again fluid.

* Plate I. fig. 5.

✓If a globule of mercury be held in a spoon of platina, over the flame of a lamp, it will be vividly agitated, and will rapidly diminish. This is owing to its becoming elastic, and flying off in gas: and by a very low temperature, which may be artificially produced by mixing together very cold snow and a salt called muriate of lime, mercury may be congealed into the solid form.

Different bodies change their states at very different temperatures. Thus mercury, which is a solid at about 40° below Fahrenheit, boils at about 660° ; sulphur, which becomes fluid at 218° , boils at 570° ; ether boils at 98° . The temperatures at which the common metals become gaseous, are generally very high, and most of them incapable of being produced by common means. Iron, manganese, platina, and some other metals, which can scarcely be fused in the best furnaces, are readily melted by electricity; and by the Voltaic apparatus a degree of heat is attained, in which platina not only fuses with readiness, but seems even to evaporate.

With respect to the conversion of solids, fluids, or gases, into ethereal substances, the proofs are not of the same distinct nature as those belonging to their conversion into each other. When the temperature of a body is raised to a certain extent, it becomes luminous; and heated bodies not only affect other bodies by direct contact, but likewise exert an influence on them at a distance, which is ascribed to what is usually called radiant heat. One solution of this phenomenon is, that particles are thrown off from heated bodies with great velocity, which by acting on our organs, produce the sensations of heat or light, and that their motion, communicated to the particles of other bodies, has the power of expanding them; thus if heat, or the force of repulsion, be so increased in an elastic fluid, as to over-

come the force of cohesion and gravitation, these particles would move in right lines through free space ; and we know of no other effects they could produce than those of heat and light. It is perhaps in favour of this opinion that all the different elastic fluids expand equally, when their temperatures are equally raised ; and from observations made on the eclipses of Jupiter's satellites, and from other phenomena presented by the heavenly bodies, it appears that the motions of light are equable.

It may, however, be said, that the radiant matters emitted by bodies in ignition, are specific substances, and that common matter is not susceptible of assuming this form ; or it may be contended, that the phenomena of radiation do, in fact, depend upon motions communicated to subtile matter every where existing in space.

10. The temperatures at which bodies change their states from fluids to solids, though in general definite, are influenced by a few circumstances, such as agitation and the pressure of the atmosphere. Water, kept perfectly at rest, may sometimes be cooled to 22° , without congelation ; but if, at a temperature below 32° , it be agitated, ice instantly forms. A saturated solution of Glauber's salt, introduced whilst warm into a bottle, from which the pressure of the atmosphere is excluded, remains liquid after cooling, but if the atmosphere be suffered to act upon it, it instantly crystallizes.* The boiling point of fluids is still less fixed, than the point of fusion of solids, and is immediately dependent upon pressure. Thus ether will boil readily at the freezing point of water, in the exhausted receiver of an air-pump ;

* [Even if atmospheric air be admitted, provided it be cooled slowly at rest, it will remain liquid for many hours ; if agitated, a portion of the salt crystallizes.]

and it appears from the researches of Professor Robison, that in a vacuum, all liquids boil about 145° lower, than in the open air. Under pressure, liquids may be heated to a high degree; water in a Papin's digester, may have its temperature raised to 300° , but at the moment the pressure is removed elastic matter is disengaged with great violence.

11. A peculiar distinction has been made by some authors between permanent elastic fluids, and elastic fluids which are condensible by pressure or cold; but these substances differ only in the degree of the point of vaporization; and steam at 500 degrees of Fahrenheit, there is every reason to believe, would be equally incondensable with air at a range of temperature such as we can command below our common temperatures; and some gases that are permanent under all common circumstances, as ammonia, are condensible by intense cold aided by pressure.

12. All bodies that boil at moderate temperatures, seem to evaporate, so as to produce a certain quantity of elastic matter in the common state of the atmosphere; and this quantity is greater in proportion as the temperature is high. According to Mr. Dalton, the force of vapour increases in geometrical progression to the temperature, but the ratio differs in different fluids. It is certain that as the temperature approaches near the point of ebullition, in liquids, the strength of the vapour, *i. e.* the quantity that would rise in free space, rapidly increases.

In hot, dry weather, it is obvious that there must be much more vapour in the atmosphere, than in cold wet weather; and the largest quantity exists in summer and in the tropical climates, when moisture is most needed for the purposes of life; and it appears to be the

aqueous vapour in the atmosphere, which, when condensed by the mixture of cold with hot air, or by other agencies occasioning a change of its temperature, is the cause of dew, mists, rain, and ultimately of springs, and rivers.

13. When solids are converted into fluids, or fluids into gases, there is always a loss of heat of temperature, and *vice versa*, when gases are converted into fluids, or fluids into solids, there is an increase of heat of temperature, and in this case it is said that *latent heat* is absorbed or given out. Thus if equal weights of snow at 32° and of water at 172° be mixed together, the whole of the snow is melted, but the temperature of the mixture is found to be 32° , so that 140 degrees of heat are lost. Again, if water be heated in a Papin's digester to 300 degrees, and the valve be raised, a quantity of steam instantly rises, which has the temperature of 212° , and the temperature of the water in the digester is found to be the same, so that a great quantity of heat of temperature is lost in converting the water into steam.

If, when the air is at 20° , a quantity of water be exposed to it in a tall glass, the water gradually cools down to 22° , without freezing, but if it be shaken, so as to be converted into ice, the temperature of the ice is found to be at 32° , so that the degree of heat is raised during the act of freezing.

If one part of steam or aqueous gas, at 212° , be mixed with six parts by weight of water at 62° , the whole of the steam will be condensed, and the temperature of the fluid will be about 212° , so that there is an immense increase of the heat of temperature, and 900° may be considered as taken from the steam, and as added to the water.

All the phenomena of these changes may be referred

to a simple general law, of which Dr. Black was the discoverer, and which has been most ably illustrated by the researches of Wilke, Watt, Irvine, and Crawford, namely, "that whenever a body changes its form, its relations to temperature are likewise changed, either increased or diminished;" and many important operations, both artificial and natural, depend upon this law. The knowledge of it, for instance, led Mr. Watt to make his great improvement of the steam engine, by which the steam is condensed out of the cylinder in which its force is efficient, and fresh gaseous matter introduced without any chance of a loss of its elasticity.

One of the most perfect modes of heating large rooms, and of procuring a uniform temperature for the purposes of manufacture, is by the condensation of steam. By the cold produced in consequence of the evaporation of water in hot climates, congelation is effected; and in the nights in Bengal, when the temperature is not below 50° , by the exposure of water in earthenware pans upon moistened bamboos, thin cakes of ice are formed, which are heaped together and preserved under ground by being kept in contact with bad conductors of heat.* The cold produced by evaporation, is likewise the cause of the formation of ice in Mr. Leslie's elegant experiment, in which sulphuric acid is placed in a vessel upon the plate of an air-pump, and water in another vessel raised above it; the surfaces both of the acid and the water being considerable. When an exhaustion is made, the sulphuric acid rapidly absorbs the vapour rising from the water; fresh vapour

* [In this instance the formation of ice seems to depend on radiation of heat rather than on evaporation,—on the same cause which produces dew, which is a common attendant. Vide Wells's Essay on Dew, first published in 1814.]

is immediately formed, and in a few minutes, if the circumstances are favourable, spicula of ice are seen to form on the surface of the water.

When aqueous vapour is condensed into fluid in the atmosphere, heat is produced; and the formation of rain, hail, and snow, tends to mitigate the severity of the winter. In the summer, evaporation is constantly tending to cool the surface. The melting of the polar ice moderates the heat that would arise in the northern regions from the constant presence of the sun during the polar summer. And the evolution of heat during the congelation of water, prevents too great a degree of cold, and renders the transitions of temperature more slow and gradual.

14. When the forms of bodies are changed by mechanical means, or when mechanical forces are made to act upon them, there is usually a change of temperature. A piece of caoutchouc extended and suffered to contract rapidly by mechanical means, becomes hot; a nail is easily made red hot by a few well directed blows of the hammer; and by the friction of solids, considerable increase of temperature is produced; thus the axle trees of carriages sometimes inflame.

By strong pressure, fluids even are made luminous, as has been lately shewn by M. Desaignes.

When an elastic fluid is compressed by mechanical means, its temperature is raised, and when the compressing forces are great and rapidly applied, the effect is such as to cause the combustion of bodies. A machine for setting fire to tinder of the agaric, by the compression of air, has been for some time in use.

When air is made to expand by removing compressing forces, a diminution of temperature is occasioned. Thus the mercury in the thermometer sinks at the time of

the rarefaction of air, by exhausting the receiver of an air pump.

In the common language of chemistry, it may be said that the capacity of elastic fluids for heat is diminished by compression, and increased by rarefaction; and it is probable that when the volumes of elastic fluids are changed by change of temperature, there is likewise a change of capacity; and on these ideas, it is easy to account for the correspondence between the diminution of the temperature of the atmosphere and its height; for if it be conceived that the capacity of air rarefied by heat, increases as it ascends, the heat of temperature which was the cause of its ascent, must, at a certain elevation, become heat of capacity: and the higher and more rarefied the air, the more it is removed from the source of heat, and the greater its power of diminishing temperature.

A very curious phenomenon is produced during the action of the fountain of Hiero, at Schemnitz in Hungary; the air in the machine is compressed by a column of water, 260 feet high, and when a stop-cock is opened so as to suffer air to escape, its sudden rarefaction produces a degree of cold which not only precipitates aqueous vapour, but causes it to congeal in a shower of snow, and the pipe from which the air issues, becomes covered with icicles. Dr. Darwin has ingeniously explained the production of snow on the tops of the highest mountains by the precipitation of vapour from the rarefied air which ascends from plains and valleys. The Andes, placed almost under the line, rise in the midst of burning sands; about the middle height is a pleasant and mild climate; the summits are covered with unchanging snows: and these ranges of temperature are always distinct; the hot winds from below, if

they ascend, become cooled in consequence of expansion, and the cold air, if by any force of the blast, it is driven downwards, is condensed, and rendered warmer as it descends.

It seems probable that the capacity of solids and fluids is increased by expansion, and diminished by condensation; and if this be the case, the additions of equal quantities of heat will give smaller increments of temperature at high than at low degrees, which must to a certain extent render the thermometer inaccurate in the higher degrees, though probably only to a very small extent, of little importance as to all practical purposes; and this cause of inaccuracy appears to be counteracted by another, that fluids seem to be more expandible by heat in proportion as their temperature is higher.

15. In all chemical changes there is an alteration of temperature; and in most instances when gases become fluids, or fluids solids, there is an increase of temperature; and *vice versa*, there is usually a diminution of temperature when solids become fluids, or fluids solids. For instance, when the highly inflammable substance called phosphorus, the properties of which will be hereafter described, is burnt in the air, it is found to condense a particular part of the air, and a high temperature is produced during the process. When a solid amalgam of bismuth, and a solid amalgam of lead, are mixed together, they become fluid, and the thermometer sinks during the time of their action.

There are, however, a number of cases in which, though gaseous bodies, or fluids are formed from solids, an increase of temperature occurs: thus, in the explosion of gunpowder a large quantity of aeriform matter is disengaged, yet a violent heat is produced.

And there is an instance in which at the time of the separation of two species of gaseous matter from each other, which is connected with expansion, there is an increase of temperature ; thus, when a little of the gas which I have named Euchlorine, and which consists of the substance called by the French chemists oxymuriatic gas, and oxygen gas, is gently heated in a small glass tube over mercury, an explosion takes place, fire appears, and yet the two gases occupy a greater volume than before the explosion.* And in the new detonating compound discovered by M. Dulong, which consists of two kinds of gaseous matter condensed in the fluid form, a flash of light occurs at the moment of their separation by heat, with a violent explosion.

16. As attempts have been made to account for attraction, by the supposition of the existence of a peculiar matter, so *calorific repulsion* has been accounted for by supposing a subtile fluid, capable of combining with bodies, and of separating their parts from each other, which has been named the *matter of heat*, or *caloric*.

Many of the phenomena admit of a happy explanation on this idea, such as the cold produced during the conversion of solids into fluids or gases, and the increase of temperature connected with the condensation of gases and fluids ; but there are other facts which are not so easily reconciled to the opinion : such are the production of heat by friction and percussion ; and some of the chemical changes which have been just referred to. When the temperature of bodies is raised by friction, there seems to be no diminution of their

* [There are instances also in which solids are formed from fluids, and even gases without the production of heat.]

capacities, using the word in its common sense; and in many chemical changes connected with an increase of temperature, there appears to be likewise an increase of capacity. A piece of iron made red hot by hammering cannot be strongly heated a second time by the same means, unless it has been previously introduced into a fire. This fact has been explained by supposing that the fluid of heat has been pressed out of it, by the percussion, which is recovered in the fire; but this is a very rude mechanical idea: the arrangements of its parts are altered by hammering in this way, and it is rendered brittle. By a moderate degree of friction, as it would appear from Rumford's experiments, the same piece of metal may be kept hot for any length of time; so that if heat be pressed out, the quantity must be inexhaustible. When any body is cooled, it occupies a smaller volume than before: it is evident, therefore, that its parts must have approached towards each other: when the body is expanded by heat, it is equally evident that its parts must have separated from each other. The immediate cause of the phenomena of heat then, as Lavoisier long ago stated, is motion, and the laws of its communication are precisely the same as the laws of the communication of motion.

Since all matter may be made to fill a smaller volume by cooling, it is evident that the particles of matter must have space between them; and since every body can communicate the power of expansion to a body of a lower temperature, that is, can give an expansive motion to its particles, it is a probable inference that its own particles are possessed of motion; but as there is no change in the position of its parts as long as its temperature is uniform, the motion, if it exist, must be a vibratory or undulatory motion, or a motion of the par-

ticles round their axes, or a motion of particles round each other.

It seems possible to account for all the phenomena of heat, if it be supposed that in solids the particles are in a constant state of vibratory motion, the particles of the hottest bodies moving with the greatest velocity, and through the greatest space; that in fluids and elastic fluids, besides the vibratory motion, which must be conceived greatest in the last, the particles have a motion round their own axes, with different velocities, the particles of elastic fluids moving with the greatest quickness; and that in ethereal substances the particles move round their own axes, and separate from each other, penetrating in right lines through space. Temperature may be conceived to depend upon the velocities of the vibrations; increase of capacity on the motion being performed in greater space; and the diminution of temperature during the conversion of solids into fluids or gases, may be explained on the idea of the loss of vibratory motion, in consequence of the revolution of particles round their axes, at the moment when the body becomes fluid or aëriform, or from the loss of rapidity of vibration, in consequence of the motion of the particles through greater space.

If a specific fluid of heat be admitted, it must be supposed liable to most of the affections which the particles of common matter are assumed to possess, to account for the phenomena; such as losing its motion when combining with bodies, producing motion when transmitted from one body to another, and gaining projectile motion, when passing into free space: so that many hypotheses must be adopted to account for its mode of agency, which renders this view of the subject less simple than the other. Very delicate experiments

have been made to show that bodies when heated do not increase in weight. This, as far as it goes, is an evidence against a specific subtile elastic fluid producing the calorific expansion; but it cannot be considered as decisive, on account of the imperfection of our instruments; a cubical inch of inflammable air requires a good balance to ascertain that it has any sensible weight, and a substance bearing the same relation to this, that this bears to platinum, could not, perhaps, be weighed by any methods in our possession.

Some arguments have been raised in favour of the existence of a specific fluid of heat, from the circumstance of the communication of heat to bodies in exhausted receivers, and from the manner in which they are affected by this heat; but there are no means known in experimental science of producing a perfect vacuum; even the best Torricellian vacuum must contain elastic matter. The great capacity of such highly rarefied matter is an obstacle to the indication of temperature; but supposing a communication of heat, the laws must be analogous to those of heat communicated to common air. If a long *cylinder* of metal, placed perpendicularly, be heated in the middle, the warmest part will be above, from the ascent of heated particles of the elastic medium; but if a *sphere* be heated in the middle, the hottest portion will be below, as the heated elastic matter must remain longer in contact with the inferior than with the superior portion.

The laws of the communication of heat, and the philosophy of its effects, are independent of this speculative question, which has been mentioned in this place merely with the view of guarding students against the adoption of the doctrine of a specific fluid of heat as a part of the philosophical principles of chemistry, and to

shew that as yet we have no decided evidence on the subject.

VI. *On Chemical Attraction, and the Laws of Combination and Decomposition.*

1. When olive oil and water are agitated together, they refuse to act upon each other, and separate according to the order of their densities, the oil swimming above the water. Oil and water will not mix intimately; they will not *combine*; and they are said to have no chemical *attraction* or *affinity* for each other. But if oil and soap lees, or solution of potassa in water, be mixed, the oil and the solution blend together, and a species of soap will be formed, which may be procured as a soft solid substance by evaporating a part of the water. This is an instance of *combination*; and solution of potassa and oil are said to *attract* each other chemically, or to have an *affinity* for each other.

2. Oil is almost insipid, but the solution of potassa is a caustic substance, which corrodes the skin, and has a strong taste. The body resulting from their union differs both from the oil and the alkali in taste, smell, colour, and in all its sensible qualities; and it is a general character of *chemical combination*, that it changes the sensible qualities of bodies.

Corrosive and pungent substances often become mild and tasteless by their union, and form what are called neutral compounds, as is the case with sulphuric acid and quick-lime, which form gypsum, or sulphate of lime.

Bodies possessed of little taste or smell often gain these qualities in a high degree by combination. Thus sulphur, when inflamed in oxygen or in common air,

dissolves and forms an elastic fluid of a most penetrating and disagreeable odour and peculiar flavour. The forms of bodies, or their densities, likewise usually alter; solids become fluids, and solids and fluids gases, and gases are often converted into fluids or solids. Thus sugar, or salt, or isinglass, dissolves in water. The consumption of charcoal in our fires depends upon its uniting with a part of the air, with which it forms an invisible elastic fluid; mercury is rendered solid by being heated with half its weight of tin, and a substance of this kind is used for silvering mirrors. The gas produced by the combustion of charcoal is condensed by another gas procured from quick-lime and sal ammoniac, when they are mixed over mercury; and the two invisible elastic fluids form a white saline solid.

3. Many substances may be made to unite by chemical affinity or attraction: thus the mineral alkali or soda, and oil of vitriol or sulphuric acid and water, will all unite together and form a peculiar salt. And the fossil alkali, sand and the glass of lead, when melted together, unite to form flint glass. And in like manner, porcelain is formed by heating together mixtures of different earths. In a number of the productions of nature likewise many substances are combined into one mass or compound. Thus many stones and gems are capable of being resolved into several elements; and in the vegetable and animal kingdom there are scarcely any compounds which do not contain more than two principles, and complexity of constitution seems uniformly connected with organization.

4. That chemical attraction may be exerted between bodies, it is necessary that they should be brought into apparent contact. Thus no body will act chemically upon another at any sensible distance.

5. A freedom of motion in the parts of bodies, or a want of cohesion, greatly assists combination ; and this circumstance is so marked, that it was formerly considered as a chemical axiom, which is still retained in some elementary books, that bodies cannot act chemically on each other unless one of them be fluid or aëriform. Such an extensive generalization is, however, incorrect ; thus crystalline muriate of lime and snow, both cooled to 0° Fahrenheit, and rubbed together, act upon each other and liquefy ; and crystals of oxalid acid and dry lime treated in the same manner readily combine. The hardest and the densest bodies, however, undergo chemical changes with the greatest difficulty. Thus the sapphire in its crystallized state, is not affected by boiling sulphuric acid ; but when in a fine powder, as alumine, it is easily dissolved. Minute division, or solution, or fusion is necessary in almost all chemical processes. In the chemical arts these circumstances are always attended to ; and in the phenomena of external nature, the commencement of chemical operations may in almost all cases be traced to the agencies of fluids or aëriform substances. Thus in the bosom of our rocks and mountains, where air and water are incapable of penetrating, all is permanent and still, without change or motion ; wherever water and air are capable of acting, decomposition slowly goes on ; and these agents gradually change the nature of the surface, render the soil fertile, and decompose and degrade the exterior of strata.

6. If equal weights of magnesia and of quick-lime, in fine powder intimately mixed, be thrown into twice their weight of dilute aqua fortis or nitric acid, and suffered to remain for some hours, it will be found by a minute examination, that a considerable part of the

lime has been dissolved, but all the magnesia will remain. Hence, it is said, that lime has a stronger *attraction* for nitric acid than magnesia has.

This is proved likewise, by another experiment of a different kind: it is easy to make a solution of magnesia, in nitric acid, by heating them together; and to make a solution of lime in water, by agitating some powdered quick-lime in distilled water. Let the solution of lime be poured into the solution of magnesia, a white powder or precipitate will separate, and gradually fall to the bottom of the vessel in which the mixture is made. This powder, when examined, is found to be magnesia, and, it is said, that magnesia is precipitated from nitric acid, in consequence of the stronger *attraction* of lime for that acid.

All bodies, that differ in their nature, combine with different degrees of force; and some very important chemical phenomena in the arts depend upon this circumstance. Thus the astringent or tanning substance, in the bark of trees, which is soluble in water, is attracted from water, by the prepared skins of animals, in consequence of their stronger affinity for it, and the skin, from being destructible by boiling water, and decomposable, becomes indestructible and permanent. In like manner, indigo, and other dyeing materials, are separated from their solutions, by vegetable or animal fibres, and new combinations of them effected; and a number of instances of the same kind might be brought forward.

7. Different bodies unite with different degrees of force; and hence, one body is capable of separating others, from certain of their combinations; and in consequence of the same circumstance, mutual decompositions of different compounds take place. This has been

called *double affinity*, or *complex chemical attraction*. Thus, if an aqueous neutral solution of lime and nitric acid, and a like solution of magnesia and sulphuric acid, be mixed together, the lime will quit the nitric acid, to unite to the sulphuric acid, and the magnesia will leave the sulphuric acid, to combine with the nitric acid. The combination of nitric acid and magnesia, will remain in solution; but the compound of lime and sulphuric acid, being only slightly soluble in water, will for the most part be precipitated, in the form of a white powder.

In many cases, decompositions, that cannot be produced by single attractions, may be produced by double affinities. Thus the elements of sulphate of baryta, or the combination of sulphuric acid, and the earth called baryta, are so firmly united, that no alkali, nor earth, will separate the acid from the baryta. Potassa, which has a very strong attraction for the acid, will not decompose it alone; but if potassa, combined with carbonic acid, be digested for some time, with powdered sulphate of baryta, there is a double decomposition; and combinations of sulphuric acid and potassa, and carbonic acid and baryta, are formed.

8. If one part of pure oxygen gas, and two parts of pure hydrogen gas, in volume, be mixed together, in a glass tube, over mercury, furnished with wires for passing the electrical spark through it, and they be inflamed by the electrical spark,* the gaseous matter will disappear, and water will result. If two parts of oxygen be employed, and two of hydrogen, one part of oxygen will remain; in whatever proportions they are mixed together, it is found that one of oxygen always condenses two of hydrogen. It is evident then, that

* See Plate I. fig. 6.

oxygen and hydrogen combine only in definite proportions, and that the water resulting is always the same in its constitution.

If a piece of well-burnt charcoal be introduced into a vessel, two-thirds filled with oxygen gas, over mercury; and the mercury be brought to the same level on the inside and on the outside of the jar, and the charcoal be inflamed by a burning-glass;* there will be at first, an expansion, but after the experiment is over, it will be found, that the volume of the gas has not perceptibly altered; and if the charcoal has been in sufficient quantity, the whole of the oxygen will be found converted into carbonic acid: now the densities of oxygen gas and carbonic acid gas, in whatever way they are formed, are always the same; and to each other, as 34 to 47 nearly. It is evident then, that carbonic acid must always contain the same weight of oxygen and charcoal. If there is twice as much oxygen in the vessel as is necessary for the consumption of the charcoal, half of it remains untouched; and if the charcoal is partly unconsumed, still the gas is the same in quality; it always contains by weight, 5·7 of charcoal, and 15 of oxygen.

If some pure lead be introduced into a retort of green glass, and the part of the retort containing the lead be heated to redness, it will become covered with a greyish powder, and by agitating the lead and continuing the heat, the metal will at length be entirely converted into a yellow glass; and if the air in the vessel be confined by a tight stopper adapted to the neck of the retort, it will be found that the part of it which supports combustion, the oxygen, is absorbed, and the lead is converted into yellow oxide by the absorption of oxygen, and the yellow oxide is always

* Plate II. fig. 7.

the same in composition, and 100 parts of lead always absorb 7·5 parts of oxygen to become the yellow oxide.

If water be poured upon pure dry quick lime, the lime will become hot, will absorb the water and fall into powder; the water unites chemically to the lime, and forms the solid substance called hydrate of lime; and this substance when dried at the temperature of boiling water is always the same in composition, always contains the same proportions of lime and water, and 1000 parts of lime, if pure, whatever be the nature of the marble or limestone from which it is made, absorbs rather more than 309 parts of water.

If 100 parts of sulphuric acid (oil of vitriol of commerce) be poured into a solution of baryta, whether in muriatic, acetic, or nitric acid, or water, the solid matter that falls down will always be the same in quantity, about 195 parts in weight. The sulphate of baryta is insoluble, and the same proportion of baryta always combines with the same proportion of sulphuric acid.

In experiments of this kind, the different manner of collecting and weighing the product, the different sizes of the vessels employed, and even the different states of the atmosphere, occasion slight differences in the results, but they are never considerable, and in taking the mean of several accurate experiments, estimations are obtained which cannot differ from the true proportions more than a few 10,000th parts, and which are sufficiently accurate as foundations for principles in the science.

The same circumstances of invariable proportions occur in all cases in which solids are formed by the combinations of fluids with each other, or with solids or gases, and in which gases are formed from solids, or solids and fluids, or fluids from gases alone, or gases

and solids. In all these cases the elements of the same compound are constantly found in the same quantities.

In instances in which fluids mix with each other, or dissolve solids or gases, the solutions may be differently constituted in different cases, that is, the fluids may be mixed in different quantities, or the solids or gases dissolved in different proportions of the same fluid, yet there is always a limit to true dissolution. Thus, water will only dissolve a certain quantity of saline, or acid, or alkaline or gaseous matters, and it is possible that solutions containing smaller quantities *may be* mixtures of the saturated solution with water. If water be carefully poured upon a saturated aqueous solution of sugar, it will swim above it without exerting any action upon it, but by agitation the two fluids will mix together and form an uniform body. In such instances the force which produces their mixture appears the same as attraction of aggregation, at least it is connected with none of those changes of sensible properties which are dependent upon the exertion of true chemical attraction. When two quantities of water or of mercury mix together to form one mass, the force that produces the effect appears of the same kind as that which occasions the mixture of saturated solutions, or solutions containing definite proportions of solid or gaseous substances with water or other fluids; and the true test of chemical union is a change of density of the acting bodies, and when there is a change of volume certain proportions are always uniform even in the action of fluids on each other. Thus, 92 parts of sulphuric acid undergo their maximum of contraction by combining with 17 parts of water, and the substance so formed congeals by a moderate degree of cold.

If a solution of the caustic vegetable alkali be poured

into any neutral solution, that is, a solution which does not exhibit alkaline or acid properties, in which magnesia is combined with an acid, it always occasions a precipitate, and whatever be the nature of the acid, the same quantity of solution of potassa always separates the same quantity of magnesia.

The case is similar when one acid is expelled by another from a neutral combination, provided the new acid likewise forms a neutral combination. Thus, the same quantity of carbonic acid gas is always expelled by the same quantity of acetic acid, whether it be combined with lime, or magnesia, or strontia, or baryta. And the proof is easily gained by exposing the carbonates, or the compounds of carbonic acid and the earths, in little graduated tubes filled with mercury, inverted in mercury, and admitting to each the same quantity of acetic acid. If the experiment be carefully made, it is found that equal quantities of carbonic acid gas are produced in all the tubes.

The law is general, if one metal precipitates another from certain solutions, provided they are neutral, and remain neutral, the same quantity of the one metal always separates the same quantity of the other metal, provided there is no decomposition of the solvent; thus, when copper is dissolved in the acetic and sulphuric acids, the same quantity of iron precipitates from each solution the same quantity of copper, but when copper is dissolved in nitric acid there is a different result, because the iron decomposes the acid.

When one gas expels another from a compound by a simple phenomenon of decomposition, in consequence of its superior attraction for the other element or elements of the compound, the ratio of the proportion is always in like manner uniform, and as the quantities of

gases are easily determined by their volumes, experiments on decompositions of this kind afford apt illustrations of the principle. Thus, if the neutral carbonates be exposed to sulphureous acid gas over mercury, there is no change of volume, though carbonic acid gas is slowly produced; a volume of carbonic acid gas is expelled for a volume of sulphureous acid gas absorbed, and whatever carbonate be used the result is similar.*

9. It is a proof of this general law of proportion, that in the instances of double decomposition there is never an excess of any one of the elements, and the neutral

* That the proportions in compound gases are definite, has long been generally acknowledged; but Mr. Higgins is, I believe, the first person who conceived that when gases combined in more than one proportion, all the proportions of the same element were equal; and he founded this idea, which was made public in 1789, on the corpuscular hypothesis, that bodies combine particle with particle, or one with two, or three, or a greater number of particles. Mr. Dalton, about 1802, adopting a similar hypothesis, apparently without the knowledge of what Mr. Higgins had written, extended his views to compounds in general. Mr. Richter seems to have been the first person to shew that in the decomposition of neutral salts by double affinity, the neutral state is preserved; and likewise that, when a metallic salt is decomposed by a metal, all the oxygen and acid is transferred, and the metal only changed, and that the new solution is as neutral as the former one. It had been ascertained, by different experiments, that in certain cases when solids dissolved in gases, the volume is unchanged, and some instances of the combination of gases were known, in which the volumes bore simple ratios to each other, as in nitrous oxide, and water; but M. Gay Lussac is the first philosopher who attempted to generalize on the phenomena, and shew that in all cases where gases unite, it is always in simple ratios of volume, 1 to 1, or 1 to 2, or 1 to 3, and that the condensation, if any, is in a simple ratio. His very ingenious ideas on this subject, were made known towards the close of 1808. Berzelius, in a work published in 1810, has determined very correctly some of the definite proportions of several important compounds. See *Higgins's Comparative View. Dalton's New Chemical Philosophy. Richter Ueber die neuen gegenstande der Chemie. Mémoires d'Arcueil*, tom. ii. *Berzelius Annales de Chemie*, tom. lxxvii. *Thomson's System of Chemistry*, vol. iii.

point of the mixture is preserved. Thus when nitrate of baryta, which consists of nitric acid and baryta, is decomposed by sulphate of potassa, no acid, no alkaline substance is rendered free, but all the potassa unites to all the nitric acid, and all the baryta to all the sulphuric acid, and two new compounds are formed.

Bodies when they form the same compound always combine in the same proportion, but they may likewise combine in different proportions, and then they form other compounds which are equally definite and invariable. Thus if sulphuric acid be poured into a solution of potassa in small quantities, the solid substance formed is a neutral salt which has no action upon vegetable colours; but if solution of potassa in small quantities be poured into sulphuric acid, the substance formed is a strong acid salt, much more fusible than the other salt; and these two different salts always contain proportions invariable for each species. And if the gas called fluoboric gas be added in small quantities to ammoniacal gas over mercury, it will form a fluid compound absorbing three times its volume; but if ammonia in small quantities be added to fluo-boric acid, there is produced a solid compound, and it combines only with an equal volume of the alkaline gas.

10. In cases when one body combines with another in more than one proportion, there is always a relation between the quantities similar to that mentioned above, that is, the greater proportions are always twice or three times or some simple multiple of the lesser. Thus in the sulphate and the acid sulphate of potassa, the quantity of potassa remaining the same, there is exactly twice as much acid in the acid sulphate as in the neutral sulphate.

In the two combinations of carbonic acid with po-

tassa, as Dr. Wollaston has shewn by a very simple experiment, the quantity of carbonic acid in one is double that in the other. This is his experiment: let a given weight of the salt called carbonate of potassa be thrown into a tube over mercury, and diluted sulphuric acid sufficient to cover it be introduced into the tube, a certain volume of carbonic acid gas will be disengaged; let an equal weight of the salt be heated to redness when it becomes a subcarbonate, and let this subcarbonate be treated in the same way, it will be found to give off exactly half as much carbonic acid gas.

In the combinations of oxygen there are likewise numerous examples; thus there are two combinations of mercury with oxygen, the black and the red; and one appears to contain twice as much oxygen as the other. There are two known combinations of iron with oxygen, the black and the red oxide of iron, and the oxygen in the first being considered as 2, that in the second must be considered as 3, that is, 100 parts of iron take 29 parts of oxygen to become the black oxide, and 43·5* to become the red.

11. As in all well-known compounds, the proportions of the elements are in certain definite ratios to each other; it is evident, that these ratios may be expressed by numbers; and if one number be employed to denote the smallest quantity in which a body combines, all other quantities of the same body will be multiples of this number; and the smallest proportions in which the undecomposed bodies enter into union being known, the constitution of the compounds they form may be learnt, and the element which unites chemically in the

* These results I have obtained very nearly, namely, 28 and 43, and they differ very little from those of Mr. Hassenfratz, Dr. Thomson, and M. Berzelius.

smallest quantity being expressed by unity, all the other elements may be represented by the relations of their quantities to unity.

Hydrogen gas, or inflammable air, is the substance of which the smallest weights seem to enter into combination; and it appears to exist in no definite compound in less proportion than in water. The specific gravity of hydrogen is to that of oxygen as 1 to 15; and as 2 volumes of hydrogen to 1 of oxygen enter into the composition of water, the ratio of the hydrogen in water will be to the oxygen as 2 to 15; and it may be regarded as composed of two proportions of hydrogen and 1 of oxygen: and the number representing hydrogen will be 1, and that representing oxygen 15.

The weights of equal volumes of azote and oxygen are to each other nearly as 13 to 15; therefore supposing the number representing the proportion, in which azote combines, gained from the composition of nitrous oxide, which contains 2 volumes of azote to 1 of oxygen, it will be represented by 26; and nitrous oxide will consist of 1 proportion of azote equal to 26, and 1 proportion of oxygen, equal to 15. Nitrous gas will consist of 1 of azote and 2 of oxygen, 26 and 30. Nitrous acid gas of 1 of azote and 4 of oxygen, 26 and 60.

Ammonia, which is decomposed by electricity into 3 volumes of hydrogen and 1 volume of azote, will consist of 6 proportions of hydrogen and 1 proportion of azote, or 6 and 26.

The weight of chlorine or oxymuriatic gas, is to that of hydrogen nearly as 33.5 to 1; and muriatic acid gas consists of equal volumes of these gases, and therefore is composed of 33.5 of chlorine, and 1 of hydrogen: but 2 of chlorine may be made to combine with 1 of oxygen in volume; and double proportions of

this gas combine to form compounds, which when decomposed by water, afford compounds containing single proportions of oxygen, so that the ratio of chlorine to oxygen, is that of 67 to 15, and the number representing chlorine is correctly stated 67.

In like manner it is easy to deduce the number representing the other undecomposed bodies; and they will be found to correspond as nearly as can be expected in whatever way they are obtained. Thus, whether the number representing the proportion in which potassium the basis of potassa combines, be gained from its combination with oxygen or with chlorine, the result will scarcely differ; for 8 grains of potassium converted into the compound of chlorine and potassium I have found gain about 7.1 grains, and when converted into potassa, they gain a grain and $\frac{6}{10}$; and as $7.1:8::67:75.4$; and as $1.6:8::15:75$, giving the number representing potassium as about 75.

It is easy to form a series of proportional numbers by taking $\frac{1}{2}$ of these numbers, on the supposition that water is composed of one proportion of hydrogen and one of oxygen; but in this case the number representing the proportion in which oxygen combines must contain a fraction; and the calculations are much expedited, and the formula rendered more simple, by considering the smallest proportion an integer.

Mr. Higgins has supposed that water is composed of one particle of oxygen and one of hydrogen, and Mr. Dalton, of an atom of each; but in the doctrine of proportions derived from facts, it is not necessary to consider the combining bodies, either as composed of indivisible particles, or even as always united, one and one, or one and two, or one and three proportions. Cases will hereafter be pointed out, in which the ratios

are very different ; and at present, as we have no means whatever of judging either of the relative numbers, figures, or weights, of those particles of bodies which are not in contact, our numerical expressions ought to relate only to the results of experiments.

If it should hereafter be discovered, that any of those substances now considered as undecomposed, consist of other elements, these elements must be represented by some division of their numbers ; and should even hydrogen be found a compounded body, it would merely be necessary to multiply all the numbers representing the other elements, by some common number which would admit of a division into proportions, representing the elements of hydrogen ; so that no discovery concerning the composition of bodies, can interfere with the general law of the definite nature of their combinations.

M. Berzelius and Dr. Wollaston have proposed to make the number representing oxygen unity, because this substance is the most important chemical agent known, and because it enters into a greater number of compounds than any other principle ; but in this case hydrogen would be represented by a fractional part, whilst the multiples of 15 are extremely simple 30, 45, 60, 75, 90, &c. If oxygen were to be represented by 1000, then hydrogen would be represented by 133·33, &c. ; the magnitude of the numbers would enable combinations to be expressed with greater accuracy, but at the same time they would be less easy to acquire and to retain in the memory.

11. If the black oxide of manganese be exposed to a strong heat, it gives off oxygen gas and becomes brown ; but no heat as yet applied is capable of depriving it of the whole of its oxygen.

Hence it is evident that when one proportion of one

substance is combined with more than one proportion of another, the first proportions may be separated with much more facility than the last. There are numbers of other instances: thus the carbonate of soda, which contains two proportions of carbonic acid to one of soda, gives off half its carbonic acid with great facility, by heat, but obstinately retains the other half. Nitric acid is easily brought to the state of nitrous gas by the abstraction of oxygen: nitrous gas with more difficulty is converted into nitrous oxide, but nitrous oxide is still less decomposable than nitrous gas.

When one proportion of a body is combined with two or more proportions of another, it seems to enter with more difficulty into new combinations, than when it is combined with one proportion. Thus iron combined with two proportions of sulphur in golden pyrites is not acted upon by diluted sulphuric acid: but when combined only with one proportion of sulphur, as in the common artificial sulphuret, it is readily acted upon.

It seems from these facts that two or more proportions of one body attract a single proportion of another body with more energy than one proportion, and that two proportions or more adhere to a single proportion with less energy than one proportion; or at least that a second or a third proportion adheres with less energy than the first.

It may possibly be said, that the effect of two or three proportions, in defending one proportion from the action of a new substance, may depend upon mechanical causes, from their more completely enveloping its parts; but the other solution of the effect seems to be the most probable.

13. M. Berthollet, to whom the first distinct views of the relations of the force of attraction to quantity are owing, has endeavoured to prove that these relations are

universal, and that elective affinities cannot strictly be said to exist. He considers the powers of bodies to combine as depending in all cases upon their relative attractions, and upon their acting masses, whatever these may be: and he conceives that in all cases of decomposition, in which two bodies act upon a third, that third is divided between them in proportion to their relative affinities, and their quantities of matter. Were this proposition strictly correct, it is evident that there could be scarcely any definite proportions: a salt crystallizing in a strong alkaline solution, would be strongly alkaline; in a weak one less alkaline; and in an acid solution, it would be acid; which does not seem to be the case. In combinations, in which gaseous bodies are concerned, the particles of which have perfect freedom of motion, the proportions are unchangeable; and in all solid compounds, which have been accurately examined, and in which there is no chance of mechanical mixture, the same law seems to hold good. It is certainly possible to dissolve different bodies in fluid menstrua, in very various proportions, but the result may be a mixture of different solutions, rather than a combination. M. Berthollet brings forward glasses and alloys of metals, as compounds, containing indefinite proportions; but it is not easy to prove, that in these, all the elements are chemically combined; and the points of fusion of alkali, glass, and certain metallic oxides, are so near each other, that transparent mixtures of them *may* be formed. It cannot but be supposed, that the attractive power of matter is general, but in the formation of aggregates, certain arrangements seem to be always uniform.

14. M. Berthollet conceives, that he has proved that a large quantity of a body having a weak affinity, may

separate a part of a second body, from a small quantity of a third, for which it has a strong affinity; but even granting this, it does not destroy the idea of definite proportions. Thus in the fact, noticed by Bergman, the decomposition of sulphate of potassa by nitric acid, one proportion of potassa may be separated from the acid; and the other proportion may combine with two proportions of acid; phenomena analogous to those of common double affinity.

M. Berthollet states, that a large quantity of potassa will separate a small quantity of sulphuric acid from sulphate of baryta; but he made his experiments in contact with the atmosphere, in which carbonic acid constantly floats; and carbonate of potassa and sulphate of baryta mutually decompose each other (7). Even allowing the correctness of his views, still he has not given a complete statement of facts. If potassa separates sulphuric acid from baryta, either there must exist an insoluble sulphate of baryta, containing more baryta than the common sulphate, and which of course may contain two proportions of baryta; or baryta, sulphuric acid, and potassa, must all be dissolved, in the same fluid, which seems highly improbable. M. Berthollet regards baryta as separable from sulphuric acid, by potassa; but has not endeavoured to show in what form it appears after the process.

15. M. Berthollet states, that soda is capable of separating a certain quantity of potassa from sulphuric acid; but, in his experiment, water was present, as the soda must have been a hydrate; and he likewise used alcohol: and the phenomenon may be a phenomenon of double attraction. Potassa has a much stronger attraction for water than soda; and the soda may quit its water, and the potassa its sulphuric acid; and the

effect may be assisted by the stronger attraction of hydrate of potassa for alcohol.

In general, when large quantities of fluid or fusible bodies, are used in experiments, the attraction of the substances which are capable of acting upon each other, is more readily brought into play. In many solutions all the elements are in chemical combination; and their separations depend not merely upon the relative attractions of their parts, but likewise on the manner in which they are acted on by water; and earths, and oxides, are usually thrown down from their solutions in union with water.

16. When an alkali precipitates an earth from its solution in an acid, the earth, according to M. Berthollet's ideas, ought to fall down in combination with a portion of acid. But if a solution of potassa be poured into a sulphuric solution of magnesia, the precipitate produced, after being well washed, affords no indication of the presence of acid; and M. Pfaff has shewn by some very decisive experiments, that magnesia has no action upon neutral combinations of the alkalies and sulphuric acid; and likewise, that the tartaric acid is entirely separated from lime, and the oxalic acid from oxide of lead, by quantities of sulphuric acid, merely sufficient to saturate the two bases; and these are distinct and simple instances of elective attraction. Again, when one metal precipitates another from an acid solution, the body that falls down is usually free both from acid and oxygen: thus zinc precipitates lead and tin; and iron, copper; and the whole of the oxygen and the acid, is transferred from one metal to the other.

17. M. Berthollet, in crystallizing sulphate of potassa, from acid solutions, states that he obtained salts, of

which the first portion contained 55·83 of acid in 100 parts, and another portion only 49·5 ; but it is far from improbable, that these salts were both mixtures of the acidulous sulphate, and the neutral sulphate of potash ; and the idea is strengthened by the circumstance, that he obtained neutral sulphate from the same solution, towards the end of the process.

In cases in which solutions of salts are formed in acid or alkaline menstrua, which are supposed incapable of decomposing them, the results must be considered as depending upon a new combination ; and in the evaporation of the water or of the menstruum, and the crystallization of the remaining constituents, the proportions, that have acted, will determine the nature of the solids which are formed. There appears no difficulty in reconciling the doctrine of definite proportions, with the influence of quantity ; none of the experiments of M. Berthollet can be considered as strictly contradictory to the doctrine, and some of the most important results of this sagacious chemist afford it confirmation.

18. M. Berthollet supposes that the attractions of bodies for each other, are inversely, as the quantities that saturate. Thus, magnesia and ammonia take up more sulphuric acid than equal quantities of potassa ; and therefore he concludes, that magnesia and ammonia have a stronger attraction for acids than potassa : yet potassa instantly separates magnesia and ammonia from acids ; and though the facility with which ammonia is expelled from a compound, may be hypothetically accounted for, by assuming that the ease, with which it takes the gaseous state, assists its escape ; yet magnesia is an opposite case ; and to account for chemical changes, by supposing the effects of forms of matter, which are about to appear, or powers not in actual existence, such

as elasticity or cohesion, is merely the solution of one difficulty, by the creation of another; and ammonia, when solid or fluid, should require a new force to render it elastic: and the cohesion, in a compound, can only be regarded as the exertion of the chemical attractions of its elements. The action between the constituents of a compound must be mutual; sulphuric acid, there is every reason to believe, has as much attraction for baryta, as baryta for sulphuric acid: and baryta is the alkaline substance, of which the largest quantity is required to saturate sulphuric acid; therefore, on M. Berthollet's view, it has the weakest affinity for that acid; but less sulphuric acid saturates this substance, than any other earthy or alkaline body; therefore, according to M. Berthollet, sulphuric acid has a stronger affinity for baryta than for any other substance; which is contradictory.

19. It cannot be laid down as a general law, that the attractions of bodies are connected with the weights of the proportions in which they combine; yet in some cases the proportions, which unite in the greatest quantity, or the bodies represented by the highest numbers, are separated by proportions combining in smaller quantity, or by bodies represented by lower numbers. Thus gold, platina, mercury, and silver, are separated in their metallic states by the common metals, which are represented by much lower numbers, and the metallic oxides by the alkalies; but there are many exceptions; and the intensity of attraction seems to be dependent upon other causes, which are intimately related to the electrical phenomena, to be discussed in the next section.

20. The uniformity of the law of condensation, when gases combine and form denser gaseous compounds, in

which the volume is unaltered, or in which one of the elements is condensed to $\frac{1}{2}$, or in which both are condensed to $\frac{1}{2}$, and the regularity of the forms of solid bodies seem to depend entirely upon the constancy of the nature of the combination, and probably upon the corpuscular aggregates being all of the same kind. If the particles of matter be supposed to be globular, or to act in spheres of attraction and repulsion, it would be easy to account for their forms, by supposing a few independent primary arrangements. Thus, four particles may compose a tetrahedron, five a tetrahedral pyramid, six an octahedron, or a trihedral prism, and eight a cube or a rhomboid.

21. It would be premature in this part of the work, to enter upon any more minute views of the laws of attraction, and the more refined details will properly follow the history of the agencies of different bodies on each other.

With respect to a power so constantly in action, it is necessary, however, even at an early period of the study, to possess some definite ideas. If it be regarded as capricious in its effects, and tending constantly to produce different arrangements, chemistry would be without a guide, without certain combinations, and no results of analysis could be perfectly alike; but fortunately for the progress of science, this is not the case: the changes of the terrestrial cycle of events, like the arrangements of the heavens, and the system of the planetary motions, are characterized by uniformity and simplicity; weight and measure can be applied to them, their order perceived, and their laws discovered.

VII. *Of Electrical Attraction and Repulsion, and their Relations to Chemical Changes.*

1. If a piece of dry silk be briskly rubbed against a warm plate of polished flint glass, it will be found to have acquired the property of adhering to it, which it will retain for some seconds; if at the time this adhesive power exists, the silk and glass be separated from each other, they will both be found to have gained the property of attracting very light substances, such as the ashes of paper or fragments of gold leaf; and the long filaments of the silk, if there be any, will be seen to repel each other.

2. These bodies are said to be *electrically excited*, and the phenomena are called *electrical* phenomena. The peculiar circumstances under which they occur, are best observed by the use of an instrument called the electrical machine; it consists of a cylinder of glass* supported upon glass pillars, and which can be made to revolve, so as to press against a cushion of silk rubbed over with a little amalgam of zinc and mercury; and of two cylinders of metal, one in contact with the cushion, and the other opposite to the glass cylinder, both supported upon glass.

3. If two gilt pith balls, suspended upon strings of silk covered with tinsel, be hung upon a wire, placed in contact with either of the metallic cylinders, and the machine be put in action, the balls will repel each other; but if one ball be attached to a wire, connected with one metallic cylinder, and the other ball be attached to a wire connected with the other, the two balls, when the machine is put into action, will attract each other; and

* Plate II. fig. 8.

at the moment that they come in contact, sparks of light will be perceived, if the experiment be made under favourable circumstances.

As the two balls, when in contact with the same cylinder, may be considered as receiving the same impulse or impression, they are said to be *similarly electrified*; but when in contact with different cylinders, they are said to be *differently electrified*; and electrified bodies that repel each other, are considered as in the same electrical states; those that attract each other as in different electrical states.

4. There are probably no two bodies differing in nature, which are not capable of exhibiting electrical phenomena, either by contact, pressure, or friction; but the first substances in which the property was observed, were *vitreous* and *resinous* bodies; and hence the different states were called states of *resinous* and *vitreous* electricity; and resinous bodies bear the same relation to flint glass, as silk. The terms, *negative* and *positive* electricity, have been likewise adopted, on the idea, that the phenomena depend upon a peculiar subtile fluid, which becomes in excess in the vitreous, and deficient in the resinous bodies; and which is conceived by its motion and transfer, to produce the electrical phenomena.

5. Flint glass and silk, silk and sulphur, sulphur and metals, resin and metals, all by friction or contact, become strongly electrical, and of course attractive, and communicate their attractive powers to small masses of matter brought in contact with them; a pith ball, or a slip of gold leaf that has been touched by flint glass, excited by silk, will be repelled by a ball or slip that has been touched by silk excited by sulphur, or by a ball or slip that has been touched by sulphur excited by

metals, so that the attractive and repellent states depend entirely upon the actions of the two substances, and not upon any power peculiar to and inherent in each.

6. It is upon this circumstance, that the *electrometer*, which might be called the *differential* one, is framed; it consists of two gold leaves attached to a metallic plate, and included in a hollow cylinder of glass,* fixed upon another metallic plate, which is connected with two pieces of tin foil, pasted upon the glass opposite to the leaves. When any electrified body is made to touch the upper plate, the gold leaves diverge; if their divergence is increased by the approach of flint glass excited by silk, they are said to have the same state as the glass, the vitreous or the positive; if their divergence is diminished, they are said to be in the opposite state, or to possess the resinous or negative electricity.

7. When luminous phenomena are connected with electrical excitation, the different states may be known by presenting a metallic point to the excited body; if rays of light issue from the point to the body, it is said to be negatively electrified: but if the point appears simply luminous, without sending off any rays, the electricity is said to be positive.

8. For measuring small degrees of electricity of bodies, as compared with those of others of the same kind, the *electrical balance* of Coulomb is applied; it consists of a gilt pith ball, placed upon a metallic rod, on the opposite extremity of which is a thin leaf of metal; the rod is suspended horizontally, by a fine metallic wire, which passes into a glass tube, to the top of which it is attached; the glass tube is inserted into a cylinder of glass, which contains a copper ball, connected with a small bar of metal, which is carried through an aperture

* Plate II. fig. 9.

in the glass cylinder, into the atmosphere ; a very small force only is required to twist the wire, and when the two balls are brought in contact, and the bar touched by the electrified body, they gain the same kind of electricity, and repel each other ; and the degree of their repulsion may be measured by a scale of degrees, made on the circumference of the cylinder.*

9. Bodies receive the electrical influence in different manners. If a rod of glass be brought in contact with any excited electrical body, it will receive the electrical influence in the part where it touched the body, and will be electrical, to a little distance, round the point of contact ; but its remote parts will not be affected. A rod of metal, on the contrary, suspended on a rod of glass, and brought in contact with an electrical surface, instantly becomes electrical throughout. The glass, in common philosophical language, is said to be a *non-conductor* of electricity, or an *insulating* substance ; the metal a *conductor*. Some bodies are affected to a much greater extent than glass, but not nearly so much as metals ; such are animal and vegetable substances, water, and fluids containing water ; they are said to be *imperfect conductors*. According to the statements of Mr. Cavendish, iron conducts 400 millions of times better than water, sea water 100 times better than distilled water, and water saturated with salt, 720 times better. The mineral acids are the best fluid conducting substances known, and after them, saline solutions, the powers of which appear to be nearly in proportion to the quantities of salts they contain. Charcoal and metals, and the greater number of inflammable metallic compounds are conductors. Alcohol and ether, are very imperfect conductors ; and sulphur, oils, resinous substances,

*. Plate II. fig. 10.

metallic oxides and compounds of chlorine, nonconductors.

10. There is a stone found in many parts of the world, called tourmaline, which is sometimes crystallized as a nine-sided prism, terminated by a three-sided and a six-sided pyramid; when this substance is gently heated, it becomes electrical, and one extremity, that terminated by the six-sided pyramid, is positive, the other is negative; to a certain extent, its electricities are exalted by increasing the temperature; when it begins to cool, it is still found electrical; but the electricities are changed, the pyramid, before positive, is now negative, and vice versa. When the stone is of considerable size, flashes of light may be seen along its surface.

There are other gems and crystallized substances, which possess a property similar to that of the tourmaline. The luminous appearance of some diamonds, when heated, probably depends upon their electrical excitation. The substance called the Boracite, which is a cube, having its edges and angles defective, becomes electrical by heat, and in one variety presents no less than eight sides, in different states, four positive, four negative; and the opposite poles are in the direction of the axes of the crystal.

11. It would appear, that in all cases of electrical action, the two electrical states are always coincident, either in different parts of the same body, or in two bodies; and that they are always equal, and capable of neutralizing each other. If a connection be made by a wire, between the positive and negative conductors of the electrical machine, during the time of its action, all electrical effects cease; and to produce a succession of effects, both conductors must be brought near bodies

connected with the ground, which gain the opposite state, in consequence of what may be called *induction*, and which will be explained in the next paragraph.

12. When a non-conductor, or imperfect conductor, provided it be a thin plate of matter, placed upon a conductor, is brought in contact with an excited electrical body, the surface, opposite to that in contact, gains the opposite electricity from that of the excited body; and if the plate be removed from the conductor and the source of electricity, it is found to possess two surfaces in opposite states. If a conductor be brought into the neighbourhood of an excited body, the air, which is a non-conductor, being between them, that extremity of the conductor, which is opposite to the excited body, gains the opposite electricity, and the other extremity, if opposite to a body connected with the ground, gains the same electricity, and the middle point is not electrical at all. This is easily proved, by examining the electricity of three sets of gilt pith balls raised on wires on the different parts of the conductor, which is thus effected by *induced* electricity.

If, instead of air, a plate of mica or glass be between the two conductors, the same phenomena will occur; so that it would appear that the conductor merely gains two opposite electricities, or polar electricities, of the same kind as those of the non-conductor. The phenomena of sparks, of discharges, and of accumulated electricity, depend upon this law. In the case of the common electrical spark, a stratum of air is charged in the same manner as a glass bottle, partially coated with tin-foil, is charged in the Leyden experiment;* when the hand is held near the positive conductor of an electrical machine, the person standing on the ground,

* Plate II. fig. 11.

the hand is rendered negative, and the states become exalted, till the polarities, as they may be called, are annihilated through the air, producing a spark, a snap, and a distinct sensation. If a number of small pith balls, placed upon a surface of metal, are caused to approach an electrified body, they are brought into the opposite state by induction, and are attracted towards the body; but when they come in contact with it, this state is destroyed, they gain the same state, and are repelled; and if they are properly placed, their alternate attractions and repulsions may be produced, as long as the machine is in action.

13. If a number of cylinders of metal, *insulated* on glass, be placed in a line with each other, but not in contact, and the last be connected with the ground;* when a powerfully electrified conductor of a machine is brought opposite to the first, they will all become electrical, and every insulated cylinder will present two poles; the negative pole of one being opposite to the positive pole of the other; and if a spark is produced by means of the last, sparks occur throughout the whole arrangement. In like manner a series of Leyden jars may be made to charge each other, the outer surface of the first rendering negative the inner surface of the second, and so on; and by connecting the surfaces that have the same kind of electricity, in the first place, and then connecting two opposite surfaces in the series, a powerful explosion† may be produced.

14. When a point connected with the ground is brought near an electrified substance, it rapidly gains the opposite state, and an immediate discharge takes place, which continues till the equilibrium is restored. Large surfaces are electrified by induction much more

* Plate III. fig. 12.

† Plate III. fig. 13.

slowly than small ones, and are capable of accumulating much more electricity, which renders the discharge from them much more violent. Indeed, the electrical powers seem entirely to belong to the surfaces of bodies, and not to be connected with the quantity of solid matter they contain.

15. It is in consequence of the principle of induction, that the condensing electrometer is so much more sensible than the common electrometer; this instrument consists of two plates of polished metal,* the surfaces of which are parallel, one connected with the plate of the electrometer, the other moveable, in connexion with the ground, and the plates are very near each other. When the body supposed to be electrical, is made to touch the top of the electrometer, and is afterwards removed, in separating the plates, the effect will be perceived.

16. The difference in what are called the conducting powers of bodies, seems to depend entirely upon the different manner in which they receive the electrical polarities, or in which their parts become capable of communicating attractive or repellent powers, to other matter. Non-conductors appear to receive polarities, only with great difficulty, but retain them for a long while, and present probably a number of different alternations of poles, within a small space, and cannot be affected to any great distance. Imperfect conductors receive polarity with more facility, but present fewer alternations, and preserve their electricities for a shorter time. Perfect conductors are easily affected throughout, but present at most only two poles, and the powers rapidly destroy each other. The difficulty with which non-conductors receive polarity, is shewn in the phe-

* Plate III. fig. 14.

nomena of charging thick and thin coated plates of glass and mica. The thin plates are capable of being charged much more highly than the thick ones, and the accumulation on the opposite surfaces is much greater.

Rarefied air, or gaseous matter, is much more susceptible of receiving polarities, than dense air or gaseous matter; and hence, the electrical spark will pass much further through rarefied air or light gases, than through dense air or heavy gases; it passes much further, likewise in gases, than in non-conducting fluids.

17. If a non-conducting surface, coated with two conducting surfaces, and charged so as to give a spark of an inch in length, through air, be connected by both its conducting surfaces, with a similar apparatus not charged; then both systems may be discharged together; but the spark they will give, will be only half as long as the single one would have given, if discharged alone. The *quantity* of the electricity in this case, is conceived not to be altered, but its *intensity* is said to be only half as great when it is discharged from a double surface; and these expressions of intensity and quantity, though it is not easy to attach any very definite ideas to them, are nevertheless useful, in giving more facility to the arrangement of some important electrical phenomena.

18. When very small conducting surfaces are used for conveying very large quantities of electricity, they become ignited; and of the different conductors that have been compared, charcoal is most easily heated by electrical discharges,* next iron, platina, gold, then

* The conclusions are drawn from experiments made by the electricity of the Voltaic apparatus.

copper, and lastly zinc. The phenomena of electrical ignition, whether taking place in gaseous, fluid, or solid bodies, always seem to be the result of a violent exertion of the electrical attractive and repellent powers, which may be connected with motions of the particles of the substances affected. That no subtile fluid, such as the matter of heat has been imagined to be, can be discharged from these substances, in consequence of the effect of the electricity, seems probable, from the circumstance, that a wire of platina may be preserved in a state of intense ignition in vacuo, by means of the Voltaic apparatus, (an instrument which will be immediately described,) for an unlimited time; and such a wire cannot be supposed to contain an inexhaustible quantity of subtile matter.

19. Certain changes in the forms of substances, are always connected with electrical effects. Thus when vapour is formed or condensed, the bodies in contact with the vapour, become electrical. If, for instance, a plate of metal, strongly heated, be placed upon an electrometer, and a drop of water be poured upon the plate, at the moment the water rises in vapour, the gold leaves of the electrometer diverge with negative electricity. Sulphur, when melted, becomes strongly electrical during the time of congelation; and the case seems to be analagous, with respect to non-conducting substances in general, when they change their forms.

20. As electricity appears to result from the general powers or agencies of matter, it is obvious, that it must be continually exhibited in nature, and that a number of important phenomena must depend upon its operation. When aqueous vapour is condensed, the clouds formed are usually more or less electrical; and the earth below them being brought into an opposite state,

by induction, a discharge takes place when the clouds approach within a certain distance, constituting lightning; and the undulation of the air, produced by the discharge, is the cause of thunder, which is more or less intense, and of longer or shorter duration, according to the quantity of air acted upon, and the distance of the place, where the report is heard from the point of the discharge. It may not be uninteresting to give a further illustration of this idea: electrical effects take place in no sensible time; it has been found, that a discharge through a circuit of four miles, is instantaneous; but sound moves at the rate of about twelve miles in a minute. Now, supposing the lightning to pass through a space of some miles, the explosion will be first heard from the point of the air agitated, nearest to the spectator; it will gradually come from the more distant parts of the course of the electricity, and last of all, will be heard from the remote extremity; and the different degrees of the agitation of the air, and likewise the difference of the distance, will account for the different intensities of the sound, and its apparent reverberations and changes.

21. In a violent thunder storm, when the sound instantly succeeds the flash, the persons who witness the circumstance, are in some danger: when the interval is a quarter of a minute, they are secure. In a thunder storm, the lowest ground is the safest place, and a horizontal posture, the least dangerous; the neighbourhood of trees, or buildings, should be avoided, particularly of trees, the living juices of which are calculated to conduct the electricity, and make part of a circuit. In a house, the cellars are the safest places, and in a room the person should stand as far as possible from the fire. The means adopted by Franklin have, however, to a

great extent, averted the destructive effects of atmospheric electricity; and by pointed conductors, the thunder cloud is disarmed of its terrors, and the lightning slowly discharged in harmless coruscations.

If a school-boy's kite be mounted high in the atmosphere, by means of a string, containing filaments of metal, fastened to a conductor, fixed on a glass rod; the conductor usually gives signs of electricity, which will be greatest, when clouds are floating in the atmosphere; and it was by means of a simple apparatus of this kind, that the American Philosopher effected his grand discovery of the identity of electricity and lightning.

The water-spout is probably the result of the operation of a weakly electrical cloud, at an inconsiderable elevation above the sea, brought into an opposite state: and the attraction of the lower part of the cloud, for the surface of the water, may be the immediate cause of this extraordinary phenomenon.

The coruscations of the *Aurora Borealis*, and *Australis*, precisely resemble strong artificial electricity, discharged through rare air; and as the poles are non-conductors, being coated with ice or snow, and as vapour must be constantly formed in the atmosphere above them; the idea of Franklin is not improbable, that the Auroras may arise from a discharge of electricity, accumulated in the atmosphere near the poles, into its rarer parts; though other solutions of the phenomena may be given on the idea, that the earth itself is endowed with electrical polarity; or that the motions of the atmosphere produce the effect: but all views on this subject must be hypothetical, and the light may result from other causes than electrical action.

22. The common exhibition of electrical effects, is in

attractions and repulsions, in which masses of matter are concerned ; but there are other effects, in which the changes that take place, operate in a manner, in small spaces of time imperceptible, and in which the effects are produced upon the chemical arrangements of bodies.

If a piece of zinc and a piece of copper be brought in contact with each other, they will form a weak electrical combination, of which the zinc will be positive, the copper negative ; this may be learnt by the use of a delicate condensing electrometer ; or by pouring zinc filings through holes, in a plate of copper, upon a common electrometer ; but the power of the combination may be most distinctly exhibited in the experiments, called *Galvanic experiments*, by connecting the two metals, which must be in contact with each other, with a nerve and muscle in the limb of an animal recently deprived of life, a frog for instance ; at the moment the contact is completed, or the circuit made, one metal touching the muscle, the other the nerve, violent contractions of the limb will be occasioned. If a piece of zinc and copper, in contact with each other in one point, be placed in contact in other points with the same portion of water ; the zinc will corrode and attract oxygen from the water, much more rapidly than if it had not been in contact with the copper ; and if a small quantity of sulphuric acid be added to the water, it will be seen that globules of inflammable air are given off from the copper, though it is not dissolved nor acted upon.

23. The connection of chemical effects, with the exhibition of electrical powers is, however, best witnessed in combinations, in which these powers are accumulated by alternations of different metals and

fluids. If plates of copper and zinc two or three inches square, and pieces of cloth of the same size soaked in a solution of salt, or sal ammoniac, or nitre, be arranged in the order of copper, zinc, moistened cloth, and so on, and made into an insulated pile, of which the series are 200 ;* several remarkable phenomena will occur.

When one hand is applied to the bottom of the pile, and the other to the top, both hands being moistened, a shock will be perceived.

When a metallic wire, having a bit of well burned charcoal at its extremity, is made to connect the two extremities of the pile, a spark will be perceived, or the point of the charcoal will become ignited.

A wire, connected with the top of the pile, brought in contact with a sensible electrometer, will cause the leaves to diverge; and by removing the wire and applying excited glass to the electrometer, it will be found that the electricity is positive; a wire connected with the bottom of the pile will affect it with negative electricity; a wire from the middle of the pile will have no influence on the instrument.

If wires of platina from the extremities of the pile be introduced into water, or into two portions of water connected by moist substances, oxygen gas will separate at the wire exhibiting the positive electricity, and hydrogen gas at the wire exhibiting the negative electricity; and the proportions are such, when the proper circumstances exist, that they will produce water when exploded by the electrical spark, that is, the volume of hydrogen will be to that of the oxygen as two to one.

If the same wires be introduced into a strong solution of sulphuric or phosphoric acid, or into metallic solutions, oxygen will separate at the positive surface, the

* See Plate III. fig. 14, 15.

inflammable or metallic matter contained in the solution at the negative surface.

When any substance rendered fluid by heat, consisting of water, oxygen and inflammable or metallic matter, is exposed to those wires, similar phenomena occur.

When any solution of a neutral salt containing acid, united to alkaline, earthy, or common metallic matter, is used; besides the other phenomena that take place, acid matter collects round the positively electrified surface; alkali, earth, or oxide, round the negative surface; and if two separate vessels are employed to contain the solution, connected by moist asbestos, it is found that the acid collected in the vessel containing the wire, positively electrified, will be in definite proportion to the matter collected in the other cup; that is, it will form with it a neutrosaline compound.

If a solution of muriatic acid in water be acted on by the wires, hydrogen will separate at the negative surface, and chlorine at the positive surface.

24. This apparatus, which exhibits in so distinct a manner the relations of electrical polarities to chemical attractions, is the grand invention of Volta, made known in the first year of this century; its electrical effects have been long known, but the phenomena of its operation in decomposing bodies, are of more recent discovery.

Several modes of constructing it have been adopted, some of which are much superior in point of convenience, to that which has been just described.

One mode is by soldering the plates of zinc and copper together, and by cementing them into troughs of baked wood, covered with cement, in the regular order, so as to form cells to be filled with the fluid menstruum;

each surface of zinc being opposite to a surface of copper; and this combination is very simple and easy of application.

Another form is that of introducing plates of copper and of zinc, fastened together by a slip of copper, into a trough of porcelain containing a number of cells corresponding to the number of the series. The different series may be introduced separately into the troughs, and taken out without the necessity of changing the fluid, or they may be attached to a piece of baked wood and (when the number is not very large) introduced into the cells, or taken out together.*

25. Similar polar electrical arrangements to those formed by zinc and copper, may be made by various alternations of conducting and imperfect conducting substances; but for the accumulation of the power, the series must consist of three substances or more, and one at least must be a conductor. Silver or copper, when brought in contact with a solution of a compound of sulphur and potassa, at one extremity, and in contact with water or a solution of nitric acid, at the other extremity, some saline solution being between the sulphuretted and the acid solutions, forms an element of a powerful combination, which will give shocks when fifty are put together; the order is copper, cloth of the same size moistened with solution of nitric acid, cloth moistened in solution of common salt, cloth moistened in solution of the compound of sulphur, copper, and so on; the specific gravities of the solutions should be in the order in which they are arranged, to prevent the mixture of the acid and sulphuretted solution; that is, the heaviest solution should be placed lowest.

The tables annexed contain some series, which form

* Plate III. fig. 16.

Voltaic electrical combinations, arranged in the order of their powers; the substance most active being named first in each column.

Table of some Electrical Arrangements, which by Combination form Voltaic Batteries, composed of two Conductors and one Imperfect Conductor.

Zinc Iron Tin Lead Copper Silver Gold Platina Charcoal	Each of these is the positive pole to all the metals below it, and negative with respect to the metals above it in the column.	Solutions of nitric acid of muriatic acid of sulphuric acid of sal-ammoniac of nitre other neutralsalts
--	--	--

Table of some Electrical Arrangements, consisting of one Conductor and two Imperfect Conductors.

Solution of sulphur and potash of potash of soda	Copper Silver Lead Tin Zinc Other metals Charcoal	Nitric Sulphuric acid Muriatic acid Any solutions containing acid
--	---	---

The metals having the strongest attraction for oxygen, are the metals which form the positive pole, in all cases in which the fluid menstrea act chemically by affording oxygen; but when the fluid menstrea afford sulphur to the metals, the metal having the strongest attraction for sulphur under the existing circumstances, determines the positive pole; thus in a series of copper and iron, introduced into a porcelain trough, the cells of which are filled with water or with

acid solutions, the iron is positive, and the copper negative ; but when the cells are filled with solutions of sulphur and potash, the copper is positive and the iron negative.

In all combinations in which one metal is concerned, the surface opposite the acid is negative, that in contact with solution of alkali and sulphur, or of alkali, is positive.

26. The energy of a combination to give repulsive or attractive powers to masses of matter or to affect the electrometer, seems to increase with the number of the series, as does the power to give shocks, and to decompose bodies ; but as long as the surface of the gold leaves in the electrometer, or of the human body, or of the water acted upon, is the same, and less than that of the acting plates, increase of surface of the plates is connected with no increase of power. In the operation upon metallic substances or charcoal, or upon good imperfect conductors, the case, however, is different. Thus, though a battery composed of plates of copper and zinc a foot square, will not affect the condensing electrometer more, nor decompose more water, nor give greater shocks to the fingers, than a battery containing plates of an inch square, yet it will ignite more than 100 times as much fine platina wire, and decompose sulphuric acid, and the water in strong saline solutions with infinitely more rapidity. This has been expressed by Mr. Cavendish in the statement, that the intensity is the same in both cases ; but that the quantity is in some ratio as the surface. The quantity in the small plates is as much or more than such imperfect conductors as water and the human body can carry off by a small surface : whilst better conductors can transmit the whole quantity afforded by the large plates, even when used

in very thin laminæ or wires. The correctness of this view may be shewn by a very simple experiment. Let two platina wires, from the extremities of a battery composed of plates of a foot square, be plunged into water, the quantity of gas disengaged from the wires will be nearly the same as from an equal number of plates of an inch square; let the fingers of each hand, moistened with water, be applied to the two extremities of the battery, a shock will be perceived nearly the same as if there had been no connection between the wires and the water. Whilst the circuit exists through the human body and through water, let a wire attached to a thin slip of charcoal be made to connect the two poles of the battery, the charcoal will become vividly ignited. The water and the animal substance discharge the electricity of a surface, probably not superior to their own surface of contact with the metals; the wires discharge all the residual electricity of the plates; and if a similar experiment be made on plates of an inch square, there will scarcely be any sensation, when the hands are made to connect the ends of the battery, a circuit being previously made through water; and no spark when charcoal is made the medium of connection, imperfect conductors having been previously applied.

The first distinct experiment upon the igniting powers of large plates was performed by MM. Fourcroy, Vauquelin, and Thenard. But the grandest combination ever constructed for exhibiting the effects of extensive surface, was made by Mr. Children: it consists of twenty double plates four feet by two; of which the whole surfaces are exposed, in a wooden trough, in cells covered with cement, to the action of diluted acids. This battery, when in full action, had no more

effect on water or on the human body than one containing an equal number of small plates; but when the circuit was made through metallic wires, the phenomena were of the most brilliant kind. A platina wire of one-thirtieth of an inch in thickness, and eighteen inches long, placed in the circuit between bars of copper, instantly became red hot, then white hot, the brilliancy of the light was soon insupportable to the eye, and in a few seconds the metal fell fused into globules. The other metals were easily fused or dissipated in vapour by this power. Points of charcoal ignited by it produced a light so vivid, that even the sunshine compared with it appeared feeble.

Mr. Children has another battery in construction, the plates of which are double the size of that just described, and which are to be arranged in pairs in single troughs, and connected by means of plates of lead in regular order.*

27. The most powerful combination that exists in which number of alternations is combined with extent of surface, is that constructed by the subscriptions of a few zealous cultivators and patrons of science, in the laboratory of the Royal Institution. It consists of two hundred instruments, connected together in regular order, each composed of ten double plates arranged in cells of porcelain, and containing in each plate thirty-two square inches; so that the whole number of double plates is 2000, and the whole surface 128,000 square inches. This battery, when the cells were filled with 60 parts of water mixed with one part of nitric acid, and one part of sulphuric acid, afforded a series of brilliant and impressive effects. When pieces of charcoal about

* [For an account of this battery and its effects, vide *Phil. Trans.* for 1815.]

an inch long and one-sixth of an inch in diameter, were brought near each other (within the thirtieth or fortieth part of an inch,) a bright spark was produced, and more than half the volume of the charcoal became ignited to whiteness, and by withdrawing the points from each other a constant discharge took place through the heated air, in a space equal at least to four inches, producing a most brilliant ascending arch of light, broad, and conical in form in the middle.* When any substance was introduced into this arch, it instantly became ignited; platina melted as readily in it as wax in the flame of a common candle; quartz, the sapphire, magnesia, lime, all entered into fusion; fragments of diamond, and points of charcoal and plumbago, rapidly disappeared, and seemed to evaporate in it, even when the connection was made in a receiver exhausted by the air pump; but there was no evidence of their having previously undergone fusion.

When the communication between the points positively and negatively electrified was made in air, rarefied in the receiver of the air pump, the distance at which the discharge took place increased as the exhaustion was made, and when the atmosphere in the vessel supported only one-fourth of an inch of mercury in the barometrical gage, the sparks passed through a space of nearly half an inch; and by withdrawing the points from each other, the discharge was made through six or seven inches, producing a most beautiful coruscation of purple light, the charcoal became intensely ignited, and some platina wire attached to it, fused with brilliant scintillations, and fell in large globules upon the plate of the pump. All the phenomena of chemical decomposition were produced with intense rapidity by this

* Plate III. fig. 17.

combination. When the points of charcoal were brought near each other in non-conducting fluids, such as oils, ether, and oxymuriatic compounds, brilliant sparks occurred, and elastic matter was rapidly generated; and such was the intensity of the electricity, that sparks were produced, even in good imperfect conductors, such as the nitric and sulphuric acids.

When the two conductors from the ends of the combination were connected with a Leyden battery, one with the internal, the other with the external coating, the battery instantly became charged, and on removing the wires, and making the proper connections, either a shock or a spark could be perceived; and the least possible time of contact was sufficient to renew the charge to its full intensity.

28. The general facts of the connection of the increase of the different powers of the battery with the increase of the number and surface of the series, are very distinct; but to determine the exact ratio of the connection is a problem not easy of solution.

MM. Gay Lussac and Thenard have announced, that the power of chemical decomposition increases only as the cube root of the number of plates; but their experiments were made with parts of piles of a construction very unfavourable for gaining accurate results; and in various trials made with great care in the laboratory of the Royal Institution, the results were altogether different. The batteries employed were parts of the great combination, carefully insulated, and similarly charged; arcs of zinc and silver presenting equal surfaces, and arranged in equal glasses filled with the same kind of fluid, were likewise used; and the tubes for collecting the gases were precisely similar, and filled with the same solution of potassa.* In these experiments ten

* Plate IV. fig. 18.

pairs of plates produced fifteen measures of gas; twenty pairs in the same time produced forty-nine: again, ten pairs produced five measures; forty pairs in the same time produced seventy-eight measures. In experiments made with arcs, and which appeared unexceptionable, four pairs produced one measure of gas; twelve pairs in the same time produced nine and $\frac{7}{10}$ of gas; six pairs produced one measure of gas; thirty pairs, under like circumstances, produced 24·5 measures; and these quantities are nearly as the squares of the numbers.

It would appear from the experiments of Vanmarum and Pfaff, confirmed by those of Messrs. Wilkinson, Cuthbertson, and Singer, that the increase of power of batteries, the plates of which have equal surfaces, is as the number. I found that ten double plates, each having a surface of a hundred square inches, ignited two inches of platina in wire of one-eightieth of an inch; twenty plates, five inches; forty plates, eleven inches: but the results of experiments on higher numbers were not satisfactory; for one hundred double plates of thirty-two square inches each, ignited three inches of platina wire of one-seventieth, and one thousand ignited only thirteen inches, and the charges of diluted acid were similar in both cases.

The power of ignition for equal numbers of plates, seems to increase in a very high ratio with the increase of surface, probably higher than even the square; for twenty double plates, containing each two square feet, did not ignite one-sixteenth as much wire as twenty, containing each eight square feet, the acid employed being of the same strength in both cases.

Numerous circumstances are opposed to the accuracy of experiments made with high numbers, or very large surfaces; the activity of combinations rapidly

diminishes in consequence of the decomposition of the menstruum used; and this decomposition is much more violent, the greater the number and surface of the alternations; the vapour rising likewise, when the action is intense, interferes by its conducting powers, and the gas by its want of conducting power; and when series containing above five hundred double plates are used, unless the insulation is very perfect, there is a considerable loss of electricity; thus the great battery of two thousand double plates belonging to the Royal Institution, will scarcely act by its true poles, when arranged on a floor of stone, and requires not merely the insulation of porcelain, but likewise of dry wood; and when arranged on a stone floor, it is hardly possible to walk near any of the approaching series without receiving shocks. In cases of the ignition of wire, the cooling influence of the substances in contact, and of that part of the chain not ignited, interferes most, when small quantities of wire are employed, and with feeble powers; and hence the effect is at first in a lower and then in a higher ratio than the number, when the whole range is small, as in the experiments above stated. If there is an imperfect connection in any of the series, a great diminution of power is the consequence. If one plate is corroded, or covered with more oxide than the rest, there is a general loss of effect. If copper is substituted for zinc, or zinc for copper, in a single series, the result is similar; and I find that a platina wire, introduced in the place of an arc of silver and zinc, in a series of thirty, diminished its power of producing gas so much, that it was equal only to that of four.

29. The circumstance most important in electricity, perhaps, is its connection with the chemical powers of matter, and the manner in which it modifies, exalts, or

destroys these powers. Most of the substances that act distinctly upon each other electrically, are likewise such as act chemically, when their particles have freedom of motion: this is the case with the different metals, with sulphur and the metals, with acid and alkaline substances; and the relations of bodies are uniform; those that have the highest attracting powers being in the relation of positive, in arrangements in which chemical changes can go on. Thus, as is shown in the tables, page 107, zinc is positive with respect to iron, iron with respect to copper, copper with respect to silver, and so on in all combinations in which oxygen is capable of being combined with the metal; but copper is positive with respect to iron in compound menstrua containing sulphur; the electrical power being in all cases apparently connected with the power of chemical combination.

Crystals of oxalic acid touched by dry quick-lime exhibit electrical powers; and the acid is negative, the lime positive.

All the acid crystals, upon which I have experimented when touched by a plate of metal, render it positive. And in Voltaic combinations with single plates or arcs of metal, as is stated in page 108, the metal is negative on the side opposed to the acid, and positive on the side or pole opposed to the alkali.

Bodies that exhibit electrical effects previous to their chemical action on each other, lose this power during combination. Thus, if a polished plate of zinc is made to touch a surface of dry mercury, and quickly separated, it is found positively electrical, and the effect is increased by heat; but if it be so heated as to amalgamate with the surface of the mercury, it no longer exhibits any marks of electricity. The case is analagous

with copper and sulphur; and iron acts more powerfully than zinc with quicksilver in a permanent electrical combination, as in the experiments of Colonel Haldane; apparently, because under common circumstances it is incapable of amalgamating with that metal. When any conducting substance, capable of combining with oxygen, has its positive electricity increased, it will attract oxygen with more energy from any imperfect conducting medium; and metallic bodies which in their common state have no action upon water, such as silver, attract oxygen from it easily, when connected with the positive pole in the Voltaic circuit; and bodies that act upon water, such as zinc and iron, so as to decompose it slowly, refuse to attract oxygen from it when they are negatively electrified in the Voltaic circuit.

Acids, which are negative with respect to alkalies, metals, and earths, are separated from these bodies in the Voltaic circuit at the positive surface; and alkalies, metals, and earths, are separated from acids at the negative surface: and such are the attracting powers of these surfaces, that acids are transferred through alkaline solutions, and alkalies through acid solutions, to the surfaces where they have their points of rest. It is easy to shew this by making a combination of three agate cups,* one containing sulphate of potassa, one weak nitric acid, and the third distilled water, and connecting them by asbestos moistened in pure water, in such a manner, that the surface of the acid is lower than the surface of the fluid in the other two cups. When two wires of platina from a powerful Voltaic apparatus are introduced into the two extreme cups, the solution of the salt being positively electrified, a decomposition will take place, and in a certain time a

* Plate IV. fig. 19.

portion of potassa will be found dissolved in the cup in contact with the negative wire, though the fluid in the middle cup will still be sensibly acid.

30. Such are the decomposing powers of electricity, that not even insoluble compounds are capable of resisting their energy; for even glass, sulphate of baryta, fluor spar, &c. when moistened and placed in contact with electrified surfaces from the Voltaic apparatus, are slowly acted upon, and the alkaline, earthy, or acid matter carried to the poles in the common order. Not even the most solid aggregates, nor the firmest compounds, are capable of resisting this mode of attack; its operation is slow, but the results are certain; and sooner or later, by means of it, bodies are resolved into simpler forms of matter.

31. It is in consequence of the phenomena of electrical decomposition, in which metals, inflammable bodies, alkalies, earths, and oxides, are determined to the negative surface, and oxygen, chlorine, and acids, to the positive surface, that for some time it was conceived, that various substances, might be composed from pure water, by means of electricity, such as potassa, soda, and muriatic acid. A strict investigation of the circumstances under which these substances appeared, led me to discover that they were always furnished from the vessels, or from impurities in the water, and enabled me to determine the general principles of electrical decomposition, and to apply this power to the resolution of some species of matter, of unknown nature, into their elements.

32. The connection of electrical phenomena and chemical changes is evident likewise in the general phenomena of the battery. The most powerful Voltaic combinations are formed by substances that act chemi-

cally with most energy upon each other; and such substances as undergo no chemical changes in the combination, exhibit no electrical powers. Thus, zinc, copper, and nitric acid form a powerful battery; whilst silver, gold, and water, which do not act chemically on each other, in series of the same number, produce no sensible effect. These circumstances led some philosophers to suppose, at an early period of the investigation of the electrical powers of metals, that they were entirely the result of chemical changes: that as heat was produced by this action, when exerted under common circumstances, so electricity resulted from it under other circumstances; and many of the phenomena were conformable to such an idea, and some ingenious inquirers adopted it to such an extent, as to suppose electricity in all cases owing to this cause.

This generalization, whether applied to Voltaic or to common electricity, seems, however, to be incorrect. Zinc and copper, as has been stated, different metals and oxalic acid, different metals and sulphur, or charcoal, exhibit electrical effects after mere contact, and that in cases when not the slightest chemical change can be observed; and if in these experiments chemical phenomena are produced by the action of menstrua, all electrical effects immediately cease: and it is not philosophical to assume a cause to account for an effect, when no such cause can be perceived.

It has been supposed that the action of the common electrical machine depends upon the oxidation of the amalgam; but I found by mounting a small machine in a glass vessel, in such a manner that it could be made to revolve in any species of gas, that it was active in hydrogen gas, and more active in carbonic acid gas than in the atmosphere (probably owing to its greater

density). The experiment has been several times repeated under different circumstances, and uniformly with the same results; and may be regarded as decisive in this important question.

33. Electrical effects are exhibited by the same bodies, when acting as masses, which produce chemical phenomena when acting by their particles; it is not therefore improbable, that the primary cause of both may be the same, and that the same arrangements of matter, or the same attractive powers, which place bodies in the relations of positive and negative, i. e. which render them attractive of each other electrically, and capable of communicating attractive powers to other matter, may likewise render their particles attractive, and enable them to combine, when they have full freedom of motion.

It is not a little in favour of this hypothesis, that heat, and sometimes heat and light, result from the exertion of both electrical and chemical attractive powers; and that by rendering bodies, which on contact are in the relation of positive to others, still more highly positive, as has been stated, page 116, their powers of combination are increased; whereas, when they are placed in a state corresponding to the negative electrical state, their powers of union are destroyed. That acids can be detached from alkalies, oxygen and chlorine from inflammable matter by metallic substances, or by a fluid menstruum highly positive, is likewise favourable to the supposition.

34. This view of the possibility of the dependence of electrical and chemical action upon the same cause, has been much misrepresented. It has been supposed that the idea was entertained, that chemical changes were occasioned by electrical changes; than which

nothing is further from the hypothesis which I have ventured to advance. They are conceived, on the contrary, to be *distinct* phenomena; but produced by the *same power*, acting in one case on masses, in the other case on particles. The hypothesis has been attempted to be controverted by experiments which are far from satisfactory, and some of which have no connection with it. It has been said that acids rendered positive by the common machine, will still combine with alkalies, and that other contradictory results may be obtained; but a non-conducting acid, though brought in contact with a positive surface, electrified by the common machine, is not rendered positive throughout; but gains a polar electricity, which extends only to a certain depth into the crystals, and the exterior surface, if electrical at all, is negative: and if a wire, positively electrified by the common machine, be introduced into an acid solution, this solution, if at all affected, when made to act upon another solution, will be negative at its point of action; that is, it will be positive near the wire, but will be in the opposite state with regard to another surface. And common electricity is too small in quantity, in its usual form of application, to influence chemical changes; for it requires a very strong machine acting upon a very small surface, to produce any sensible polar decompositions of bodies.

35. The power of action of the Voltaic apparatus, seems to depend upon causes similar to those which produce the accumulation in the Leyden battery, namely, the property of non-conductors and imperfect conductors to receive electrical polarities from, and to communicate them to conductors; but its permanent action is connected with the decomposition of the chemical menstrua between the plates. Each plate of

zinc is made positive, and each plate of copper negative, by contact; and all the plates are so arranged with respect to each other as to have their electricities exalted by induction, so that every single polar arrangement, heightens the electricity of every other polar arrangement; and the accumulation of power increases with the number of the series. When the battery is connected in a circle, the effects are demonstrated by its constant exhibition of chemical agencies, and the powers exist as long as there is any menstruum to decompose: but when it is insulated, and the extreme poles of zinc and copper are unconnected, no effects whatever are perceived to take place, no chemical changes go on, and it exhibits its influence only by communicating very weak charges to the electrometer, the end terminated by zinc communicating a positive charge, that terminated by copper, a negative charge.

That each plate of the most oxidable metal in the apparatus, is in the relation of positive, and each plate of the least oxidable, in the relation of negative, and that every series is possessed of similar and equal polarity, is shewn by a very simple experiment; forty rods of zinc of the same size, connected with forty silver wires precisely similar, were introduced in the regular order into similar glasses filled with a solution of muriate of ammonia, rendered slightly acid by muriatic acid; as long as the extreme parts remained unconnected, no gas was disengaged from the silver, and the zinc was scarcely acted upon; when they were connected, all the plates of zinc were dissolved much more rapidly, and hydrogen gas was evolved from every silver wire. And in another experiment, in which several of these wires at equal distances were introduced

into small glass tubes, it was found that equal quantities of hydrogen were produced.

36. It seems absolutely necessary for the exhibition of the powers of the Voltaic apparatus, that the fluid between the plates should be susceptible of chemical change, which appears to be connected with the property of double polarity, of being rendered positive at one surface, and negative at the other. There are substances that are imperfect conductors, which are capable of receiving only one kind of electricity, when made parts of the Voltaic circuit, and which M. Eberman, who discovered them, has named *unipolar* bodies. Perfectly dry soap, and the flame of phosphorus, when connected with the two extremities of the Voltaic apparatus, and with the ground, discharge only the negative electricity. The flames of alcohol, hydrogen, wax, and oil, discharge under like circumstances only the positive electricity; but all these bodies when connected with one pole only of the pile, and with the ground, destroy the divergence of the leaves of the electrometer connected with that end. It is not difficult to exhibit these phenomena when the atmosphere is dry, by means of two hundred pair of plates carefully insulated: an insulated gold leaf electrometer having a moveable wire attached to it, should be connected with each end of the pile: when either electrometer is brought in contact with soap, the soap being connected with the ground, the slight divergence of the leaves will cease; when the soap is connected with both electrometers and with the ground, the divergence of the leaves of the electrometer connected with the end terminated by the zinc, will continue, the leaves of the other electrometer will collapse. The opposite effect

occurs when the flame of a taper is connected with both electrometers and with the ground.

The unipolar conductors are incapable of being active in any part of the pile, and in this respect agree with non-conductors; many of which, it is probable, if examined in their relations to electricities of low intensity, would exhibit similar differences.

37. There are no fluids known, except such as contain water, which are capable of being made the medium of connection between the metals, or metal of the Voltaic apparatus; and in cases in which Voltaic batteries have been said to be constructed by metals and paper, or metals and starch, or other like substances, the feeble effects produced are merely owing to the small quantity of water adhering to these substances, which will not act when carefully dried. The instrument, called by M. de Luc, the electrical column, formed of zinc, Dutch leaf, and paper, and which he appears to consider as a different combination from the pile of Volta, seems to be merely a feeble Voltaic apparatus, in which the quantity of electricity is not sufficiently great to produce any chemical changes, or distinct phenomena of ignition; but in which the intensity of the small quantity existing, when the combination amounts to 400 or 500, is sufficient to enable it to affect the electrometer, and to act through a plate of air.

It is very probable that the power of water to receive double polarities, and to evolve oxygen and hydrogen, is necessary to the constant operation of the connected apparatus; and that acids, or saline bodies, increase the action, by affording elements which possess opposite electricities to each other, when mutually excited; the action of the chemical menstrea exposes continually

new surfaces of metal; and the electrical equilibrium may be conceived in consequence, to be alternately destroyed and restored, the changes taking place in imperceptible portions of time.

The manner in which aqueous fluids receive and communicate electrical polarity, is shewn by a very simple experiment; let a number of fine metallic surfaces, or flattened wires (of tin for instance) be made to swim in a narrow trough containing water; and let two wires from the extremities of a Voltaic battery of 1000 double plates, be plunged into the remote ends of the trough, one into one end, the other into the other end. The metals swimming on the water will immediately acquire electrical polarity; and the positive and negative poles will be regularly opposed to each other, the pole of the metal opposite to the wire positively electrified, will be found to be negative, giving off hydrogen, the other pole will deposit oxide; the next wire to this will present the alternate order, which will be preserved in all of them; those most remote from the right line of the circuit, will be least affected. If the battery be in a highly active state, the different wires will attract each other by their opposite poles, and the circle will at length be closed with the production of brilliant sparks. The phenomena are precisely analogous to those phenomena in *magnetism*, presented by a number of flattened wires of soft iron, made to swim upon water, and rendered magnetic by the opposite poles of two powerful magnets; each wire has a north pole and a south pole; and in the alternation, the different poles are attractive of each other.

38. That the decomposition of the chemical agents is connected with the energies of the pile, is evident from all the experiments that have been made; as yet no

sound objection has been urged against the theory that the contact of the metals destroys the electrical equilibrium, and that the chemical changes restore it; and in consequence that the action exists as long as the decompositions continue; and this conclusion is confirmed by the late researches made by MM. Gay Lussac and Thenard, on the great pile constructed by order of the French government. The manner in which chemical changes tend to restore the electrical equilibrium, is shewn by a remarkable experiment on the electrization of mercury, which I have very lately made. A few globules of mercury are placed in a vessel containing common pump water; or any water that contains a small quantity of saline impregnation; wires from a battery of 1000 double plates, not very strongly charged, are introduced into the vessel opposite to each other, so as to reach the bottom; as soon as the circle is completed, the mercury will be violently agitated, each globule will become elongated towards the positive pole, but will retain its circular outline in the part opposite to the negative pole; oxide will be given off from this part, which is positive, but no hydrogen from the part which is negative, and the oxide will pass in a rapid current from the positive towards the negative pole. As long as no hydrogen is given off, the globule is in continued agitation, and a stream of oxide flows with great rapidity from the positive to the negative surfaces; and the negative surfaces of the mercury approach rapidly towards the positive, which are at rest; if the conducting power of the water is exalted by the addition of more of the saline impregnation, or if the charge of the battery be increased, hydrogen will be given off from the negative poles; and the instant this happens the globules become sta-

tionary; as if the same power which gave motion to the mercury was neutralized by, or employed in, the evolution of the hydrogen. There are many other remarkable phenomena connected with the operation of electricity on mercury, in contact with water; which may be urged in favour of the idea, that chemical and electrical attraction depend upon the same cause, and which will possibly lead to new views respecting the elements of matter; but the consideration of them properly belongs to a more advanced division of this work.

39. The illustrious inventor of the new electrical apparatus, has given it the name of the electromotive apparatus, and has founded his theory of its operation upon the Franklinian idea of an electrical fluid, for which certain bodies have stronger attractions than others: and he conceives, that in his pile the upper plate of zinc attracts electricity from the copper, the copper from the water, the water again from the next plate of zinc, the next plate of zinc from the next plate of copper, and so on.

This hypothesis applies very happily to most of the phenomena of the action of the insulated pile, and the pile connected by either of its extremities with the ground; but does not explain with the same facility, the powers of the apparatus connected in a circle, in which each plate of zinc must be supposed to have the same quantity of electricity as each plate of copper; for it can only receive as much as the copper can give, unless indeed the phenomena of the circular apparatus be considered as depending upon the constant and rapid circulation of the natural quantity of electricity, in the different series; which requires the proof of a constant power in a substance to attract electricity from

one body, at the same time that it is giving it off, to another.

40. Whatever may be the happiest approximation to the true theory of the Voltaic instrument, it can scarcely be doubted that the electrical organs of certain animals depend upon similar arrangements of exciting bodies. The shock of the *Gymnotus Electricus*, and the *Torpedo*, resemble the Voltaic shock; and the power resides in organs which consist of a number of similar alternations of different substances. The effects are analogous to those which a Voltaic apparatus of small surface, consisting of very numerous but not very powerful series, would produce. It has been conceived that other phenomena of living action may be connected with the operation of weak electrical powers; such as secretion; and some ingenious hints on this subject have been advanced by Dr. Wollaston and Sir Everard Home, and some experiments relating to the subject instituted by Mr. Brande. Such inquiries are worthy of further pursuit, as they may tend to elucidate some important functions of the animal economy; but they must not be confounded with certain vague speculations, that have been advanced by some authors, on the general dependence of nervous or sensitive action, and muscular or irritable action, upon electricity; such speculations are mere associations of words derived from known phenomena, and applied illogically to unknown things. The laws of dead and living nature appear to be perfectly distinct: material powers are made subservient to the purposes of life, and the elements of matter are newly arranged in living organs; but they are merely the instruments of a superior principle.

As electrical changes are almost constantly taking

place in the atmosphere, and as the different substances composing the exterior of the globe, bear different electrical relations to each other, it is very probable that many of the chemical changes taking place on the surface, are influenced by the action of weak electrical powers: such as the decomposition of the surfaces of rocks, the modifications of soils, the formation of acid, and development of alkaline compounds; and the mutual agencies of the elements in the earth, the sea, and the atmosphere, may be assisted or modified by the circumstances of general electrical action.

41. With regard to the great speculative questions, whether the electrical phenomena depend upon one fluid, in excess in the bodies positively electrified, and in deficiency in the bodies negatively electrified, or upon two different fluids, capable by their combination of producing heat and light, or whether they may be particular exertions of the general attractive powers of matter, it is perhaps impossible to decide in the present imperfect state of our knowledge. The application of electricity as an instrument of chemical decomposition, and the study of its effects, may be carried on independent of any hypothetical ideas concerning the origin of the phenomena; and these ideas are dangerous only when they are confounded with facts. Some modern writers have asserted the existence of an electrical fluid with as much confidence as they would assert the existence of water, and have even attempted to shew that it is composed of several other elements; but it is impossible in sound philosophy to adopt such hasty generalizations; Franklin, Cavendish, Epinus, and Volta, the illustrious advocates for the idea of a single electrical fluid, have advanced it only as hypothetical, as accounting in a happy way for most of the pheno-

mena; and none of the facts that have been brought forward in favour of the actual existence either of one or of two fluids, can be considered as conclusive.

From a very ingeniously contrived experiment of Mr. Cuthbertson, it appears that when a stream of electrical sparks is passed through the flame of a candle between two electrified surfaces, the surface which is negative is most heated; and it has been argued that a current must pass from the positive surface to the negative.

But it is more probable, that this phenomenon depends upon the positive *unipolar quality* of the flame of wax or tallow referred to above; for supposing this flame to become positive, which would seem to be the case, it must be attracted by the negative, and not by the positive surface; and this view is confirmed by an experiment I made on an arch of flame between the two poles of the great Voltaic apparatus of 2000 plates. Platina melted with more facility in the arch at the positive than at the negative extremity, and this arch was common air intensely ignited, through which the electricity was discharged; and if any mechanical current existed from the positive pole to the negative, the maximum of heat must have been produced at the negative. When a wire of platina was made positive, and brought in contact with charcoal rendered negative, it became ignited much sooner, and fused into larger globules, than when made negative, and brought in contact with the charcoal rendered positive; and that the effect did not depend upon the greater heat of the charcoal, appears from the circumstance, that similar phenomena occurred when the experiment was made by contact with mercury. But when an imperfectly conducting fluid, such as sulphuric acid, was used, the result was reversed. The wire being negatively electri-

fied, and the acid positively, the point in contact with the surface of the acid instantly became white hot; in the opposite case a spark of blue light only was produced.

The different appearance of the light on points positively and negatively electrified, has been urged in favour of the idea of a fluid proceeding from the positive to the negative surface. This phenomenon occurs as well in the Voltaic, as in the common discharge: for when the arch of flame passes between two points of charcoal, a vivid spot of white light is always perceived on the negative point, and rays seem to diverge from the positive point. The effect of the difference of the appearance of differently electrified points, I find, does not depend upon the nature of the elastic medium, for it takes place in hydrogen, carbonic acid, and chlorine, though it is less distinct in the heavier gases, probably from their being worse conductors; but the affections of light in passing from the different parts of the circuit, can with no more propriety be urged in favour of a specific fluid, than the chemical changes produced by the different poles.

When folds of paper are perforated by a discharge from an electrical jar, there is a burr on both sides, which may be urged as an argument against any fluid passing through; for it could penetrate in one direction only, and the experiment is favourable to the idea that electricity is an exhibition of attractive powers acting in peculiar combinations, for the substance of the paper, which was negative, may be conceived violently attracted to the positive surface, and the part which was positive, to the negative, at the moment the discharge takes place.

It will be useless to pursue any further this recondite

part of the subject; whatever view is taken, active powers must be supposed to be bestowed upon species of matter, and the impulse must be ultimately derived from the same source. In the universe nothing can be said to be automatic, as nothing can be said to be without design. An imperfect parallel may be found in human inventions; springs may move springs, and wheels, indexes; but the motion and the regulation must be derived from the artist; sounds may be produced by undulations in the air, undulations of the air by vibrations of musical strings; but the impulse and the melody must arise from the master.

VIII. *On Analysis and Synthesis; on the Circumstances to be attended to in these Operations, and on the Arrangement of Undecomposed Bodies.*

1. When a substance is capable of being resolved into other forms of matter, it is said to be compounded; thus, if mild magnesia, (subcarbonate of magnesia) be strongly heated for an hour in a green glass retort, having its beak connected with a flaccid bladder, elastic matter will collect in the bladder; and the magnesia, when examined, will be found to have lost in weight, and to be altered in its properties, it will not effervesce with acids, and it is harsher to the feel. The weight of the elastic matter collected in the bladder is exactly equal to that lost by the magnesia; it cannot by any means be converted into magnesia, and the mild magnesia gives only a limited quantity of it; so it is evident that mild magnesia consists of a matter which can be rendered permanently gaseous, and a fixed substance; it is a *compounded body*.

The metal called zinc, if heated strongly in close

vessels, rises in the elastic form, but when condensed by cold, it appears unaltered in its properties. It may be distilled any number of times, but it will be still the same: nothing permanently elastic will be given off from it; and if the operations be conducted with care, it will be found undiminished in weight. Not even the intense heat of the Voltaic battery applied in a vessel exhausted of air, effects any change in it; it easily enters into new combinations, but can be resolved into no other forms of matter; it is considered as an *undecompounded* body.

The term *element* is used as synonymous with *undecompounded* body; but in modern chemistry its application is limited to the results of experiments. The improvements taking place in the methods of examining bodies, are constantly changing the opinions of chemists with respect to their nature, and there is no reason to suppose that any real *indestructible principle* has been yet discovered. Matter may ultimately be found to be the same in essence, differing only in the arrangements of its particles; or two or three *simple* substances may produce all the varieties of compound bodies. The results of our operations must be considered as offering at best approximations only to the true knowledge of things, and should never be exalted as a standard to estimate the resources of nature.

2. By *analysis* compounded bodies are resolved into their constituents; by *synthesis* they are produced in consequence of the union of these constituents; and when the weight of the compound corresponds to that of the constituents, the processes are considered as accurate.

The words analysis and synthesis are applied in cases when bodies are resolved into, or compounded from, any

other forms of matter, without relation to the elementary nature of these forms;—thus crystals of Glauber's salt may be resolved analytically into sulphate of soda and water, or compounded synthetically from these substances; and sulphate of soda may be formed by synthesis from sulphuric acid and soda, both of which are compounded bodies.

3. In all conclusions upon the results of analytical and synthetical experiments, it is of the greatest importance that the agencies of all the substances concerned, should be accurately known, that no circumstance should be taken for granted, and that the nature of the real constituents of the body should be shewn to be unchanged during the process.

Whatever instruments of experiments be used, their relations to the substances acted upon should be well known, and their influence (if any) estimated. Thus if a hard stone be pulverized in a mortar of porcelain, agate, or iron, the comparison of weights before and after the process should be carefully made, to ascertain what quantity of matter may have been abraded from the mortar. When substances are fused, or heated, in vessels on which they are capable of acting, the same precautions should be taken. It should either be shewn, that the vessel has been unchanged during the operation, or the nature and extent of the change should be demonstrated.

Many celebrated chemists have been led into error in the infancy of their investigations, from a want of attention to these circumstances. Thus the illustrious Scheele for some time supposed that siliceous earth was composed of fluoric acid and water, because he obtained it by mixing together an acid gas, (procured from fluor spar) and water; but subsequent experiments, by de-

monstrating the loss of weight of the glass vessels in which his operations were conducted, shewed that the matter which produced the siliceous earth was derived from these vessels.

4. Water is the great solvent employed in chemical processes, and its operation therefore should be strictly attended to. It has been too much the custom to consider its elements as almost passive in the processes of dissolution and decomposition; but there are a number of instances in which these elements are newly arranged, and in which their transfer and changes produce very important phenomena.

When oxymuriatic or chlorine gas is exposed to light, it undergoes no change, but when a solution of it in water is placed under the same circumstances, oxygen gas is given off, and a solution of muriatic acid is found in the water. Hence it was concluded, without any reference to weights, that oxymuriatic gas consists of muriatic gas and oxygen, and that the water acted in no other way than in assisting the expulsion of the oxygen, by its attraction for the muriatic acid gas.

This inference, however, is now known to be incorrect, and it affords a striking instance of the present object of discussion. If aqueous vapour in small quantities, and chlorine gas, be passed through an ignited tube of glass, the steam entirely disappears, and oxygen gas, and muriatic acid gas are formed; therefore the water must have entered into the composition of the muriatic acid gas, or must have been decomposed; its hydrogen combined with the chlorine, to form muriatic acid gas, and the oxygen gas set free; and that hydrogen actually enters into the composition of muriatic acid gas, is proved by the experiment, which will be detailed hereafter; nor can any oxygen gas be procured

in any experiments upon chlorine, in which bodies not known to contain oxygen alone are concerned; nor have any means been found by which this substance can be decomposed.

To give another example: when concentrated oil of vitriol, which consists of sulphuric acid and water, is poured upon common salt, and they are heated together, muriatic acid gas flies off, and sulphate of soda is obtained; hence it was concluded, that common salt consists of muriatic acid gas and soda; and that the sulphuric acid merely displaced the muriatic acid gas; and no account was taken of the water of the sulphuric acid in the operation; yet the whole change depends upon this water: and no soda and no muriatic acid can be procured from common salt, without water; and common salt is made directly by heating sodium, the metal which I discovered to be the basis of soda, and chlorine together, and these are both as yet undecomposed bodies; and if 92 parts of oil of vitriol, which consists of 75 parts, by weight, of sulphuric acid, and 17 parts of water, be made to act upon 111 parts of common salt, which consists of 44 sodium and 67 chlorine, the water will be decomposed, 15 of oxygen will combine with the sodium to form 59 of soda, and 2 of hydrogen will combine with 67 of chlorine to form 69 of muriatic acid gas, and the sulphate of soda will be 134 parts.

5. There are many substances which possess an attraction of a peculiar kind for water; they absorb water without undergoing any remarkable change in their properties, and in small proportions. Such are charcoal, different earths, and animal and vegetable substances. If well-burnt charcoal be exposed to the atmosphere for some days, it will increase in weight from 10 to 14

grains per cent., and the increase is almost entirely owing to its absorbing water, which existed in the form of vapour in the air; and by heating charcoal that has been exposed to air, in close vessels, the water may be collected unaltered. Baryta, strontia, and lime, absorb definite proportions of water, and form what are called hydrates, in which the water is in chemical combination, and requires an intense heat for its expulsion. And magnesia, alumina, silica, glucina, and zircona, likewise increase in weight by attracting aqueous vapour from the atmosphere, and seem to form analogous combinations; but in which the water is less powerfully retained; it is expelled at a temperature of dull ignition.

Compounds of the earths in fine powder, that have been heated red, increase in weight, from the absorption of atmospheric moisture: and the case is the same with almost all substances, except the metals, and certain inflammable bodies; so that in all experiments of analysis, the solid products obtained should be strongly heated, and weighed whilst warm, and before they have been long exposed to the atmosphere, or the quantity of water they have absorbed should be exactly known. And the same precautions should be used, and more strictly, with respect to alkaline, acid, and saline bodies that enter into chemical combination with water, and attract it rapidly from the atmosphere.

6. Gaseous bodies are usually procured from substances that contain water, and many of them are collected over water; it is therefore of considerable importance in analytical processes, that their relations to this substance should be distinctly understood.

It has been already stated, that common air contains aqueous vapour, or water in an invisible elastic form, which is greater in proportion as the temperature is

high, air at the temperature of 65° Fahrenheit, containing about $\frac{1}{50}$ of its volume. From the experiments of Desormes and Clement, it appears that all the gases not absorbable to any extent by water, such as oxygen, azote, carbonic acid, and hydrogen gases, contain in equal volumes, the same quantity of vapour as common air; so that vapour, when it exists in these bodies, can only be regarded as mixed with them; and it is separated by substances that have a strong chemical attraction for water, such as lime, muriate of lime, sulphuric acid, hydrate of potassa, &c.; and in all accurate experiments in which gases are examined, they should be previously freed from vapour by exposure for some hours to substances that have a strong attraction for water, but possessed of no chemical action on the gas.

The relations of water to gases with which it is capable of combining chemically, (which gases will be described hereafter,) are very different. It is evident that no pure aqueous vapour can exist in them in a state of *mixture*, but they may, and probably in almost all cases do contain a gaseous compound of water, and the peculiar elastic fluid. If a drop of water be introduced into a flask filled with ammoniacal gas, it rapidly absorbs the gas, and increases in size; but if a minute drop of a concentrated solution of ammonia be introduced, and the temperature of the flask be gently raised, the drop disappears, and continues invisible, as long as the heat is preserved uniform. The instances are similar when analogous experiments are made upon muriatic acid and silicated fluoric acid gases; and I have found that these elastic fluids collected at the temperature of 75° deposited a slight dew, consisting of strong solution of acid in water, when intensely cooled by a freezing mixture. There is reason to believe that

the case must be the same with fluoboric acid, and that this body may contain a minute quantity of the compound which may be called hydrate of fluoboric acid; and this is confirmed by the phenomena of the action of potassium upon the gas, for I have never been able to decompose it by this substance, without procuring small quantities of hydrogen.

The quantity of water in the gases for which it has a chemical attraction, must depend upon the degree of volatility of the fluid compound of the gas and water, and upon the proportion of water it contains. Sulphureous acid gas, which has only a weak attraction for water, would, there is every reason to believe, contain most of the gaseous hydrate; but even in this it is most likely there must be less water than in common air at the same temperature; ammonia would probably be next in order, then silicated fluoric gas, muriatic acid gas, nitrous acid gas, and, last of all, fluoboric gas.

The temperatures at which the compounds of water and gases rise in vapour, seem to depend upon the strength of the attraction, by which they are combined, and upon the degree of volatility of the gaseous element. All solutions of sulphurous acid, and ammonia, boil at temperatures which differ very little from the boiling point of water. The highest point of the ebullition of solution of muriatic acid gas in water is about 232° Fahrenheit; that solution of nitric acid which gives a compound vapour, does not boil at a temperature below 248° ; the temperature at which hydrated fluoric acid boils, according to MM. Gay Lussac and Thenard, is not very high; but the vapour contains a considerable quantity of water, compared with other acid vapours.

Whether substances will attract water from the absorbable gases, must depend upon the strength of their

affinity for water, as compared with that of the gas. Dry hydrate of potassa will slowly attract moisture from ammonia, and dry muriate of lime from sulphurous acid gas; but muriate of lime does not appear to act upon the water in muriatic acid gas. Silicated fluoric acid gas and fluoboric gas, instantly render cloudy sulphurous acid gas by attracting moisture from it, and fluoboric gas, if mixed with silicated fluoric acid gas, renders it very slightly cloudy. Probably there are no known substances which will attract water from the vapour of hydrated fluoboric acid; but the quantity is too minute to influence to any extent, the results of experiments on gases containing it.

In cases when elastic fluids are produced in contact with substances which afford peculiar vapours, such as volatile oils, alcohol, ether, &c.; these vapours should be separated either by agitating the gases in water, or solutions of substances which are capable of absorbing them, such as solution of potassa, &c. and the aqueous vapour separated afterwards by the means above mentioned.

7. In stating the weights of bodies which are the result of analytical experiments, the temperature should be noticed; and in the case of elastic fluids, the degree of pressure of the atmosphere, as indicated by the barometer. When gaseous compounds are resolved into simpler gaseous bodies, or when gases are compared with each other, as they are all similarly affected by heat and pressure, there is no necessity for any specific statements of these circumstances, and in describing the specific gravity of a gaseous body, it is necessary only to give the relation of its weight to that of air; thus the weight of air being 1000, that of oxygen gas will be 1097. As hydrogen gas is much

lighter than any other elastic fluid, and as it is the body which combines with other substances in the smallest proportions, it would perhaps assist the progress of chemical inquiry to denote its specific gravity, by unity, which would harmonize with the idea of representing the proportion in which it combines likewise by unity, and would facilitate the means of comparing the absolute weights of gaseous bodies concerned in experiments, with the numerical symbols representing their elements. The specific gravity of hydrogen being considered as 1, that of common air will be 13·7, and that of oxygen, as has been stated in page 81.

8. In treating of the different substances which, by their agencies, combinations, or decompositions, produce the phenomena of chemistry—*radiant* or *ethereal matters* will be first considered, as their principal effects seem rather to depend upon their communicating motion to the particles of common matter, or modifying their attractions, than to their actually entering into combination with them; and as from the laws of their motions, or from their extreme subtileness, they are incapable of being weighed.

The *undecomposed* substances which are permanent in their forms, will be considered in an order of arrangement depending upon their electrical relations; *those* determined to the *positive surface* in the Voltaic electrical circuit, being arranged in one class, and *those* determined to the *negative surface* in another; and the subdivisions of the classes will be made according to their natural relations.

The general principle adopted will be, that no compounded body shall be treated of, till its constituents have been described.

The relations of bodies derived from their electrical

powers, are coincident with those dependent upon their agencies in combustion; that is, one class contains supporters of combustion, and the other class combustible bodies; but as the heat and light produced in combustion seem to be merely indications of the strength of attraction of the acting substances; and as these phenomena occur in cases in which inflammable matters act upon each other, combustibility can scarcely be considered as a definite idea; though the importance of the common phenomena of combustion have made them the grand objects in all the early theories of chemistry.

DIVISION II.

OF RADIANT OR ETHEREAL MATTER.

I. *Of the Effects of Radiant Matter, in producing the Phenomena of Vision.*

1. The phenomena of vision depend upon the presence of the sun, of the heavenly bodies, or on the mutual action of certain substances on the surface of the earth.

2. When light is entirely intercepted by a body placed between the luminous object and the eye, that body is said to be opaque; and the manner in which the light is intercepted, proves that it proceeds in right lines or rays from that luminous body.

3. Luminous objects may be seen through certain substances; and these bodies are said to be transparent. Bodies differ considerably in the degree of their transparency; some transmit many more rays than others, and there are gradations from perfect opacity, when all the rays are intercepted, to a higher degree of trans-

parency, when by far the greater number are transmitted.

4. Of the rays that are not transmitted, some are lost, as it were absorbed by the body; others are thrown back again, or reflected from its exterior or inner surfaces, and these rays are called reflected rays.

5. The rays of light in their transmission through bodies, or reflection from their surfaces, undergo certain modifications, of great importance in their connection with the laws of vision, and the general properties of radiant matter.

6. If rays of light pass from one transparent substance not crystallized into another, in an oblique direction, their path is altered, and they are bent downwards or upwards, according as the medium is more or less dense, or according as it differs in chemical qualities; inflammable substances, or compounds containing inflammable substances, having the highest power of bending towards the perpendicular or of refracting, as it is called, the rays of light; and, in the same substances, the sines of the angles of refraction bear always the same relations to those of the angles of incidence.

7. The rays of light in passing through obliquangular crystalline bodies, follow different laws. If a ray of light be received perpendicularly upon a plain surface of island crystal, or rhomboidal carbonate of lime, one part of it passes through without altering its direction; another part, on the contrary, is refracted in a plane parallel to the diagonal, joining the two obtuse angles of the crystal, so that images seen through the crystal appear double; and this phenomenon, first scientifically reasoned upon by Huygens, is called the phenomenon of double refraction.

If a ray of light which has suffered double refraction

from one crystal, be received by another crystal placed in a similar and parallel position, there will be no new division of rays, and no change in their direction; but if the second crystal be placed, so that its planes of perpendicular refraction are at right angles to those of the first crystal, then there will be a new phenomenon, and that part of the ray which before passed through the ordinary refraction, will receive the extraordinary refraction, and reciprocally that which underwent the extraordinary will suffer the ordinary refraction. If the second crystal be turned gradually round in the same plane, when it has made a quarter of a revolution, there will be four divisions of the ray, and they will be reduced to two in the half of the revolution, so that the refracting power depends upon the relations of the position of the particles of the crystals, to the rays passing through them.

Similar phenomena to those presented by island crystal, are exhibited in a greater or less degree by other crystalline bodies, and probably would belong to all of them, if they were sufficiently transparent to admit of the passage of light through strata of considerable thickness. Very thin pieces of the rhomboidal carbonate of lime even do not give perceptible, double images.

8. When light is reflected from bodies; under most circumstances it is unaltered in its relations to the refractive powers of transparent substances, and the angle of reflection is equal to the angle of incidence. But, in certain cases, at angles which differ for different bodies, the reflected rays have the same property as the extraordinary refracted rays, that have passed through island crystal. This important fact, discovered by M. Malus, is easily exemplified. If the flame of a taper reflected at an angle of $52^{\circ} 45'$ from the surface of water, be viewed

through a piece of double refracting spar, one of the images will vanish every time that the crystal makes a quarter of a revolution.

If a ray of light be reflected from a surface of glass, at an angle of $54^{\circ} 35'$, and fall upon another plane of glass at the same angle, it will suffer no new reflection, and will pass through the glass unaltered, provided that the planes of reflection or refraction be perpendicular to each other; but if they are in the same direction, nothing remarkable will happen.

Direct light is most copiously reflected, as its incidence is less perpendicular; but light once reflected follows different laws, and the quantity that suffers the second reflection, depends entirely upon the relations of the angles made by the reflecting surfaces with the rays.

9. When a ray of light is made to pass through a triangular prism of glass, and the transmitted light is received upon a sheet of white paper, it is found to be of different colours; the most distinct of which are red, orange, yellow, green, blue, indigo, and violet. Newton has ascertained, that if the coloured image, or spectrum, as it is called, be divided into 360 parts, the red will occupy 45 of these parts, the orange 27, the yellow 48, the green 60, the blue 60, the indigo 40, and the violet 80. The red rays are least refracted, the violet rays most, and the other coloured rays are refrangible inversely, in the order in which they have been named.

According to Dr. Wollaston, when the beam of light is only $\frac{1}{20}$ of an inch broad, and received by the eye at the distance of ten feet, through a clear prism of flint glass, four colours only appear—red, yellowish green, blue, and violet.

If the differently coloured rays of light separated by

the prism, be concentrated upon one spot by means of lenses, they produce white light; and Newton has beautifully explained the different colours of bodies, by supposing that they retain certain of the coloured rays of light, and reflect others; thus red bodies are supposed to reflect red light, and to absorb all the other coloured rays.

The different coloured rays of light, as has been shewn by Sir Wm. Herschel, differ in their power of rendering objects visible; at least in the state of division, which is obtained by means of a prism. If an equal portion of these rays be made to illuminate a printed page, the words may be seen from the greatest distance, when exposed to the lightest green or deepest yellow light; and the effects of illumination for equal quantities of the rays, diminish from the central parts towards the extremities of the spectrum. It may, however, be said that there are *more* green rays in a given part of the spectrum than blue rays, and the difference of illuminating power may depend on this circumstance.

10. The rays separated by one prism are not capable of being further divided by being passed through another; and in their relations to double refraction and reflection, they appear to agree with direct light; an object illuminated by any of the rays in the spectrum, is seen double through island crystal, in the same manner as if it had been visible by white light.

11. The minute investigation of the properties of radiant matter, in their relations to the phenomena of vision, constitutes the object of a particular branch of science—*Optics*. The few statements that have been made on this subject, it will be found in the following pages, are connected with the chemical effects and nature of radiant matter; and it will immediately be

seen, that the same causes which produce the most numerous and important of our sensations, and which give, as it were, language to the external world, are likewise subservient to the orderly succession of events in the economy of nature.

II. *Of the Operation of Radiant Matter in producing Heat.*

1. When similar thermometers are placed in the different parts of the solar beam, separated by the prism, it is found that different effects are produced in the different coloured rays. The greatest heat is exhibited in the red rays, the least in the violet rays; and in a space beyond the red rays, where there is no visible light, there is still an increase of temperature. The important discovery of the production of heat where there is no visible light was made by Sir Wm. Herschel:* he estimates the power of heating in the red rays, to be to that of the green rays as 55 to 26, and to that of the violet rays as 55 to 16; and he states that beyond the red rays the increase of temperature is greater than in the red rays in a ratio as 9 to 7.

It is evident, however, that these results must differ according to the manner in which the experiment is made, or the nature of the apparatus and the size of the thermometers used. M. Prevost has obtained results which differ from those of Sir Wm. Herschel as to the ratio of the intensity of the different rays, and M. Berard fixes the maximum of heat in the solar spectrum within the extreme red rays, and not in the space where there is no visible light. In experiments which I saw

* Philosophical Transactions, 1800, p. 261.

made by Sir Henry Englefield,* the greatest effect was produced when the largest part of the ball of the thermometer was entirely out of the spectrum.

2. From these facts it is certain that matter proceeding from, or set in motion by the sun, has the power of producing heat without light, and that its rays are less refrangible than the visible rays.

Some persons have concluded from the phenomena, that all the rays which produce heat in the solar beam, are distinct from those that produce light; but this does not seem to be warranted by the experiments, for if it were the case, they would, probably, be entirely separated from the coloured rays by the prism, as the coloured rays are from each other. It has been used as an argument, in favour of the distinctness of the rays producing light, and those producing heat, that the beams from the moon illuminate without heating; but it is possible that a greater number of the most heat-making rays, than of the other rays, may be absorbed by that planet: and supposing *all* the rays reflected that fall upon the moon, yet still their intensity would be 95,890 times less than that of the solar rays, at the surface of the earth: and it appears from experiment, that the real intensity of the light of the moon to that of the sun is less than 1 to 300,000; such rays concentrated by the most powerful lenses, could not be expected to produce any effect on common thermometers; and as yet no very delicate experiments have been published on the subject.

* In these experiments the chamber was not darkened; hence probably the extreme red rays were not *visible*. In M. Berard's experiment a helioscope was used, and the room darkened; and I am inclined to put more confidence in these results. It is, however, probable that if the temperature of a *very minute* space beyond the red could be measured, the elevation would be greatest in that space.

3. The invisible rays that produce heat, are capable of reflection as well as refraction, in the same manner as the visible rays. It is well known that an intense degree of heat may be produced by the concentration of the solar beams upon a single spot, by a concave mirror, or by several mirrors; and there is no reason to disbelieve the *possibility* of the inventions ascribed to Archimedes, who, it is said, by the combined effect of a number of plane mirrors, set fire to the Roman ships during the siege of Syracuse; though the immense means and labour required for such an operation, render the narrative very doubtful.

4. Rays capable of producing heat with and without light proceed from bodies at the surface of the globe under peculiar agencies or changes, as well as from the sun; and the phenomena that are usually called the phenomena of the *radiation of terrestrial heat* are of great extent and importance, and well worthy of being studied.

5. If a thermometer be held near an ignited body it receives an impression connected with an elevation of temperature: this is partly produced by the conducting powers of the air; but it is likewise partly produced by another impulse which is instantaneously communicated, even to a considerable distance. If a large concave metallic mirror be placed upon the ground, and the hot body be raised some feet above it, a thermometer will instantly rise in the focus of the mirror, though it is evident that no current of hot air can pass downwards from the body.

This effect is commonly called the radiation of terrestrial heat. It is best observed, by employing two mirrors parallel to each other, and to the surface of the earth. If the mirrors be of copper tinned, and be well polished, and as much as 2 feet in diameter, and distant

only 12 feet, a small pan of red hot charcoal, placed in the focus* of the upper mirror, will cause gunpowder to explode in the focus of the lower mirror.

6. In cases when no light is emitted from a hot body, the effect of the concentration of heat by the mirrors may still be produced. Thus, if a vessel filled with boiling water, be placed in the focus of the upper mirror, a thermometer placed in the focus of the lower one will have its temperature increased.

These phenomena of the radiation of terrestrial matter producing heat, were made known by the academicians Del Cimento, Hooke, Scheele, and Pictet: and there is another fact, still more extraordinary, which has been called *the radiation of cold*, first observed by the Italian philosophers, and afterwards by Pictet. If in the arrangement of the two parallel mirrors, a piece of ice be introduced into the lower focus, the thermometer in the upper focus will indicate a diminution of temperature.

7. All these phenomena coincide with the phenomena of the reflection of the solar beams; and even the *apparent radiation of cold* is in harmony with them; for if it be supposed that rays capable of producing heat, emanate from all terrestrial bodies, but in quantities greater in some increasing proportion as their temperatures are higher, then the introduction of a cold body into the focus of one mirror, ought to diminish the temperature of a thermometer in the focus of the other, in the same manner as a black body placed in one focus, would diminish the quantity of light in the other focus; and

* In the usual form of the experiment, the mirrors are placed opposite to each other on the ground. This arrangement, which I have been long in the habit of employing, in the demonstrations in the Theatre of the Royal Institution, more distinctly exhibits the effect. See Plate V. fig. 20.

the eye is to the rays producing light, a measure, similar to that which the thermometer is to rays producing heat.

8. If a large lens be placed before an ignited body, and the light be refracted to a focus, a thermometer placed in the focus will be very slowly affected, and the increase of temperature will be infinitely less than that produced in the focus of a metallic mirror of the same size. Likewise, if glass screens be placed between the two mirrors, in the experiments in which the ignited coals, or hot water, or ice, are used, the effect is in great measure destroyed. This establishes a difference between the agency of the radiant matter producing heat on the surface of the earth, and of that from the sun. Mr. Leslie has supposed that the phenomena of the radiation of terrestrial heat, depend upon certain pulsations or undulations of the atmosphere capable of being reflected, but not of being refracted; but none of his facts prove this ingenious hypothesis. I had an apparatus made, by which platina wire could be heated in any elastic medium, or in vacuo; and by which the effects of radiation could be distinctly exhibited by two mirrors, the heat being excited by a Voltaic battery. In several experiments, in which the same powers were employed to produce the ignition, it was found that the temperature of a thermometer rose nearly three times as much in the focus of radiation, when the air in the receiver was exhausted to $\frac{1}{120}$, as when it was in its natural state of condensation.* The cooling power, by contact of the rarefied air, was much less than that of the air in its common state, for the glow of the platina was more intense in the first case than in the last: but this circumstance does not render the experiment less

* See Plate VI. fig. 21.

conclusive, and the results are entirely opposed to the idea that the terrestrial radiation of heat is dependent upon any motions or affections of the atmosphere.

9. It appears that the higher the temperature of the body radiating heat, the more readily the heat is transmitted through diaphanous solid or fluid media. I found that when a small lens was placed before the brilliant star of light produced by the discharge of the Voltaic battery of 2000 double plates by points of charcoal, and a thermometer placed in the focus of the lens it was instantly affected; though when the same lens was made to concentrate the rays from a candle, or the fire on the thermometer, there was no immediate effect produced. Dr. Delaroche has lately made a number of experiments on the radiation of terrestrial heat, and has deduced the following propositions from them.

First Proposition.—Invisible radiant heat may in some circumstances pass directly through glass.

Second Proposition.—The quantity of radiant heat which passes directly through glass is so much greater relative to the whole heat emitted in the same direction, as the temperature of the source of heat is more elevated.

Third Proposition.—The calorific rays which have already passed through a screen of glass, experience in passing through a second glass screen of a similar nature a much smaller diminution of their intensity than they did in passing through the first screen.

Fourth Proposition.—The rays emitted by a hot body differ from each other in their power of passing through glass.

Fifth Proposition.—A thick glass, though as much or more permeable to light than a thin glass of worse quality, allows a much smaller quantity of radiant heat

to pass. The difference is so much the less as the temperature of the radiating source is more elevated.

Sixth Proposition.—The quantity of heat which a hot body yields in a given time by radiation to a cold body situated at a distance, increases *cæteris paribus*, in a greater ratio than the excess of temperature of the first body above that of the second.

10. The manner in which the temperatures of bodies are affected by rays producing heat is different for different substances, and is very much connected with their colours. The bodies that absorb, as it is called, most light, and of course that reflect least, are most heated when exposed either to solar or terrestrial rays. Black bodies in general are more heated than red; red more than green; green more than yellow; and yellow more than white.

Metals are less heated than earthy or stony bodies, or than animal or vegetable matters. Polished surfaces are less heated than rough surfaces.

11. The bodies that have their temperatures most easily raised by the action of rays producing heat, are likewise those that are most easily cooled by their own radiation, or that at the same temperature emit most heat-making rays. Metals radiate less heat than glass: glass less than vegetable substances; and charcoal has the highest radiating powers of any body yet made the subject of experiment.

12. As the radiant matter producing heat is liable to reflection and refraction like the radiant matter producing light, so it is likewise susceptible of polarization. This has been proved by the accurate and ingenious experiments of M. Berard; who has shewn that the radiant heat from an ignited globe of metal is polarized and reflected at an angle which cannot differ consider-

ably from that at which the rays producing light are reflected.

13. Mr. Leslie has concluded from his researches on the radiating power of different bodies that the radiating power of finely divided charcoal (lamp black) being taken as 100; the following substances radiate in a proportion that may be thus expressed. Sealing wax 95; crown glass 90; China ink 88; ice 85; minium 80; isinglass 80; plumbago 75; tarnished lead 45; clean lead 19; polished iron 15; tin plate, gold, silver, and copper 12.

14. There are some practical applications of the doctrines of radiant heat, to the economy of some of the useful arts and processes of common life.

Vessels that are intended to retain their heat, should be metallic and highly polished; and, independent of elegance and delicacy, there is a reason obvious, from the preceding facts, why metallic vessels for the purposes of the table, should be kept as bright as possible. Steam or air pipes for warming houses, should be polished in those parts where the heat is not required to be communicated, and covered with some radiating substance, such as lamp black, or plumbago, in those rooms which are to be heated by them.

Culinary implements should be blackened, and not polished on those parts, which are to receive heat. The heated surfaces of fire-places or stoves should not be metallic; but of earthy or stony materials, and in this case much more heat will be communicated by radiation.*

* Count Rumford. Phil. Trans. 1804, page 178.

III. *Of the Effects of Radiant Matter in producing Chemical Changes.*

1. A number of the effects of radiant matter in producing chemical changes, may be ascribed to its powers of heating bodies. The heat produced by the concentration of rays has been already referred to, and the focus of a powerful lens or mirror exposed to the sun, offers means of exciting heat inferior only to those afforded by Voltaic electricity. In some cases the direct solar light produces effects similar to those produced by a degree of heat much higher than could be excited by their influence upon an opaque body, thus as M. Gay Lussac, Thenard, and Dalton have shewn, chlorine and hydrogen detonate when a mixture of them is exposed to the solar beams, though the same effect is not produced by the application of a heat below that of ignition. This has been explained by supposing that the temperature of *particles* of substances is raised *considerably* and *instantaneously* by the rays; but it may likewise, and with more probability, be supposed to depend upon a specific and peculiar influence of radiant matter, and that such an influence exists, is proved by many circumstances.

2. If moist horn silver (chloride of silver) be exposed to the different rays in the prismatic spectrum, it will be found that no effect is produced upon it in the least refrangible rays, which occasion heat without light; a slight discoloration only will be occasioned by the red rays; the effect of blackening will be greater towards the violet part of the spectrum; and in a space beyond the violet, where there is no sensible heat or light, the chemical effect will be very distinct.

This observation made by M. Ritter, and Dr. Wollaston, proves, that there are rays more refrangible than the rays producing light and heat; and from the observations of M. Berthollet, it appears, that muriatic acid gas is formed when horn silver is blackened by light.

3. It has been supposed that these peculiar rays are mixed with the other rays throughout the coloured part of the spectrum; but it seems equally probable, that the same rays that produce light, may likewise produce chemical effects, and effects of heat. Dr. Young has shewn that these invisible rays are liable to the same affections as visible rays, when they are reflected from thin plates of air, as in the phenomena of coloured rings; and M. Berard has proved that they are susceptible of polarization in the same manner as the rays producing heat and light.

As yet it has not been clearly demonstrated what is the peculiar chemical energy of the least refrangible rays. In acting upon the compounds of silver they produce new combinations or a disengagement of hydrogen by decomposing water; and they have no action whatever on these substances when dry.

4. I found that chlorine was much more rapidly absorbed by water, and muriatic acid much more rapidly formed when the vessels were exposed in the violet extremity of the spectrum than when they were exposed to the other rays; but that the combinations of chlorine and hydrogen in dry vessels took place more rapidly in the red rays. M. Berard has stated, however, an opposite result: but new experiments are wanting to decide this question. I made my experiment by concentrating the red rays and the rays beyond the red rays by a lens, and throwing them upon a small sphere

of copper within the vessel, so that the heat communicated to the metal may have interfered with the result: and in M. Berard's trials the vessels were moist, so that his result may have been connected with a decomposition of water.

I found that the black oxide of mercury exposed in the red rays concentrated by a lens gradually became red; in which case oxygen was probably absorbed; but the same oxide exposed in the violet rays concentrated in the same manner was not changed; and these rays produced no effect upon dry red oxide of mercury, but upon moistened red oxide occasioned the same change as a current of hydrogen gas.

Dr. Wollaston has shewn that a change of colour is produced upon gum guaiacum by the rays beyond the violet rays connected with an absorption of oxygen; and hence he has proposed to call the invisible rays in this part of the spectrum, chemical rays, and M. Berard has adopted this distinction; but supposing that hydrogen is disengaged in the experiment on guaiacum it may in its nascent state absorb oxygen, and the change of colour may depend not upon the absorption of oxygen, but the separation of hydrogen or its new combination. The facts that I have stated above, sufficiently demonstrate that the rays producing heat are capable of assisting certain species of chemical action, and there seems no more reason for asserting that the rays producing no heat are the rays most efficacious in occasioning chemical changes, than for asserting with MM. Gay Lussac and Thenard, that light produces all its chemical effects by producing heat.

5. In general, in nature, the effects of the solar rays are very compounded. Healthy vegetation depends upon the presence of the solar beams, or of light, and

whilst the heat gives fluidity and mobility to the vegetable juices, chemical effects likewise are occasioned, oxygen is separated from them, and inflammable compounds formed. Plants deprived of light become white, and contain an excess of saccharine and aqueous particles; and flowers owe the variety of their hues to the influence of the solar beams.

Even animals require the presence of the rays of the sun, and their colours seem materially to depend upon the chemical influence of these rays; a comparison between the polar and tropical animals, and between the parts of their bodies exposed, and those not exposed, to light, shews the correctness of this opinion.

IV. *Of the Nature of the Motions or Affections of Radiant Matter.*

1. Two hypotheses have been invented to account for the principal operations of radiant matter. In the first it is supposed that the universe contains a highly rare elastic substance, which when put into a state of *undulation* produces those effects on our organs of sight, which constitute the sensations of vision, and the other phenomena occasioned by solar and terrestrial rays. In the second it is conceived that *particles* are *emitted* or sent off from luminous or heat-making bodies with great velocity, and that they produce their effects by communicating their motions to substances, or by entering into them, and changing their composition.

2. The first of these suppositions was adopted by Hooke, Huygens, and Euler;—the second by Newton, and the philosophers of the Newtonian School. Many of the phenomena may be accounted for by either hypothesis, but the Newtonian doctrine applies much more happily to most of the facts discovered respecting the

modifications of light by double refraction and reflection. Indeed it does not seem possible, as Newton has shewn, to account for the circumstance,—that a ray which has suffered extraordinary refraction in passing through one crystal, should suffer ordinary refraction in passing in another direction, through another like crystal, on the idea of the effect being a mere undulation of an ethereal medium; but it may be explained by supposing the rays to consist of particles endowed with rectilinear motion, and possessed of a certain polarity, that is, parts attractive with respect to some surfaces of the crystal, and repulsive with respect to others.

3. M. Malus has supposed, in his ingenious speculations on these remarkable phenomena, that the molecules producing light, are possessed of three rectangular axes, of which one is always in the direction of the ray, and the other two are made by the influence of the repulsive forces exerted by the crystalline medium, perpendicular to the direction of these forces; and such a form, and such an effect would correspond with the idea of the luminous particles being octohedrous.

4. As the coloured rays separated by the prism bear the same relation to double refraction, that direct light bears, it follows that the polarity of the different particles must be of the same kind, and this is what might be expected. The same crystalline substance always affects the same primary forms. When a tourmaline is broken into pieces, the pieces are found to possess similar electrical powers to the original crystal, and a large rhomb of calcareous spar, easily breaks into a number of small rhombs.

5. Newton has attempted to explain the different refrangibility of the rays of light, by supposing them composed of particles differing in size, and this hypo-

thesis is not contradictory to the idea of their being regular solids endowed with similar polarities. The same great man has put the query whether light and common matter are not convertible into each other; and adopting the idea that the phenomena of sensible heat depend upon vibrations of the particles of bodies, supposes that a certain intensity of vibrations may send off particles into free space, and that particles in rapid motion in right lines, in losing their own motion, may communicate a vibratory motion to the particles of terrestrial bodies.*

* The views of Newton are so clearly developed in the following passages, and they are so much connected with the refined philosophy of Chemistry, that the reader probably will require no apology for the insertion of them in a note.

Quere 29. Are not the rays of light very small bodies emitted from shining substances? For such bodies will pass through uniform mediums in right lines, without bending into the shadow, which is the nature of the rays of light. They will also be capable of several properties, and be able to conserve their properties unchanged, in passing through several mediums which is another condition of the rays of light. Pellucid substances act upon the rays of light at a distance, in refracting, reflecting, and inflecting them; and the rays mutually agitate the parts of these substances at a distance for heating them; and this action and re-action at a distance, very much resembles an attractive force between bodies. If refraction be performed by attraction of the rays, the sines of incidence must be to the sines of refraction in a given proportion; as we shewed in our principles of philosophy, and this rule is true by experience. The rays of light in going out of a glass into a *vacuum*, are bent towards the glass; and if they fall too obliquely on the *vacuum*, they are bent backwards into the glass, and totally reflected; and this reflection cannot be ascribed to the resistance of an absolute *vacuum*, but must be caused by the power of the glass attracting the rays at their going out of it into the *vacuum*, and bringing them back. For if the farther surface of the glass be moistened with water, or clear oil, or liquid and clear honey, the rays which would otherwise be reflected, will go into the water, oil, or honey, and therefore are not reflected before they arrive at the farther surface of the glass, and begin to go out of it. But if they go out of it into a *vacuum*, which has no attrac-

6. As particles of any gaseous medium when put into a state of undulatory motion are capable of producing

tion to balance that of the glass, the attraction of the glass either bends and refracts them, or brings them and reflects them. And this is still more evident by laying together two prisms of glass, or two object-glasses of very long telescopes, the one plain, the other a little convex, and so compressing them that they do not fully touch, nor are too far asunder. For the light, which falls upon the surface of the first glass, where the interval between the glasses is not above the ten hundred thousandth part of an inch, will go through that surface, and through the air, or *vacuum* between the glasses, and enter into the second glass, as was explained in the first, fourth, and eighth observations of the first *part* of the Second Book. But if the second glass be taken away, the light, which goes out of the second surface of the first glass into the air, or *vacuum*, will not go on forwards, but turns back into the first glass, and is reflected ; and therefore it is drawn back by the power of the first glass, there being nothing else to turn it back. Nothing more is requisite for producing all the variety of colours, and degrees of refrangibility, than that the rays of light be bodies of different sizes ; the least of which may make a violet, the weakest and darkest, of the colours, and the more easily diverted by refracting surfaces from the right course ; and the rest as they are bigger and bigger, may make the stronger and more lucid colours, blue, green, yellow, and red, and be more and more difficultly diverted. Nothing more is requisite for putting the rays of light into fits of easy reflection and easy transmission, than that they be small bodies, which by their attractive powers, or some other forces, stir up vibrations in what they act upon ; which vibrations being swifter than the rays, overtake them successively, and agitate them, so as by turns to increase and decrease their velocities, and thereby put them into those fits. And lastly, the unusual refraction of island crystal, looks very much as if it were performed by some kind of attractive virtue, lodged in certain sides, both of the rays, and of the particles of the crystal, and not in their other sides : for were it not for some kind of disposition or virtue lodged in some sides of the particles of the crystal, and which inclines and bends the rays towards the coast of unusual refraction, the rays which fall perpendicularly on the crystal, would not be refracted towards that coast, rather than towards any other coast, both at their incidence, and at their emergence, so as to emerge perpendicularly, by a contrary situation of the coast of unusual refraction, at the second surface ; the crystal acting upon the rays, after they have passed through it, and are emerging into the air, or, if you please, into a *vacuum*. And since the crystal, by this disposition, or virtue, does not

the sensation of sound by acting upon the auditory organs, so it may be conceived, that certain particles or aggregates of particles from any matter moving with great and equal velocity may occasion the sensations of vision, and the other effects of the solar beams; and the difficulty of refracting terrestrial radiant heat, may be conceived to depend upon the greater size of the aggregated particles; and according to the Newtonian hypothesis, any matter moving with considerable quickness in right lines may be conceived capable of communicating an expansive motion to the particles of bodies.

7. If specific highly rare imponderable fluids be assumed to account for the phenomena, as many must be adopted as there are different series of effects produced by different rays. There must be a matter of

act upon the rays, unless when one of their sides of unusual refraction looks towards that coast; this argues a virtue or disposition in those sides of the rays, which answers to, and sympathises with that virtue or disposition of the crystal, as the poles of two magnets answer to one another. And as magnetism may be intended and remitted, and is found only in the magnet and in iron, so this virtue of refracting the perpendicular rays is greater in island crystal, less in crystal of the rock, and is not yet found in other bodies. I do not say that this virtue is magnetical; it seems to be of another kind; I only say, that whatever it be, it is difficult to conceive how the rays of light, unless they be bodies, can have a permanent virtue in two of their sides which is not in their other sides; and this without any regard to their position to the space, or medium, through which they pass."—Optics.

May not the experiments of Dr. Young, *Phil. Trans.* 1804, page 2, which he considers as proving that homogeneous light, at certain equal distances, in the direction of its motion, is possessed of opposite qualities capable of neutralizing each other, and of extinguishing the light when they happen to be united; be explained on the idea of attractive poles in opposite sides of the particles of light. That able philosopher considered them as favourable to the theory of undulation; but if the attractions of other matter can destroy the motions of light, as in the case of its action on black bodies, may not the same result be produced by the attractions of its particles for each other?

violet light, a matter of blue light, and so on : and likewise a deoxydating ethereal matter, a calorific solar matter, and a calorific terrestrial matter, which is very contradictory to the usual simplicity of causes observable in the economy of things ; and the idea likewise is rendered improbable, by experiments on solar phosphori. When a mixture of calcined oyster shells, and sulphur, that have been heated together, is exposed to the solar rays, it forms a good solar phosphorus, it becomes luminous, and continues so for some minutes in the dark : and to whichever of the prismatic rays it be exposed, its light is always the same, pale yellow. It is easy to explain the phenomena, on the idea that vibratory motion is communicated to particles of the substance by the rays, in consequence of which, some of its own particles are slowly sent off, or that the particles have been formed into new aggregates in consequence of the attraction of the substance ; but if light be supposed specific in its kind, and absorbed and emitted ; then when the phosphorus is exposed to blue rays, blue rays alone ought to be emitted, which is not the case.

8. Many authors have written of the *combinations* of light and heat ; but from the views that have been developed, even taking the Newtonian theory of emission for granted, it is evident that such combinations are merely hypothetical. When the solar rays are *absorbed*, as it is called, by a black body, it must be conceived, on this theory, that their motion is communicated to the particles of the body, but whether they adhere to it, or are thrown off in new aggregates, as radiant heat, cannot well be discussed, for we have no means accurate enough to determine whether in such cases there is an increase of weight ; and this is the only

test to be depended upon, of true chemical combination, or of mechanical mixture.

The fire produced in a number of chemical processes, particularly in combustion, on the Newtonian view, may be ascribed to particles sent into free space, in consequence of the repulsion exerted by other particles at the moment of their entering into chemical union. Any solid bodies may be made to emit light, when exposed to a blast of air very hot, though not luminous; the light is always of the same kind, and this circumstance is favourable to the idea of the possibility of the conversion of common matter into radiant matter.

Many phenomena which have been attributed to combined light, appear to be electrical, or to be merely the effect of the ignition of the substances, for whenever heat rises beyond a certain degree, bodies become luminous; pieces of quartz rubbed together are rendered electrical; and by percussion or friction any hard bodies may be intensely heated.

During the putrefaction of certain animal and vegetable substances, light is emitted; and this is no more difficult to account for, than the heat produced during similar operations.

The light emitted by certain living insects, appears to depend upon the secretion of a substance very easy of decomposition: and any chemical change may be supposed adequate to the production of light.

It has been sometimes supposed that a specific imponderable substance, capable of producing light, is contained in oxygen gas; and it has been also imagined that such a substance exists in inflammable bodies; but the facts are contradictory to the hypothesis. Iron, when heated to whiteness, burns with amazing brilliancy in oxygen gas, throwing off sparks intensely luminous;

but when heated to 600° Fahrenheit, it combines slowly with the oxygen, producing heat without light; the chemical change is of the same nature in both cases; the only difference is in its rapidity and energy.

9. The later investigations on light, teach us that there is still much to learn with respect to the affections and motions of radiant matter; and this subject, when fully understood, promises to connect together chemical and mechanical science, and to offer new and more comprehensive views of the corpuscular arrangements of matter.

In radiant matter, the particles act almost independently of the common laws of attraction; and by prismatic refraction, the difference of their actions is determined, and it seems probable that the relations of the different particles to the crystalline arrangements of matter, will be found connected with those powers which they possess analogous to electrical qualities.

If that sublime idea of the ancient Philosophers which has been sanctioned by the approbation of Newton, should be true, namely, that there is only one species of matter, the different chemical, as well as mechanical forms of which are owing to the different arrangement of its particles, then a method of analysing those forms may probably be found in their relations to radiant matter. Newton supposed that the luminous particles at the violet end of the spectrum were smallest in size, and those at the red end, largest in size, and those producing the intermediate colours, intermediate. On this idea, the calorific invisible particles would be the largest in the solar beam, and the calorific particles emitted by terrestrial bodies, may be imagined of still greater size, so as to be incapable of passing through the pores of dense transparent media. The rays at the

red end of the spectrum in their chemical powers heat bodies, and probably produce all those combinations which depend upon elevation of temperature ; those at the opposite end tend to produce combinations in which hydrogen is concerned ; and this is the lightest chemical element in nature, and may be conceived to be composed, on the corpuscular hypothesis, of the smallest particles, and therefore likely to be acted on in a higher ratio than other elements by the smallest particles of light.

10. The idea that light is not a specific fluid, is confirmed by some practical results relating to the economy of light. Count Rumford has lately shewn that the quantity of light emitted by a given portion of inflammable matter in combustion, is proportional in some high ratio to the elevation of temperature ; and that a lamp having many wicks very near each other, so to communicate heat, burns with much more brilliancy in proportion to the consumption of oil than the Argand's lamps in common use.

DIVISION III.

OF EMPYREAL UNDECOMPOUNDED SUBSTANCES, OR UNDECOMPOUNDED SUBSTANCES THAT SUPPORT COMBUSTION, AND THEIR COMBINATION WITH EACH OTHER.

I. *General Observations.*

1. It has been mentioned that almost all cases of vivid chemical action are connected with the increase of temperature of the acting bodies, and a greater radiation of heat from them ; and in a number of instances, light is also produced, see p. 64, and 119.

The strength of the attraction of the acting bodies determines the rapidity of combination, and in proportion as this is greater, so likewise is there more intensity of heat and light. In the phlogistic doctrine of chemistry, all changes in which heat and light are manifested, were explained by supposing that the acting bodies contained the principle of inflammability; in the anti-phlogistic doctrine, most of them have been accounted for by imagining the position or transfer of oxygen: but all the later researches seem to shew that no *peculiar* substance, or form of matter is necessary for the effect; that it is a *general* result of the actions of any substances possessed of strong chemical attractions, or different electrical relations, and that it takes place in all cases in which an intense and violent motion can be conceived to be communicated to the corpuscles of bodies.

2. Many bodies which have not yet been decomposed, and which cannot well be conceived to contain oxygen, produce heat and light by their mutual chemical action—such are some metallic substances, potassium, for instance, in combining with arsenic and tellurium; and sulphur and certain metals become ignited during their union. In naming a class of bodies by their relations to combustion, or to their efficacy in producing the phenomena of fire, it is only intended to signify that the production of heat and light is more characteristic of their actions, than of those of any other substances; and they are likewise opposed to all other undecomposed substances by their electrical relations, being always in voltaic combinations attracted to, or elicited from the *positive* surface; whereas all other known undecomposed substances are separated at the *negative* surface. Only two undecomposed empyreal substances have been as yet discovered. They

will be described, and their actions on each other discussed in the two following sections.

II. *Of Oxygen Gas.*

1. *Oxygen* gas was discovered by Dr. Priestley, in August, 1774. To procure it, a quantity of manganese (a mineral substance found in abundance near Exeter, and in many other places) is introduced into a glass retort furnished with a ground stopper, a quantity of oil of vitriol (sulphuric acid) sufficient to moisten the manganese, is added, and they are mixed together by means of a glass rod; the bottom of the retort is then gently heated by means of a lamp, and the extremity of its neck introduced under an inverted cylinder filled with water in the hydro-pneumatic apparatus.* Globules of gas will soon rise through the water; the first portions collected must be thrown away, being principally the common air contained in the retort; when a quantity more than equal to the capacity of the retort has been thus disposed of, the remainder may be preserved for use.

There are many other modes of obtaining oxygen gas. Manganese heated to redness in an iron tube, such as a gun-barrel, the touch-hole of which is closed, will afford a considerable quantity of the substance, which may be collected by means of a tube fastened into the neck of the barrel, and having its extremity in the hydro-pneumatic apparatus.†

Nitre heated strongly in a porcelain retort gives off oxygen gas: puce-coloured, or red oxide of lead offers a similar result; and from any of the salts called hy-

* See Plate VI. fig. 22.

† See Plate VI. fig. 23.

peroxymuriates (chlorates), oxygen is procured by a dull red heat ; a retort of glass may be employed in the process ; and a charcoal fire in a small chafing-dish.

100 grains of the hyperoxymuriate of potassa, afford 114 cubical inches of oxygen gas, under common circumstances.

The oxygen gas procured from nitre and the metallic substances above-mentioned, is mixed with larger or smaller quantities of other permanent gaseous matter ; the gas from hyperoxymuriate of potassa is free from such adulteration, and when collected over mercury, contains nothing but aqueous vapour, from which it may be purified by means of the salt called dry muriate of lime (chloride of calcium), or by sticks of common potash. The elastic fluid from nitre contains more foreign gaseous matter than that from the metallic oxides. The gas from manganese and sulphuric acid, when collected in the mercurial apparatus, seldom affords more than $\frac{1}{50}$ of adulteration ; when collected over water, it is mixed with from $\frac{1}{20}$ to $\frac{1}{30}$, in consequence of mixture with the common air expelled from the water.

The degree of purity of oxygen gas procured in these modes, is easily ascertained by filling a small curved tube closed at one end, with mercury, and passing into it some of the dry gas, so as to occupy about $\frac{1}{4}$ of its capacity, which is measured ; a bit of phosphorus in the proportion of half a grain to a cubical inch of gas, is introduced and made to pass into the curved part of the tube ; the phosphorus is inflamed by the application of the heat of a spirit lamp ; at the moment of the expansion of the gas, the open extremity of the tube is closed under the mercury by the finger, and heat is applied till no light can be perceived in the tube ; when the tube is cool, the finger is taken off. The mercury will instantly

rise into the tube ; all the oxygen will have been absorbed, and the gas remaining, when measured and compared with the original quantity, will indicate the impurity.

2. Oxygen gas is distinguished from all other gaseous matter by several important properties.

Inflammable substances burn in it under the same circumstances as in common air, but with much greater vividness.

If a taper, the flame of which has been extinguished, the wick only remaining ignited, be plunged into a bottle filled with it, the flame will be instantly re-kindled, and will be very brilliant, and accompanied by a crackling noise.

If a steel wire, or thin file, having a sharp point, armed with a bit of wood in inflammation, be introduced into a jar filled with the gas, the steel will take fire, and its combustion will continue producing a most brilliant phenomenon.

The specific gravity of oxygen gas has been already referred to, page 81 ; it is to that of hydrogen gas, as 15 to 1 ; 100 cubical inches under an atmospherical pressure equal to that of 30 inches of mercury, and at the temperature of 60° Fahrenheit, weigh about 34 grains. Its power of refracting light is stated by Biot and Arago to be to that of hydrogen, nearly as 1958 to 1000. Its capacity for heat, according to Dr. Crawford, is nearly as 4·7 to 21·4.

Oxygen gas is slightly absorbable by water. From Dr. Henry's experiments, it appears that this fluid takes up $\frac{1}{27}$ of its bulk at 60° Fahrenheit, whatever be the density of the gas.

Oxygen gas is respirable ; a small animal confined in a jar filled with this gas, lives four or five times as long

as in an equal quantity of common air;—hence it has been called vital air.

3. The proportion in which oxygen gas unites to bodies, has been referred to, page 82, and the number representing it may be considered as 15; various elucidations of the correctness of this conclusion, will be found in the following pages.

4. Oxygen gas forms the most important part of our atmosphere. It is easy to prove this by many very simple experiments.

If phosphorus be inflamed in a tube half filled with atmospherical air, in the same manner as in the experiment for ascertaining the purity of oxygen, a quantity of elastic matter will be absorbed equal to about one-fifth of the volume of the confined air, and the same substance will be produced as that formed by burning phosphorus in oxygen: the remaining elastic fluid will not support flame, and animals will not live in it; it is called azote or nitrogen gas; and if four parts of it be mixed with about one part of pure oxygen gas, they constitute a mixture resembling exactly atmospheric air. That the oxygen obtained artificially is the same chemical substance as that found in the air, is proved by the phenomena of the calcination of mercury. If running mercury be preserved in a heat at which it boils slowly, in a retort, the beak of which is plunged in mercury, and the process be continued for some days, there will be a gradual diminution of the air, and after a certain time, the remainder will not support flame, and a part of the mercury will be found converted into a red powder. It will have gained in weight, as much as the air has lost; and the red powder, if heated to ignition, will give off a quantity of oxygen, that, added to the residual elastic substance, will reconstitute common air, and it will be

restored to the state of mercury. And this oxygen, if a part of it be compared with the oxygen procured in other modes from mineral substances, or artificial compounds, is found in no respect different; its specific gravity, refractive power, and chemical properties, are precisely identical.

There have been several substances proposed for ascertaining with facility the quantity of oxygen in air; they have been called eudiometrical substances; and the instruments in which they have been employed, are named eudiometers. The solution obtained by water from sulphur and pearl ashes, or sulphur and lime that have been fused together, slowly absorbs oxygen; a solution of tin in muriatic acid has a similar property; and likewise solutions of iron, into which nitrous gas has been passed till they become coloured. A tube of glass graduated to 100 parts, forms a good eudiometer; and when filled with air, it is plunged into any solution that will absorb oxygen, and suffered to remain there, till the process is complete.

It was formerly supposed that there are great differences in the quantity of oxygen in air, in different places, and at different times; but all late researches shew that this opinion is erroneous.

Air, analyzed in different quarters of the globe, in cities, and in the country, on sea and land, has been found not perceptibly different in composition; the accurate proportions of oxygen and azote are 21 and 79.

It has been shewn by the experiments of Dr. Priestley, Mr. Dalton, and M. Berthollet, that different elastic fluids have a tendency to rapid equable mixture, even when at rest, and exposed to each other on small surfaces only; and the mixture of the parts of the atmosphere is constantly assisted by winds, by currents of

air, and by all the motions taking place on the surface of the earth.

5. In all processes of combustion in the atmosphere, oxygen is either fixed in the combustible body, or it dissolves it, or forms a new compound with it. In respiration, as will be more fully explained in the last part of this work, the volume of air is not changed; but a part of its oxygen disappears, and an equal bulk of carbonic acid gas is found in its place.

As the constitution of the atmosphere constantly remains the same, it is evident that there must be some processes in nature, by which a quantity of oxygen is produced equal to that consumed. One principal cause of the renovation of oxygen appears to be in the process of vegetation; healthy plants exposed in the sunshine to air containing small quantities of carbonic acid gas, destroy that elastic fluid, and evolve oxygen gas; so that the two great classes of organized beings are dependent upon each other. Carbonic acid gas, which is formed in many processes of combustion, as well as in respiration, if not removed from air, by its excess would be deleterious to animals, but it is a healthy food of vegetables; and vegetables produce oxygen, which is necessary to the existence of animals, and thus this part of the economy of nature is preserved, by the very functions to which it is subservient; and the order displayed in the arrangement, demonstrates the intelligence by which it was designed.

6. No other forms of matter have been produced from oxygen by any processes to which it has been submitted; but it readily enters into combination, and no substance is more active as a chemical agent. It is known to be a constituent part of most of the acids and earths, and of all the alkalies except one, and the his-

tory of its compounds, forms the most extensive and important part of modern chemistry.

Its operations, as will be seen hereafter, are connected with many of the arts; with the processes of bleaching, dyeing, colour-making, and metallurgy; and in its various applications to the production of fire, it is absolutely essential to cultivation, and to the comforts and enjoyments of social life.

In the phenomena of nature, it occasions a wonderful diversity of effects. It is active in most of the changes taking place on the surface of the globe, and its constant tendency is to unite different substances in forms adapted for the purposes of organized life.

II. *Chlorine, or Oxymuriatic Gas.*

1. This elastic substance was discovered by Scheele in 1774. It may be procured in the hydro-pneumatic apparatus, by a process very similar to that first described in the last section for procuring oxygen gas, but the manganese is to be mixed with common salt, and the oil of vitriol diluted with an equal quantity of water. The best proportions are three parts of common salt in weight, one part of manganese finely powdered, and two parts of oil of vitriol. Instead of manganese, red, or puce-coloured oxide of lead may be used; and instead of the common salt, and oil of vitriol, a solution of muriatic acid in water (spirits of salt. C.)

2. *Chlorine* is of a yellowish green colour, and it is this property which suggested its name.* Its odour is extremely disagreeable. It is not capable of being respired, and even when mixed in very small quantities,

* $\chi\lambda\omega\rho\omicron\varsigma$, green.

with common air, renders the air extremely pernicious to the lungs.

Its specific gravity is to that of hydrogen, nearly as 33·5 to 1, and 100 cubical inches of it weigh at mean temperature and pressure between 76 and 77 grains.

It is absorbable to a certain extent by water : at the temperature of 60° Fahrenheit, water dissolves about double its volume, and acquires a strong acrid taste, and a disagreeable smell.

When an inflamed taper is introduced into a phial filled with it, the light continues, but of a dull red colour, and a dark carbonaceous smoke rises from the flame.

Many of the metals introduced into it in thin filaments, or leaves, or powder, take fire, and burn spontaneously at common temperatures ; such as copper, tin, arsenic, zinc, antimony, and the alkaline metals.

Phosphorus burns in it spontaneously, with a pale white light, producing a white volatile powder.

Sulphur melted or sublimed in it, does not burn, but forms with it a volatile red liquor.

The gas does not change by any action of heat or cold;* but its aqueous solution freezes more readily than water, namely, at about 40° Fahrenheit.

When freed from vapour by muriate of lime, the gas does not act upon perfectly dry substances tinged with vegetable colours ; but when moisture is present in the gas or the coloured bodies, their colours are speedily destroyed, they are rendered white, or brought to a dull yellow ; and this last tint is almost the only one not changed by the combined action of water and chlorine.

3. Chlorine and oxygen are capable of existing in combination, and they form a peculiar gaseous matter.

* [By condensation it is rendered liquid. Vide Vol. I. p. 281.]

They do not unite, when mixed together, but when existing in certain solids, they may be detached in union.

To make the compound of chlorine and oxygen, hyperoxymuriate of potassa is introduced into a small retort of glass; and twice as much muriatic acid as will cover it, diluted with an equal volume of water. By the application of a gentle heat, the gas is evolved, and it must be collected over mercury.

I discovered this elastic substance in its pure form in January 1811, and gave to it the name of *Euchlorine*,* from its bright yellow-green colour.

Its tint is much more lively than that of chlorine, and more inclined to yellow.

Its smell is very different, being not unlike that of burnt sugar.

It is not respirable.

It is soluble in water, to which it gives a lemon colour, water takes up 8 or 10 times its volume.

Its specific gravity is to that of hydrogen, nearly as 33 to 1. 100 cubical inches weigh at mean temperature and pressure between 74 and 75 grains.

It must be collected and examined with great care, and only in small quantities at a time; a very gentle heat causes it to explode, sometimes even the heat of the hand; and its elements separate from each other with great violence, producing light.

From the facility with which *euchlorine* decomposes, it is not easy to ascertain the action of combustible bodies upon it. None of the metals that burn in chlorine, act upon this gas at common temperatures; but when the oxygen is separated, they then inflame in the chlorine. It is easy to witness this. Let a little Dutch foil be introduced into a bottle filled with *euchlorine*, it will

* *Ev* and *χλωρος*.

undergo no change, and will not even tarnish. Let a heated glass tube be applied to the gas in the neck of the bottle a decomposition will take place, and the oxygen and chlorine will be detached from each other, and at the same moment the foil will inflame, and burn with great brilliancy.

Chlorine is rapidly absorbed by mercury; euchlorine has no action upon it, and chlorine may be separated from euchlorine, by agitation over mercury, and the last obtained pure.

When phosphorus is introduced into euchlorine, it is instantly decomposed, and the phosphorus burns as it would do in a mixture of 2 parts in volume of chlorine, and 1 part of oxygen.

The inflamed taper, and inflamed sulphur, instantly decompose it, and exhibit the same phenomena as in a mixture of 2 parts of chlorine, and 1 part of oxygen.

That the gas is actually composed of these elements, is shewn by causing it to detonate in a glass tube over pure mercury. It loses its brilliant colour, and becomes chlorine and oxygen. 50 parts treated in this way, expand so as to become about 60 parts, which consist of 40 parts of chlorine, and 20 parts of oxygen.

When euchlorine freed from water, is made to act upon dry vegetable colours, it gradually destroys them, but first gives to the blues, a tint of red; from which, and its absorbability by water, and the taste of its solution, which is strongly acrid approaching to sour, it may be considered as approximating to an acid in its nature.

4. The proportion in which chlorine combines with bodies, may be learnt from the decomposition of euchlorine; the oxygen in which is to the chlorine, as 15 to 67 in weight. If euchlorine be considered as consisting

of one proportion of oxygen to one of chlorine, then 67 will be the number representing chlorine, which is most convenient, as being a whole number. If euchlorine be supposed to contain two proportions of chlorine and one of oxygen, then the number representing chlorine, will be 33·5. It will hereafter be shewn that whichever of these data be assumed, the relations of the number will harmonize with those gained from various other combinations.

5. Scheele considered chlorine as an element of the muriatic acid, and hence called it dephlogisticated marine acid. By that chemist, it was regarded as an undecomposed body.

Lavoisier and Berthollet asserted that it was a compound of muriatic acid gas, and oxygen. This idea is now universally given up; but some chemists in France and Scotland, conceive that it is a compound of oxygen, and an unknown body, which they call dry muriatic acid. The weight of chlorine, its absorbability by water, its colour, and the analogy of some of its combinations to bodies, known to contain oxygen, are arguments in favour of its being a compound; and it is possible that oxygen may be one of its elements, or that oxygen and chlorine are similarly constituted. I have made a number of experiments with the hope of detecting oxygen in it, but without success; none of its compounds with inflammable bodies or metals will afford this principle; charcoal intensely ignited in it, undergoes no change, nor is it altered by the strongest powers of electricity. Should oxygen ever be procured from it, some other form of matter, possibly a new one, will at the same time be discovered, as entering into its constitution, and till it is decomposed, it must be re-

garded, according to the just logic of chemistry, as an elementary substance.*

6. Chlorine has never been found pure in nature; but exists in many compounds, particularly in common salt, as is evident from the mode of its production from that substance. It is a substance of considerable importance in its relation to the art of bleaching, an application first made by the sagacity of M. Berthollet.

In the ancient process of bleaching, the cloths of linen and cotton, after being treated with alkaline lixivium, to free them from resinous and oily matters, and in some cases with very diluted oil of vitriol, to cleanse them from stains produced by iron, were exposed upon grass to dew and air, and some weeks, or even months, were required to give them their perfect whiteness.

By a warm solution of chlorine in water, they are bleached in a very short period; but their texture is injured; for at the same time that oxygen is added to the colouring matters, muriatic acid gas is formed and dissolved in the water, which corrodes the vegetable fibres.

The gas has been condensed in alkaline lixivium, and in lime water. The substance called oxymuriate of lime is commonly used for bleaching; but though the solution of this substance does not injure so much as that of the gas, yet it tends to weaken the texture of linen.

* MM. Gay Lussac, Thenard, and Curaudau, since 1808, have laid claim to the ideas of oxymuriatic gas being a simple body, and of muriatic acid gas being composed of this substance and hydrogen. But these opinions were started by the illustrious discoverer of the gas, in 1774. In the papers in the Philosophical Transactions in which I have endeavoured to shew that it is a peculiar acidifying and solvent principle, I have merely followed and extended his views, and I referred to them in the first paper I published on the subject.

I have found that the fluid produced by the condensation of the gas in water containing magnesia diffused through it, bleaches without injuring the vegetable fibre. It acts much more slowly and gradually than any of the other compounds employed for the same purpose, and has been applied at my suggestion in Ireland, within the last few months,* with success, in whitening printed calicos, and when properly used, it does not destroy even reds or yellows fixed by mordants. Magnesia may be easily procured from sea-water, or from the residual liquor of salt works; and there is a probability that this new bleaching fluid may, at no very distant period, come into common use.

M. Berthollet supposed that chlorine destroyed colours by parting with its oxygen; the new experiments shew that the oxygen is derived from the water, which is decomposed by double affinity; that of hydrogen for chlorine, and of the colouring matters for oxygen.

The salts which are called hyperoxymuriates, and oxymuriates, are compounds of metallic bodies with chlorine and oxygen; and the oxygen is held in them by a very weak attraction, and therefore is easily given off to colouring or inflammable matters.

The great circumstance, in bleaching with these compounds, is that the salt remaining after the abstraction of oxygen, should not act upon the linen; linen boiled in a strong solution of the salt called muriate of lime, the substance remaining in the solution, when oxymuriate of lime is used, I have found is considerably weakened. Solution of muriate of magnesia has no action of this kind, and therefore the new bleaching liquor can hardly be injurious to the manufacture.

* By Mr. Duffy, of Dublin, a very enlightened calico-printer, [in 1811.]

These general views respecting chlorine, and the uses and mode of agency of the combinations of chlorine and oxygen, will be found to be confirmed by a number of statements, to be given in the progress of this work. Certain conclusions have been anticipated,—but the most important application of chlorine could not with propriety be separated from its history as an undecomposed body, though depending upon its power of detaching oxygen, which appears to be the true bleaching principle, from compounds.

DIVISION IV.

OF UNDECOMPOUNDED INFLAMMABLE OR ACIDIFEROUS SUBSTANCES NOT METALLIC, AND THEIR BINARY COMBINATIONS WITH OXYGEN AND CHLORINE, OR WITH EACH OTHER.

I. *Preliminary Observations.*

The bodies to be considered in this division, are six, hydrogen, azote, sulphur, phosphorus, carbon, and boracium, or boron. Amongst these, hydrogen is distinguished from all the rest by very singular properties. Sulphur and phosphorus are the most analogous to each other. All these substances are capable of combining with oxygen, and all except azote and charcoal, with chlorine.* They are separated in Voltaic combinations, at the negative surface, and in their electrical relations, as well as chemical powers, are opposed to oxygen and chlorine.

[* Azote and charcoal have since been combined with chlorine.]

II. *Hydrogen Gas, or Inflammable Air.*

1. This elastic substance was first examined in its pure form, by Mr. Cavendish, in 1766.

It may be procured in the hydro-pneumatic apparatus from zinc or iron filings; by means of oil of vitriol diluted with eight times its weight of water; a retort, or a bottle furnished with a tube may be used; no artificial heat is required in the process. It may likewise be produced by passing steam over turnings of iron heated to redness in a gun-barrel.

2. *Hydrogen* is distinguished from all other gaseous bodies, by its extreme lightness. The relation of its weight to that of oxygen and air, has been already stated. 100 cubical inches of it at mean temperature, and pressure, weigh about two grains and a quarter.

It is very slightly absorbable by water; that fluid takes up $\frac{1}{50}$ only of its volume.

Hydrogen gas has no taste, a slight but disagreeable smell. It is capable of being taken into the lungs, but cannot be breathed by man for more than a minute. Small animals die in it in a much shorter time.

When an inflamed taper is plunged into a long narrow jar filled with hydrogen, and opened in the atmosphere, it is extinguished; but the gas takes fire, and burns in contact with the atmosphere.

One part mixed with two or three parts of air explodes violently by the action of an inflamed body, or an electrical spark.

3. Hydrogen gas, as has been stated, combines with oxygen gas, and to this circumstance its inflammation in the air is owing. If the two gases be pure, *water* is the only result, and the proportions are 2 of hydrogen

to 15 of oxygen in weight, or 2 to 1 in volume. The union may be effected by the electric spark as described in page 73, over mercury, or the hydrogen may be introduced into a vessel full of oxygen through a narrow tube, by means of pressure, and inflamed by electricity, or the oxygen may be made to burn in the hydrogen in a similar manner.* When a stream of oxygen is thrown into a stream of inflamed hydrogen, the heat produced is very intense, and far exceeds the highest heat of our furnaces, and may be used to fuse bodies, intractable by any other fire raised by combustion.

The nature of water may be shewn synthetically as well as analytically.

It is separated into 2 of hydrogen in volume and 1 of oxygen in the Voltaic circuit; the oxygen appears at the positive, the hydrogen at the negative metallic surfaces; and by means of platina wires, hermetically sealed into glass tubes, the products are collected.

When 10 grains of the metal called potassium are added to about 2 grains of water in a glass tube, there is a violent action, much hydrogen is disengaged, and by heating the results, the operation is completed. The same effect is produced upon the potassium, as would be produced by heating it strongly in contact with a small quantity of oxygen; it becomes united to oxygen, and its increase of weight is in proportion to the weight of the hydrogen, as 15 to 2.

It will be needless to dwell upon the properties of water; it is scarcely ever found in nature pure, usually holding saline or gaseous matters in solution. It becomes solid at 32° Fahrenheit and elastic at 212°, and in the state of steam has been applied for the production

* See Plate VII. fig. 24.

of the most important mechanical effects in the steam engine.

To describe the uses of water in the operations of nature, or to point out its applications to the purposes of the arts, and common life, would demand a volume. Animals and vegetables depend upon it for their existence. Water occupies nearly two-thirds of the surface of the globe; and whether existing in the ocean united to salts, or in the atmosphere as vapour, or poured down upon the surface as rain, dew, hail, or snow, or collected in lakes, rivers, and springs, its effects are constantly connected with the order of the economy of our system.

4. Hydrogen and chlorine unite with still more readiness than hydrogen and oxygen. To make the combination, it is only necessary to expose a mixture of equal volumes of the two gases to common day light, over dry mercury, or in a vessel furnished with a stop-cock previously exhausted. In a certain time, the chlorine will have lost its colour, and have combined with the hydrogen. If the gases have been freed from aqueous vapour, there will be no notable condensation, and the result is a peculiar elastic fluid, *muriatic acid gas*. By exposure to direct solar light as has been stated before, they explode; they likewise explode by the electrical spark; the results in this case, as I have found, are the same, 1 in volume of hydrogen unites to 1 in volume of chlorine, or 1 in weight to 17.

The nature of muriatic acid gas may be proved by analysis, as well as synthesis. If some pure grain tin be kept melted for some time, in a little curved tube containing muriatic acid gas over dry mercury, the tin will be converted into the same substance as that pro-

duced by its direct action upon chlorine, Libavius's liquor; and the hydrogen gas, when accurately measured, will be found to be equal to one-half the volume of the muriatic acid gas.

Those persons who suppose chlorine to be a compound of an unknown body, and oxygen, conceive muriatic acid gas to be a compound of $\frac{1}{4}$ of its weight of water, and the same hypothetical substance; but as no oxygen has yet been shewn to exist in chlorine, so no such combined moisture has been proved to exist in muriatic acid gas. It contains minute quantities in the vapour of hydrated muriatic acid; but no water except this can be procured from it, unless by substances that contain oxygen; and the quantity produced, is exactly proportional to the oxygen contained in the substance, and the hydrogen in the muriatic acid gas, and the other result is the same as the substance combined with the oxygen would produce *directly* by its action upon chlorine.

Five grains of red oxide of mercury, heated to redness, gave off a cubical inch, and $\frac{1}{8}$ of oxygen gas. Five grains of the same substance were made to act on muriatic acid gas by a spirit lamp in a curved tube over mercury; corrosive sublimate was formed, and water which absorbed muriatic acid gas, and 5 cubical inches of muriatic acid gas disappeared; and of these 4 cubical inches and $\frac{1}{2}$ at least, must have been decomposed by the oxide of mercury, their chlorine united to the metal, and their hydrogen to the oxygen; and the additional half a cubic inch, as will appear from the facts about to be stated, is nearly the quantity that ought to be absorbed by the water; the barometer in this experiment stood at 30.3; the thermometer at 54° Fahrenheit. Corrosive sublimate is produced by the direct combina-

tion of mercury and chlorine; and the results of this experiment can only be logically explained, on the idea of muriatic acid gas being composed of hydrogen and chlorine.

For the purposes of experiments, muriatic acid gas is procured by the action of oil of vitriol on certain salts, such as common salt, or sal ammoniac. It rises without the application of heat, when the substances are mixed together; a glass retort should be used with a ground stopper, the salt should be in large pieces, not in powder, and some bibulous paper should be introduced into the neck of the retort, to prevent any fluid acid from soiling the mercury, over which it must be collected.

Muriatic acid gas instantly extinguishes flame. It reddens dry litmus paper. When suffered to pass into the atmosphere, it produces a white smoke by uniting to the aqueous vapour in the air. Its taste is intensely acid. Its smell, pungent and disagreeable.

The specific gravity of muriatic acid gas is to that of hydrogen, nearly as 17 to 1; 100 cubical inches of it weigh at mean temperature and pressure, between 39 and 40 grains. Muriatic acid gas is rapidly absorbed by water; at the temperature of 40° Fahrenheit, water absorbs about 480 times its bulk of gas, and forms solution of muriatic acid gas in water, the specific gravity of which is 1.2109.

The table which follows, exhibits the quantity of muriatic acid gas in solutions of different specific gravities constructed after experiments made in the Laboratory of the Royal Institution.*

* 47.25 grains of water at 43° Fahrenheit, barometer being at 30.2, absorbed 34.8 grains of gas, and formed a solution of specific gravity,

At temperature 45° Fahrenheit.		Barometer 30.
100 parts of solution of muriatic acid gas in water of specific gravity.		Of Muriatic Acid Gas, parts.
1·21	Contain	42·43
1·20*		40·80
1·19		38·38
1·18		36·36
1·17		34·34
1·16		32·32
1·15		30·30
1·14		28·28
1·13		26·26
1·12		24·44
1·11*		22·3
1·10		20·20
1·09		18·18
1·08		16·16
1·07		14·14
1·06		12·12
1·05		10·10
1·04		8·08
1·03		6·06
1·02		4·04
1·01		2·02

1·21, and the whole precipitated by nitrate of silver, afforded about 132 grains of dry horn silver.

Again, 57·5 grains of water at 44°, barometer being 30·1, gained nearly 38 grains by absorbing acid gas, and formed a solution of specific gravity, 1·2.

Thermometer, 49° Fahrenheit; barometer 29; 45·5 grains of water by absorbing 13·4 grains of gas gained a specific gravity of 1·114. The two last results, which are marked in the table, agree with those gained by calculation from the first experiment. When about 150 grains of the strongest solution of muriatic acid in water, were mixed with distilled water, both being at 63°, the temperature rose to 75°; so that the real specific gravity of solutions mixed with water, is probably a little greater than the mean, though to no amount that can interfere with the use of the table. To find the composition of an acid of specific gravity not marked in the table, find the difference between the two specific gravities nearest to it in the table d , and the difference between their quantities of gas b , likewise the difference between the given specific gravity, and that nearest to it, c , then d is to $b :: c : x$, which added to the quantity of the lower specific gravity, is the quantity of acid gas sought.

The compound of water, and muriatic acid gas existing in vapour in muriatic acid gas, alluded to page 137, is probably of the same constitution as the most saturated solution at the same temperature, and at 45° must contain 57.57 per cent. of water; but in common cases the quantity of this vapour is too small to influence to any extent the results of experiments on muriatic acid gas; for I found that 200 cubical inches of gas at 75° passed slowly through a thin tube of glass cooled to 10° below 0 of Fahrenheit, did not increase its weight $\frac{1}{18}$ of a grain, but the deposition of fluid was very distinct.

5. The number representing hydrogen, as is evident from the details given page 81, and those just stated, and as will appear from a number of other evidences, may be considered as unity.

6. Of all gaseous substances, hydrogen is most distinctly characterized as an element; and in its relations it is opposed to oxygen.

Its extreme lightness, and the small quantities in which it enters into combination, render it unlikely that it should be resolved into other forms of ponderable matter, by any instruments or processes at present within our power. Some extraordinary phenomena which have been explained in the idea of its being a compound, and which will be referred to towards the end of this volume, are more satisfactorily accounted for on the idea of its being simple, or at least a form of elementary matter.

Hydrogen gas is employed for filling balloons, and its low specific gravity renders it well fitted for aërostatic purposes. It is an important principle in animal and vegetable bodies; and exists in larger or smaller quantities in all organized compounds. It is the body which

gives the power of burning with flame to all the substances used for the economical production of heat and light.

III. *Of Azote, or Nitrogen Gas.*

1. *Azote* was discovered by Dr. Rutherford in 1772. It may be procured by extracting oxygen from common air, in the manner described page 170. It is formed directly by dissolving animal matters, such as glue or muscular fibre, in diluted aquafortis, or fuming nitrous acid mixed with ten or twelve times its weight of water. It may be collected over water.

Azote extinguishes flame. It is very slightly absorbable by water; that fluid, according to Dr. Henry's experiments, takes up only $\frac{1}{75}$ part of its volume. Its specific gravity, as was mentioned in page 81, is to that of hydrogen as 13 to 1. 100 cubical inches of it, at mean temperature and pressure, weigh between 29 and 30 grains. According to Biot and Arago, its refractive power is 58976.

Its capacity for heat, according to Dr. Crawford, is .7936.

2. There are several compounds containing azote and oxygen in different proportions; three of which have been already referred to, page 81. Their nature is more easily demonstrated by analysis than synthesis; though the most important of them, nitrous acid in its union with water, may be made by the direct combination of azote and oxygen with that fluid.

Dr. Priestley ascertained that acid matter was formed by passing electrical sparks through a mixture of azote and oxygen over water, and Mr. Cavendish, by a series of beautiful experiments, proved that the two gases

combined with the water, and formed the same acid as that procured from nitre by oil of vitriol. The other compounds of azote and oxygen are always formed from the decomposition of this acid, or some of its compounds; but as *nitrous acid* exists in different states, its properties will be best understood after the more simple combinations of azote and oxygen have been described.

3. *Nitrous oxide*, the compound containing the smallest quantity of oxygen, was discovered by Dr. Priestley in 1772, and named by him dephlogisticated nitrous air.

It is a gaseous body, which may be produced by heating nitrate of ammonia; a glass retort is employed to contain the salt; the flame of an Argand lamp is sufficient to produce the gas. It may likewise be obtained during the solution of zinc in very weak nitric acid, but in this case it is not pure.

Nitrous oxide may be preserved over water; but it is absorbed by this fluid, which takes up $\frac{9}{10}$ of its volume nearly, and for accurate experiments, it should be collected in the mercurial apparatus.

Its degree of purity may be learnt from the quantity absorbed by water.

Nitrous oxide exhibits the following properties. A taper, plunged into it, burns with great brilliancy, and the flame gradually becomes surrounded with a blueish halo. Phosphorus may be melted and sublimed in it without inflaming; but when introduced into it in a state of vivid combustion, the brilliancy of the flame is greatly increased. Sulphur, and most other combustible bodies, require a higher degree of heat for their combustion in it than they require in oxygen, or in the atmosphere.

Its specific gravity, according to my experiments, is to that of hydrogen, nearly as 21 to 1. 100 cubical inches of it at mean temperature and pressure, weigh between 48 and 49 grains.

Its taste is sweetish, its odour slight but agreeable.

It is respirable, but not fitted to support life. I ascertained in 1799, that when it was respired, it produced effects analogous to those produced by drinking fermented liquors,—usually a transient intoxication, or violent exhilaration. Individuals that differ in temperament are, however, as might be expected, differently affected.

The composition of nitrous oxide has been stated, page 81. One in volume of this gas is decomposed by one volume of hydrogen, water is formed, and one in volume of azote remains.

Or if well-burnt charcoal be inflamed in a volume of it by a burning glass, 1 in volume of it affords as much carbonic acid as half a volume of oxygen, and when this carbonic acid is absorbed, a volume of azote remains; so that it consists of 26 in weight of azote, and 15 of oxygen.

4. *Nitrous gas* was noticed by Dr. Hales, but its properties as a specific elastic fluid were first described by Dr. Priestley in 1772; it is procured during the solution of various bodies in nitric acid; sugar, silver, mercury, copper, bismuth, afford it very readily. Filings of copper are usually employed; and a retort, or a bottle having a tube, inserted into it is used; the acid (if the common acid of commerce) should be diluted with 6 or 8 times its weight of water; the production of the gas may be assisted by a gentle heat.

It may be collected over water, which absorbs only

about $\frac{1}{20}$ of its volume ; but for accurate purposes mercury should be employed.

The degree of purity of nitrous gas may be known by agitating it in contact with an aqueous solution of green sulphate of iron. Nitrous gas is quickly absorbed by this substance.

When a jar of nitrous gas is opened in the atmosphere, red fumes appear. When an inflamed taper is plunged into it, the light is instantly extinguished.

Inflamed sulphur is extinguished by it; but inflamed phosphorus burns in it with great brilliancy. It cannot be made to detonate when mixed with hydrogen, by the electric spark.

Its specific gravity is to that of hydrogen, as fourteen to one. 100 cubical inches of it weigh about 32 grains.

Whether it is respirable, or has taste or smell, cannot be ascertained, as it instantly unites with the oxygen in air, producing red fumes, which are nitrous acid gas.

The composition of nitrous gas has been already referred to, page 81.

It is decomposable by several of the metals when they are heated in it, such as arsenic, zinc, potassium in excess; it oxidates them, and affords half its volume of azote. In an experiment in which I decomposed a small quantity by igniting charcoal in it by a burning glass, I found that it afforded about half a volume of carbonic acid, and half a volume of azote; so that it consists of 26 of azote to 30 of oxygen.

When it is exposed to certain bodies, such as the salts called sulphites, solution of tin in muriatic acid, or solutions of alkaline sulphurets, it is converted into nitrous oxide; I have found that in accurate experi-

ments of this kind, two in volume of nitrous gas become one in volume of nitrous oxide ; a circumstance harmonizing precisely with their relative proportions of oxygen and nitrogen.

5. It has been mentioned that the red fumes produced by the action of oxygen and nitrous gas, are owing to the production of *nitrous acid gas*.

It is not easy to ascertain the exact nature of this change, as the substance formed acts both upon mercury and upon water ; and over water very different proportions of the gases may be made to condense each other. When large quantities of nitrous gas are added to small quantities of oxygen, in vessels of large diameter, from two to three in volume of nitrous gas disappear for one of oxygen. When large quantities of oxygen are added to small quantities of nitrous gas in narrow tubes, the absorption is from 1 to 1.5 of oxygen in volume, and 2 of nitrous gas. From a series of experiments on the decomposition of nitre, and others on the mixture of nitrous gas and oxygen, executed with great care in exhausted vessels furnished with glass stop-cocks, I am inclined to believe that the acid obtained over water by the condensation of mixtures of nitrous gas and oxygen is never fully saturated with oxygen, and that the pale fluid called nitric acid consists of water united to two in volume of nitrous gas, and one and a half of oxygen ; and this acid, according to its different degrees of dilution, may be made to absorb different quantities of nitrous gas, when it becomes yellow, orange, blue, or blueish green ; and in this last state it is saturated with nitrous gas.

When two of nitrous gas, and one of oxygen freed from moisture, are mixed together in a vessel previously exhausted of air, they become condensed to about one-

half of their volume, and form a deep orange-coloured elastic fluid, which may be called *nitrous acid gas*.

This substance has the following properties: a taper burns in it with considerable brilliancy. Sulphur inflamed, does not burn in it; but the combustion of phosphorus continues with great vividness.

Tin, copper, and mercury, act upon it slowly; iron ignited to whiteness is rapidly cooled in it.

Charcoal inflamed continues to burn in it with a dull red light.

When a portion of water is exposed to it, there is a rapid absorption, and the water gains a tint of green.

Its smell is very disagreeable, its taste sour; when applied to animal substances, it renders them yellow; it reddens litmus paper.

Calculating from the condensation, the specific gravity of nitrous acid gas is to that of hydrogen as about 29 to 1; and 100 cubical inches of it weigh 65·3 grains, at mean temperature and pressure.

6. I have attempted to procure a permanent elastic fluid, consisting of 2 parts in volume of nitrous gas, and 1·5 oxygen, by mixing oxygen in excess with nitrous gas; but the condensation was always such as to indicate the formation of nitrous acid gas, and the colour was deep orange; so that the existence of nitric acid as a *pure body* consisting of 1·5 of oxygen and 2 of nitrous gas is problematical; the gaseous combination of nitrous gas and oxygen probably always contains 2 of nitrous gas, and 1 of oxygen; and some basis seems necessary for the union of 2 of nitrous gas, and 1·5 of oxygen; such as water, alkalies, or oxides.

M. Gay Lussac supposes that there is a compound of three of nitrous gas,* and one of oxygen, capable of

* It is stated that this combination can only be made over a large

combining with water and alkalies without decomposition. I have tried many experiments on this subject, but have never been able to make a strong coloured aquafortis containing more than 2 of nitrous gas to 1 of oxygen in volume; when nitrous acid gas is passed into alkaline solutions, a portion of nitrous gas is always evolved; and when one in volume of oxygen is added to two of nitrous gas, and dry azote introduced to mark the condensation, no change takes place on the mixture of the gas with fresh portions of dry nitrous gas.

Aquafortis or *nitric acid*, is procured for the purposes of chemistry by the distillation of nitre and oil of vitriol; about 2 parts of nitre should be used to 1 part of oil of vitriol, and the retort heated in a sand bath connected with a receiver kept cool by moistened cloths. The acid thus obtained is usually coloured, but becomes pale by exposure to air. If the nitre is dry, its specific gravity is from 1.520 to 1.55. This substance acts with great violence on all the metals anciently known, except gold and platina, and causes volatile oils to inflame. When it is passed through a porcelain tube heated to redness, oxygen is given off from it, and nitrous acid gas; and the same effect is produced upon the residual acid, as if it had been mixed with water; so that it is proved by this experiment to be composed of nitrous acid gas, oxygen, and water; and 4 in volume of nitrous gas, and 2 of oxygen gas condensed in water, I find, absorb 1 in volume of oxygen to become nitric acid.

surface of water, which shews either that the air in the water is concerned in the condensation, or that the water itself absorbs nitrous gas: the *size* of the vessel can have no influence on the compound formed, and it is supposed by M. Gay Lussac, readily absorbable by water. Mem. D'Arcueil, t. ii. p. 241.

7. To enter upon a description of all the experiments that have been made to ascertain the quantities of water in acids of different strengths, would be unfitted to the nature of an elementary treatise.

From my own experiments compared with those of Kirwan, Wenzel, and Berthollet, I am inclined to believe that the strongest acids contain from 14 to 15 per cent. of water, and according to the principles of the French nomenclatures, they ought to be called *hydro-nitric acids*.

Aquafortis, or hydro-nitric acid, when its specific gravity is below 1.4, strengthens by being boiled; when stronger than 1.45, it becomes weaker by boiling. According to Mr. Dalton, the acid of 1.42 distils unaltered at 248° Fahrenheit. It is probable that the acid of 1.55 consists of 1 proportion of water and 1 of acid, and that which rises unaltered at 248° of 1 proportion of acid and 2 of water.

If nitrous gas be considered as represented by 56, that is, by 1 proportion of azote, and 2 of oxygen, 26 and 30, then nitrous acid gas will be represented by 86, or 1 of azote, 26, and 4 of oxygen, 60; and 101 will be the number for the acid contained in the pale acids, and in the salts called nitrates, and it will consist of 1 of azote and 5 of oxygen.

And the strongest acid will contain 17 water and 101 acid, and the acid of 1.42 34 water and 101 acid.

Hydro-nitric acid is of great use in many of the common arts. It is employed in medicine, for dissolving metals, for etching, for making compounds used in dyeing, and it is one of the constituent parts of nitre, a substance essential in the manufacture of gun-powder.

8. Azote and chlorine combine together and form a very singular substance which was discovered by M.

Dulong in 1811. It is produced by exposing chlorine to an aqueous solution of any ammoniacal salt, and suffering them to remain in contact. The chlorine gradually disappears and a yellow oil is formed, which is the compound in question. It must be examined with great care, for it explodes with extraordinary violence at a temperature below that of boiling water: M. Dulong, the discoverer, was deprived of an eye in a detonation produced by it; and in the first attempts that I made to analyse it, I received a severe wound in the eye.

By using a mask, furnished with a plate of thick glass, I have since been able to make many experiments upon it without any accident.

It easily rises in vapour. It does not congeal at 25° below 0 Fahrenheit. Its specific gravity is 1.653. It detonates violently by the contact of phosphorus or of any fixed oil, and half a grain of it exploded under water, produces a loud report, and breaks in pieces the vessels if of earthenware or glass in which the experiment is made.

It is slowly acted upon by mercury, and affords corrosive sublimate and azote.

In solution of muriatic acid it produces more than its own weight of chlorine, and muriate of ammonia is formed.

Its detonation, like that of euchlorine, appears to depend merely upon the separation of its elements. From various experiments, I conclude that it contains 1 volume of azote, and 4 of chlorine, or 1 proportion of azote 26, and 4 proportions of chlorine, 268.

9. Azote and hydrogen exist in combination in ammonia or the volatile alkali. It is not easy to produce their union, yet when azote is exposed to moist substances giving off hydrogen, a little ammonia is

found after some time in the water;—for instance, when azote is placed in contact with moist iron filings above mercury. Priestley first procured ammonia in its pure form; and his experiments, and those of Scheele, repeated and illustrated in an elaborate manner by Berthollet, led to the knowledge of its elements; indeed the last chemist must be considered as the true discoverer of the composition of ammonia.

To procure ammonia, equal parts of well burnt lime and dry sal-ammoniac or muriate of ammonia, are heated in a retort of glass, the beak of which is plunged under dry mercury. Gaseous matter comes over, which, when the common air of the retort has been all expelled, must be collected in inverted jars filled with mercury.

Ammonia at common temperatures is a permanent gas; according to Guyton de Morveau, it becomes a liquid at about 70 below 0 of Fahrenheit's scale: but his experiments were made in glass balloons, and the conclusions drawn from the appearance of fluid; so that the evidence, though strong, cannot be regarded as perfectly satisfactory, as ammonia contains vapour which must be condensed to a great extent by so intense a cold.*

The specific gravity of ammonia is to that of hydrogen, nearly as 8 to 1. 100 cubical inches of it weigh at mean temperature and pressure about 18 grains.

When an inflamed taper is plunged into a jar of ammonia, the light is extinguished; but a slight inflammation of the gas occurs where it is in contact with air.

Its taste is extremely acrid: it cannot indeed be safely applied to the organs of taste or smell except when mixed with much common air. It is the principle which gives pungency to the common concrete volatile alkali.

* [It has been liquefied at 50° by a pressure equal to 6·5 atmospheres.]

It instantly reddens paper tinged with turmeric, and gives a green colour to most vegetable blues and reds; and this property, and its other properties, characterise it as an alkaline body.

It is rapidly absorbed by water. At the temperature of 50° under a pressure equal to 29.8 inches, water, I find, absorbs about 670 times its volume of gas, and becomes of specific gravity, .875.

The following table containing approximations to the quantities of ammoniacal gas in aqueous solutions of different specific gravities, was constructed after experiments made with great care for the purpose.

100 parts of specific gravity.		Of Ammo- nia.
*8750	Contain.	32.5
8875		29.25
9000		26.00
9054*		25.37
9166		22.07
2255		19.54
9326		17.52
9385		15.88
9435		14.53
9476		13.46
9513		12.40
9545		11.56
9573		10.82
9597		19.17
9619		9.60
9692*		9.50

The constitution of ammonia may be easily demonstrated by analytical experiments, it is decomposed by electrical sparks, or by being passed through a tube

* The three results marked by the asterisk were gained by experiments, the other numbers by calculation.

heated to redness; its volume is increased, and it becomes hydrogen and azote.

M. Berthollet, jun., conceives that its volume is doubled when it is decomposed over mercury by electrical sparks. In Dr. Henry's experiments and those that I have made, the expansion is a little less; but this is probably owing to the unavoidable imperfection of the process. I once thought that a small quantity of water was formed in the operation, but very delicate experiments have convinced me that this is not the case. I decomposed a quantity of ammonia by electric sparks in a close vessel, in which its elements could not expand, and yet no moisture was deposited.

In the most accurate experiments in which the hydrogen has been separated from the azote by successive detonations with small quantities of oxygen; the volumes have been 3 of hydrogen to 1 of nitrogen, so that ammonia consists in weight of 3 of hydrogen and 13 of nitrogen, and supposing the number representing hydrogen unity, the same number is gained to represent azote as from the proportions of the elements in its compounds with oxygen; and ammonia consists of 1 proportion of azote and 6 of hydrogen, and is represented by 32. That all these conclusions are correct is shewn by the decomposition of the compound made from nitric acid and ammonia.

When this salt, which is called nitrate of ammonia, is exposed to a heat gradually raised, it is decomposed into water and nitrous oxide; and this could not happen unless it were constituted by definite proportions, which must be 101 of acid and 32 of alkaline matter; for 6 of hydrogen require 45 of oxygen to produce water, and 52 of azote, i. e. 26 in the acid, and 26 in the alkali require 30 of oxygen to produce nitrous oxide.

Ammonia is employed in medicine, and its compounds are used in processes of dyeing, and in some of the metallurgical arts.

10. Azote has not as yet been resolved into any other forms of matter. I volatilized the highly combustible metal potassium in azote over mercury, and passed the Voltaic flame of 2000 double plates through the vapour; but the azote underwent no change. I have made many other attempts to decompose it, but they have been unsuccessful.

The strongest arguments for the compound nature of azote are derived from its slight tendency to combination; and from its being found abundantly in the organs of animals which feed on substances that do not contain it.

Its uses in the economy of the globe are little understood; this likewise is favourable to the idea that its real chemical nature is as yet unknown, and that it is not actually an undecomposable substance.

It would appear that azote and oxygen combine slowly under circumstances in natural operations, when lime or alkaline substances are present; thus nitrous salts are produced in nitre beds in warm climates, and the process is assisted by, or the combination formed during the fermentation of animal and vegetable substances.

III. *Of Sulphur.*

1. Sulphur is a well known substance, found native in many parts of the world, and it may be procured by the distillation of the mineral, called pyrites.

It is brittle, moderately hard, and of a yellowish colour, and has a peculiar taste and smell. It is a non-

conductor of electricity. Its specific gravity is 1990. It is often found in regular solid forms, usually octohedral, when it is more or less transparent. Its power of refracting light, according to Dr. Wollaston, is to that of water as $\cdot 204$ to $\cdot 1336$: its capacity for heat is stated to be as $\cdot 19$ to 1. It fuses at about 220° Fahrenheit, and volatilizes slowly even before it fuses; at the temperature of 560° it becomes an elastic fluid, and in this state inflames if in contact with air, and burns with a pale blue flame.

2. If sulphur be heated above 300° Fahrenheit, it gradually becomes thick and viscid, when if it be poured into a bason of water, it will be found of a red colour, and ductile like wax. In this state it is used for receiving impressions of seals or medals. Its specific gravity is increased, according to Dr. Thomson, to 2325 during this process; from which it seems probable that its parts undergo a new arrangement, and approach nearer, than in its common crystalline form. It has been supposed that the change depends upon its combining with oxygen; but in some experiments made expressly to ascertain this point, it was not found that any oxygen was absorbed when sulphur was long kept heated in contact with it in close vessels, and I have observed after Dr. Irvine, jun., that the change of colour takes place independent of the presence of air.

3. The only well known compound consisting of sulphur and oxygen alone, is a gaseous substance, called in the modern nomenclature, *sulphurous acid gas*. It may be procured by heating sulphur in oxygen gas; the experiment may be performed in a glass retort, and the sulphur inflamed by a spirit lamp; it burns with a beautiful violet flame, and if the oxygen gas has been carefully freed from water, sulphurous acid gas will be

the product. It may be formed likewise by heating mercury or copper filings, in oil of vitriol, and collected over mercury.

Sulphurous acid gas has a very disagreeable smell; it is the smell of burning sulphur. It reddens vegetable blues, and gradually destroys most of them. It whitens many animal and vegetable substances, silk and straw, for instance, and hence the vapours of burning sulphur are employed in bleaching.

Its specific gravity is to that of hydrogen as 30 to 1., and 100 cubical inches of it at mean temperature and pressure, weigh about 68 grains.

It is absorbed by water; this fluid takes up about 30 times its bulk, gains a nauseous subacid taste, and according to Dr. Thomson, becomes of specific gravity 1.0513.

That sulphurous acid gas consists of sulphur and oxygen, is evident from the phenomena of its production by combustion.

In several experiments in which I burnt sulphur, procured from iron pyrites out of the contact of air or moisture, in dry oxygen gas over mercury, I found that the volume of the oxygen was very little altered; the condensation was never more than $\frac{1}{12}$, and seldom so much as $\frac{1}{15}$, and I am inclined to attribute the loss to the formation of a little oxide of sulphur, or to a little hydrogen loosely combined with the sulphur, so that there is every reason to believe that sulphurous acid is constituted by sulphur dissolved in a volume of oxygen.

This conclusion is confirmed by some experiments on the action of cinnabar, which consists of mercury united to sulphur, and sulphur itself, on metallic oxides.

Two equal quantities of red oxide of mercury, each

weighing 10 grains, were heated, one alone, the other mixed with sulphur. They afforded nearly equal volumes of gas. One, which equalled 2 cubical inches and $\frac{2}{10}$, was oxygen; the other, which equalled 2 cubical inches and $\frac{1}{10}$, was pure sulphurous acid gas. Similar experiments were made, cinnabar being substituted for sulphur, with like results.

If the specific gravities of sulphurous acid gas and oxygen be compared, and the last subtracted from the first, it will appear that sulphurous acid consists nearly of equal parts of oxygen and sulphur by weight.

4. If a solution of sulphurous acid gas in water be exposed to the air, it loses its peculiar flavour, and becomes strongly sour; and experiments on the action of the solution on air, shew that oxygen is absorbed.

Sulphurous acid gas is easily driven off from water in the recent solution, but after it has been changed by exposure to air, water only rises when it is heated; and if the evaporation is carried on till the temperature is 546° , the residuum is found to be *oil of vitriol*; nothing but water will have been given off; and therefore oil of vitriol contains sulphur combined with more oxygen than in sulphurous acid. That it likewise contains water, is shewn by another experiment, which, if made with accuracy, affords perfect evidence of its nature and composition. Let a porcelain tube be heated red, and the strongest oil of vitriol passed through it in vapour, a part of it will be decomposed, the gaseous products will be 2 parts of sulphurous acid gas, and 1 part of oxygen gas; and the fluid product will be a weaker acid, such as would be produced by diluting the acid which is the subject of experiment.

The compounds made by adding oil of vitriol to the alkaline earths, free, as far as our knowledge extends,

from water, give off moisture when they are heated to redness, and if the quantity of water in the strongest oil of vitriol, be estimated from experiments of this kind, it may be concluded that it contains about 19 per cent. of water; and its composition may be thus expressed, 30 of sulphur, 45 of oxygen, and 17 of water.

In the common process of manufacture, oil of vitriol is made by burning sulphur mixed with about $\frac{1}{7}$ of its weight of nitre in pans of iron or lead communicating with a chamber of lead, the bottom of which is covered to the depth of several inches with water. The true theory of this process is not given in any chemical book.

The sulphur by burning, forms sulphurous acid gas, and the acid in the nitre is decomposed, giving off nitrous gas; this coming in contact with the oxygen of the atmosphere, produces nitrous acid gas, which has no action upon sulphurous acid, to convert it to sulphuric acid, unless water be present, and if this substance be only in a certain proportion, the water, the nitrous acid gas, and the sulphurous acid gas, combine, and form a white crystalline solid. By the large quantity of water usually employed, this compound is instantly decomposed, oil of vitriol formed, and nitrous gas given off, which in the air, again becomes nitrous acid gas, and the process continues according to the same principle of combination and decomposition, till the water at the bottom of the chamber is become strongly acid. It is easy to prove the truth of these reasonings; let dry sulphurous acid gas, and nitrous acid gas, be mixed together, by suffering the sulphurous gas to pass into a glass globe partially exhausted, and containing nitrous acid gas. There will be no action between the gases. But if a drop of water be introduced, there will be an immediate condensation, and a beautiful white crystal-

line solid will line the interior of the vessel. Whereas if the globe contain plenty of water, nitrous gas will be given off with great violence, and the water will be found to be a solution of oil of vitriol.

In the solid crystalline compound, it is evident, from the products of its decomposition by water, there must be four volumes of sulphurous acid gas, and three of nitrous acid gas, probably in two or three proportions, with a single proportion of water; for nitrous acid gas contains $\frac{2}{3}$ of its volume of loosely combined oxygen, and sulphurous acid gas requires half its volume of oxygen to become, when condensed in water, solution of oil of vitriol.

Mr. Dalton, who has adopted MM. Clement's and Desormes' idea of nitrous acid gas being decomposable by sulphurous acid gas, which is not correct, supposes that there is a solid sulphuric acid, the oxygen in which is to that in sulphurous acid, as 3 to 2; but the body which he supposes to be sulphuric acid is the crystalline substance, the nature of which is demonstrated above: and no substance to which the name of pure sulphuric acid ought to be given, i. e. a substance consisting of 30 of sulphur and 45 of oxygen, has yet been discovered in an insulated state.

The term sulphuric acid, is improperly applied to the strongest oil of vitriol; this substance, according to the principles of the French nomenclature, ought to be called *hydrosulphuric acid*.

The oil of vitriol of commerce, which is of specific gravity 1.85, rises in vapour at about 550° Fahrenheit, and distils unaltered; whereas weaker acids, by being boiled, lose water, and are brought to this state of concentration. There is a diluted acid of specific gravity 1.78, which congeals at any temperature below 46°

Fahrenheit. This acid, as Mr. Dalton has stated, contains exactly twice as much water as the acid of 1·85. It is composed, according to my experiments, of 30 of sulphur, 45 of oxygen, and 34 of water.

Pure oil of vitriol is a corrosive substance. It acts with great energy upon animal and vegetable matter. It rapidly attracts moisture from the air, and produces much heat when mixed with water. It reddens vegetable blues; and acts with great violence upon alkaline substances, and upon certain earths and metallic oxides; and neutral salts are produced by the union of its sulphur and oxygen with these bodies.

The number representing sulphur, as learnt from the constitution of sulphurous acid gas, is nearly 30; and as this gas contains two proportions of oxygen twice 15, it would seem probable that an oxide of sulphur may exist, consisting of 30 of sulphur, and 15 of oxygen.

I have examined some highly coloured specimens of Sicilian sulphur, which seemed to contain oxygen, and as has been just stated, it is possible that a little oxygen may be condensed in the combustion of sulphur in the residuum; but as yet no body is known that can with propriety be called *oxide of sulphur*.

5. Sulphur and chlorine are possessed of a chemical attraction for each other. The first combination of them was made by Dr. Thomson in 1804, by passing chlorine over flowers of sulphur. It may be made more expeditiously by heating sulphur in a retort filled with chlorine. The sulphur and the chlorine unite and form a fluid substance, which is volatile below 200° Fahrenheit, and distils into the cold part of the retort. This substance, seen by reflected light, appears of a red colour, but is yellowish green, when seen by transmitted light. It smokes when exposed to air, and has an

odour somewhat resembling that of sea-weed, but much stronger; it affects the eyes like the smoke of peat. Its specific gravity, according to Dr. Thompson, is 1.6.

It does not redden perfectly dry paper tinged with litmus: when it is agitated in contact with water, the water becomes cloudy from the appearance of sulphur, and strongly acid, and it is found to contain oil of vitriol.

According to my experiments, 10 grains of pure sulphur absorb nearly 30 cubical inches of chlorine; so that the compound contains about 30 of sulphur to 68.4 of chlorine; 20 of sulphur to 67 of chlorine, would give 1 proportion of sulphur to 2 of chlorine; which, there is every reason to believe, must be the just estimation; for my experiments were made in retorts furnished with metallic stop-cocks, by which a little chlorine must have been absorbed.

The compound formed in the manner above described cannot be made to unite to more chlorine; but I find it dissolves a considerable portion of sulphur by heat, and becomes of a tawney yellow colour.

Dr. Thomson called this substance sulphuretted muriatic acid, but there is no proof that it contains muriatic acid. According to an idea which I ventured to propose in the Philosophical Transactions for 1811, that of calling the compounds of chlorine by the name of their bases, with a termination in "ane" its name would be *sulphurane*.*

6. Sulphur and hydrogen combine. Their union may be effected by causing sulphur to sublime in dry hydrogen in a retort. There is no change of volume: but only a part of the hydrogen can be combined with sulphur in this mode of operating.

* [It is now called chloride of sulphur.]

The gaseous compound of sulphur and hydrogen was discovered by Scheele, in 1777. It is usually made by the action of diluted sulphuric acid upon a mixture of 3 parts of iron filings, and 2 parts of sulphur that have been ignited together; for the purposes of accurate experiments, it should be collected over mercury.

Sulphuretted hydrogen inflames when a lighted taper is brought in contact with it, exposed to the air: it burns with a pale blue flame, depositing sulphur. Its smell is extremely fetid, resembling that of rotten eggs. Its taste is sour. It reddens vegetable blues. It is absorbable by water; that fluid takes up more than an equal volume of gas. Its specific gravity, according to MM. Gay Lussac and Thenard, is to that of air as 1.1912 to 1. From my experiments it would appear to be a little less; but I am inclined to adopt the results of the French chemists rather than my own, as their gas was weighed in larger quantity, and dried. Its weight to that of hydrogen may be considered as 16 to 1, and 100 cubical inches of it, at mean temperature and pressure, weigh between 36 and 37 grains.

The composition of sulphuretted hydrogen is demonstrated by the change produced in it by electricity; if platina wires be ignited in it by the voltaic apparatus, it is rapidly decomposed. Sulphur is deposited, and an equal volume of hydrogen remains; the same change is effected more slowly by electrical sparks.

The proportion of its elements are shewn to be the same, both by the analytical and synthetical experiments. They must be 15 of sulphur to 1 of hydrogen; and the results give as nearly as possible the same number representing sulphur, as its compounds with oxygen and chlorine: and sulphuretted hydrogen may

be considered as consisting of two proportions of hydrogen 2, and 1 of sulphur, 30.

This body combines with an equal volume of ammonia; and unites to alkalies and oxides, so that it has all the characters of an acid.

7. There is another compound of hydrogen and sulphur, the proportion of the elements of which have not yet been accurately ascertained; but it probably will be found to contain at least one proportion more of sulphur.

It may be procured by decomposing, by muriatic acid, the substance formed by melting sulphur and hydrate of potassa together. It is a fluid body, and was first described by Berthollet in 1808. It affords sulphur and sulphuretted hydrogen when heated. Very few experiments have been made upon its properties, and none upon the proportions of its elements.*

8. Sulphur has no chemical attraction for azote, at least no compound of these bodies has as yet been formed.

9. Sulphur has been placed amongst the undecomposed bodies, because as yet nothing certain is known respecting its elements. When Sicilian sulphur was fused and exposed to the action of platina points intensely ignited by Voltaic electricity, excited by 1000 double plates, permanent gas was given off from it, which proved to be sulphuretted hydrogen: a small quantity of sulphuretted hydrogen is given off likewise during the action of copper filings upon sulphur; and much hydrogen is evolved during the action of sulphur on the compound of potassium and tellurium. It may

* [It is a bisulphuret according to Dr. Dalton; from the later experiments of M. Thenard, it appears to be a persulphuret, variable in the proportions of its constituents.]

however be questioned whether hydrogen is essential to the constitution of sulphur. Sulphur may possibly contain in its common forms a little moisture, or a little of a solid compound of hydrogen and sulphur; and till the gas can be separated from it in definite proportion, and be proved to be combined with some other matter, no accurate conclusions can be formed upon the subject.

Sulphur is employed in medicine, particularly as an external application in cutaneous complaints. Its use in bleaching has been already referred to. Its most important application is in oil of vitriol, and the compounds formed from it, which are used in various processes of dyeing and calico printing.

IV. *Of Phosphorus.*

1. *Phosphorus* was discovered by Brandt in 1669. It may be made by the following process.

100 parts of burnt bones in powder, are to be mixed with 40 parts of oil of vitriol, and they are to be suffered to remain in contact for a couple of days, the mixture being frequently stirred. The whole is then to be poured upon a filter of cloth, and the liquor that passes through is to be added to a nitrous solution of lead; a white powder will be formed; this must be mixed with about $\frac{1}{5}$ of its weight of charcoal powder, and exposed to a strong red heat in a porcelain retort, the beak of which is plunged in water; much gaseous matter will come over, some of which will inflame spontaneously, and at length a substance will drop out of the neck of the retort, and congeal under the water, which is phosphorus. It may be purified by melting it

in water, and passing it under water through chamois leather.

2. Phosphorus is semi-transparent, and of a yellowish colour. It is as soft as and more ductile than wax. It is insoluble in water. Its specific gravity is about 1.77. It melts at the temperature of 90° , and boils at about 550° .

When phosphorus is exposed to air at common temperatures, it emits a white smoke, which appears luminous in the dark. This depends upon its combining with oxygen, and forming an acid which unites to the aqueous vapour in the atmosphere, and they fall down in the fluid form. Phosphorus, I find, does not smoke in air perfectly dry; and in this case the acid adheres to it, and in a short time prevents it from being luminous.

When phosphorus is heated to about 148° , it takes fire, and burns with intense brilliancy, throwing off dense white smoke, which is a strong solid acid that soon becomes liquid by attracting moisture from the air; and a red substance usually remains.

3. The manner in which phosphorus acts upon the air, as has been shewn, page 170, proves that it is capable of combining with oxygen; and there is every reason to believe in at least 3 proportions.

When phosphorus is inflamed in oxygen gas over mercury, and the white substance produced strongly heated, the oxygen being in excess, for every grain of phosphorus burnt four cubical inches and a half of oxygen gas are absorbed. The substance so procured is called *phosphoric acid*. It becomes fluid at a red heat; it is not volatile even at a white heat. It has no smell; its taste is intensely, but not disagreeably acid. It dissolves in water, producing great heat; and its

saturated solution is of the consistence of syrup. It acts upon, and corrodes glass, and unites to alkalies and oxides.

4. When phosphorus is heated in highly rarefied air, three products are formed from it; one is phosphoric acid: one is easily volatile, and appears as a white powder; and the other is a red solid, comparatively fixed, and requiring a heat above that of boiling water for its fusion. The volatile substance is soluble in water, and gives it acid properties. It contains less oxygen than phosphoric acid; for it burns and becomes fixed when heated strongly in the air. Its taste is sour, with a peculiar pungency, and it emits a smell not unlike that of garlic. It is mixed with phosphorus, but is principally the substance which, according to the French nomenclature, should be called *phosphorous acid*, and which in chemical works, is inaccurately described as a fluid body. The red substance requires less oxygen than phosphorus to convert it into phosphoric acid, and must be considered as an *oxide of phosphorus*.

I have never been able to procure phosphorous acid by combustion free from admixture or combination with other substances. In the common mode in which it is said to be obtained, namely, by exposing phosphorus to free air; there is always a large quantity of phosphoric acid formed.

A pure solid *hydro-phosphorous acid*, that is, a combination of it with water, may, I find, be produced by the following process.

Phosphorus is sublimed through corrosive sublimate in powder in a glass tube; a limpid fluid comes over, which must be mixed with water, and the solution heated till it is of the thickness of syrup. It is a combination of water and pure phosphorous acid. It red-

dens vegetable blues, and combines with alkalies, and has all the characters of a strong acid. It forms a white crystalline solid on cooling.

It becomes solution of phosphoric acid slowly when exposed to air, absorbing oxygen. When it is gently heated, it takes fire, and burns with great brilliancy, emitting globules of gas that inflame in contact with air; a red oxide of phosphorus is deposited in the bottom of the vessel, and solid phosphoric acid is formed.

The substance produced by passing phosphorus through corrosive sublimate, as will be immediately shewn, is a compound of phosphorus and chlorine; and when it acts upon water, hydrogen is afforded to the chlorine, and oxygen to the phosphorus; there are no products but muriatic acid gas and phosphorous acid, and the quantity of hydrogen in the muriatic acid gas formed, being known, the quantity of oxygen in the phosphorous acid is likewise known. By two experiments made with great care, in which the quantity of chlorine in the liquor from the phosphorus and corrosive sublimate, was estimated by its combination with silver, I ascertained that ten grains of phosphorus required for their conversion into phosphorous acid, such as exists in the hydrate just described, 7·7 grains of oxygen; and it is evident from this result, compared with that on the combustion of phosphorus, in which phosphoric acid is formed, that the quantity of phosphorus being the same, it requires twice as much oxygen to become phosphoric acid, as it requires to form phosphorous acid; and from these data, the number representing phosphorus, must be regarded as about 20, and phosphorous acid will consist of 20 phos-

phorus, and 15 oxygen, or 35 ; and phosphoric acid of 20 phosphorus and 30 oxygen, or 50.

That the hydro-phosphorous acid is a compound of phosphorus, oxygen, and water, is shewn by heating it in contact with ammonia over mercury; the ammonia unites to the pure acid, and water is expelled. I find by experiments on the quantity of water it affords, that it consists of four proportions of phosphorous acid, and two of water.

I have made no experiments on the proportion of oxygen in the red oxide. It possibly will be found to consist of two proportions of phosphorus, and one of oxygen.

5. Phosphorus and chlorine combine with great facility when brought in contact with each other at common temperatures; and compounds may be formed from their union, containing different proportions of the two elements.

When chlorine is introduced into a receiver exhausted of air, and containing phosphorus, the phosphorus takes fire and burns with a pale flame throwing off sparks, and a white substance rises and condenses on the sides of the retort.

If the chlorine be in considerable quantity, as much as twelve cubical inches to a grain of phosphorus, the phosphorus will entirely disappear, and nothing but the white powder will be formed; and about 9 cubical inches of the chlorine will be found to be absorbed; and no new gaseous matter will be produced.

The powder is a compound of phosphorus and chlorine. I first described it as a peculiar body in 1810, and various analytical and synthetical experiments which I have made, prove that it is composed of about 1 of phosphorus and 6.8 of chlorine in weight.

Its properties are very peculiar. It is a snow-white

substance. It is very volatile, and rises in a gaseous form at a temperature much below that of boiling water; under pressure it may be fused, and then it crystallizes in prisms which are transparent.

It acts violently upon water, which it decomposes. Its phosphorus combines with the oxygen, producing phosphoric acid, and its chlorine with hydrogen forms muriatic acid.

It produces flame when exposed to a lighted taper; and when passed through a glass tube heated red, with oxygen, is decomposed; the oxygen forms phosphoric acid with the phosphorus, and the chlorine is disengaged. Dry litmus paper exposed to its vapour in a vessel exhausted of air is reddened. It combines with ammonia when it is introduced into a vessel containing it, with much heat; and they form together a compound insoluble in water, indecomposable by acid or alkaline solutions, and having characters analogous to an earth.

It is evident from the analysis, that it consists of two proportions of chlorine, supposing the number representing chlorine, 67, or of four, supposing it 33·5, 134 to one of phosphorus 20, and the number representing it is analogous to an acid in many of its properties. According to the principles of nomenclature which I have ventured to propose, its name will be *phosphorana* [per-chloride].

6. I have already referred to the substance produced by passing phosphorus through corrosive sublimate. It is a fluid as clear as water; its specific gravity is to that of water as 1·45 to 1. It may be called *phosphorane* [chloride of phosphorus]. I first obtained it in a pure form, in 1809. It appears from the circumstances already detailed, that it consists of 1 proportion of phosphorus 22, and 1 of chlorine 67, and the

number representing it is 87.* It emits acid fumes when exposed to air, decomposing the vapour in the air, and if made to moisten paper, it is converted into acid in the air, without any inflammation. It does not redden dry litmus paper plunged into it; the vapour from it burns in the flame of a candle; its action upon water has been already referred to. When it is introduced into a vessel containing chlorine, it is converted into phosphorana: if made to act upon ammonia, phosphorus is produced, and the same compound as that formed by phosphorana and ammonia.

7. When phosphorus is gently heated in phosphorane, a part of it dissolves, and the fluid, when exposed to the air gives off acid fumes from its action upon the vapour it contains, and a thin film of phosphorus is left behind, which usually inflames by the heat generated from the decomposition of the vapour. The first compound of this kind was obtained by MM. Gay Lussac and Thenard, by distilling phosphorus and calomel together in 1808, and they imagined it to be a peculiar combination of phosphorus, oxygen, and muriatic acid.

No experiments have been as yet made to determine the quantity of phosphorus which phosphorane will dissolve; probably a definite combination may be obtained, in which the proportions of chlorine will correspond to the proportions of oxygen in the oxide of phosphorus.

8. An elastic fluid, which has the peculiar property of inflaming in contact with the atmosphere, may be procured by heating together slacked lime, or strong solution of potassa or soda, and phosphorus. It is expedient to deprive the air contained in the vessel in

* 13·6 grains of it decomposed by nitrate of silver, afforded 43 grains of hornsilver, and 100 grains of silver absorb 32·5 of chlorine to become hornsilver.

which it is generated of oxygen, by burning phosphorus or a taper in it; the gas should be preserved over mercury; it soon becomes adulterated by exposure to water containing air.

This gas differs in properties according to the manner in which it is made; I have obtained it from phosphorus and alkaline lixivia, of specific gravities varying from 4 to 7, 1 being the standard of hydrogen: its smell is very disagreeable: water absorbs about $\frac{1}{40}$ of its volume of the gas. It detonates when brought into contact with chlorine, producing a brilliant green light; but the results of the detonation have never been minutely examined. It explodes with a most intense white light in oxygen gas; the heaviest spontaneously inflammable gas that I have ever made, absorbed rather less than an equal volume of oxygen.

When electrical sparks are passed through gases of this kind for a long time, a reddish film which burns like phosphorus is deposited; usually there is no change of volume, and the remaining gas is hydrogen. When a gas, the specific gravity of which was 6, was heated for some time over mercury in contact with zinc filings, there was an expansion of volume to more than $\frac{1}{3}$; a substance was formed superficially on the zinc, which had the characters of a compound of phosphorus and zinc. There was an expansion when finely divided platina was heated in a portion of the same gas. Potassium in excess made to act upon it by a spirit-lamp produced a rapid increase of its volume; 2 parts became rather more than 3. The potassium was affected as it would have been by a union with phosphorus, and the gas was found to be pure hydrogen.

This substance, which was discovered by M. Gen-gembre in 1783 has been called *phosphoretted hydrogen*,

9. When solid hydrat of phosphorous acid is heated in a retort out of the contact of air, solid phosphoric acid is formed, and a large quantity of elastic fluid is generated, which has very peculiar properties; I discovered it at the same time as the solid hydrat of phosphorous acid, namely, in February 1812. This gas has a disagreeable smell, but not nearly so fetid as that of phosphoretted hydrogen: it does not explode spontaneously, but detonates violently when heated in contact with oxygen to about 300° Fahrenheit. It explodes in chlorine with a white flame. Water absorbs $\frac{1}{8}$ of its volume of this gas. In an experiment in which a small quantity only was weighed, its specific gravity appeared to be to that of hydrogen as about 12 to 1.

When potassium is heated in it, its volume is doubled, and the gas produced is pure hydrogen. When sulphur is sublimed in 1 volume of it, a sulphuret of phosphorus is formed, and nearly 2 volumes of sulphuretted hydrogen produced. When detonated with oxygen in excess, 3 in volume of it absorb more than 5 in volume of oxygen, and a little phosphorus is always thrown down; when 8 of it in volume are detonated by an electrical spark with 2 of oxygen, there is a considerable deposition of phosphorus, and 9 of gas, which has the odour of common phosphoretted hydrogen, remain; 1 volume of it absorbed nearly 4 volumes of chlorine.

I venture to propose the name of *hydroposphoric* gas for this elastic fluid. It appears to be composed of 1 proportion of phosphorus and 4 of hydrogen, 2 volumes of hydrogen being compressed in the space of 1, and the number representing it is 24.

It is probable that the gas called phosphoretted hydrogen sometimes contains this gas, mixed with com-

mon hydrogen, and perhaps a *peculiar elastic* fluid, consisting of 1 proportion of phosphorus, and 2 of hydrogen, which has the property of spontaneous inflammation. Hydrophosphoric gas, I find, does not become spontaneously explosive by mixture with hydrogen.

There is not, perhaps, in the whole series of chemical phenomena, a more beautiful illustration of the theory of definite proportions, than that offered in the decomposition of hydrophosphorous acid into phosphoric acid and hydrophosphoric gas.

4 proportions of the acid contain 4 proportions of phosphorus, and 4 of oxygen; 2 proportions of water contain 4 proportions of hydrogen, and 2 of oxygen. The 6 proportions of oxygen unite to 3 proportions of phosphorus to form 3 of phosphoric acid, and the 4 proportions of hydrogen combine with 1 of phosphorus to form 1 proportion of hydrophosphoric gas; and there are no other products.

10. Phosphorus and sulphur are capable of combining; they may be united by fusing them together in a tube exhausted of air, or under water; but in the last case they must be used only in small quantities; as at the moment of their action water is decomposed, sometimes with explosions. This compound, which has been called sulphuret of phosphorus, was described by Margraff in 1762. He formed it of equal parts of the two substances, but phosphorus and sulphur may be united into one mass in a variety of proportions; and these mixtures are more fusible than either of their constituents. The most fusible compound I have found, is that formed by the two bodies united in the proportion of one and a half of sulphur, to two of phosphorus. This remains liquid at 40° of Fahrenheit; and would appear to be composed of one proportion of sulphur 30,

and two of phosphorus 40. When solid its colour is yellowish-white. It is more combustible than phosphorus, and rises undecomposed by a strong heat.

The points of fusion and evaporation of phosphorus and sulphur, are so near each other, that it is not easy to ascertain the difference between two chemical combinations of these bodies in different proportions, and mixtures of the chemical compounds, with the bodies themselves; 8 parts of phosphorus in weight united to 1 of sulphur, remain fluid at 68° of Fahrenheit; and 1 of phosphorus with 3 of sulphur, congeal at about 100° .

11. When phosphorus is fused and exposed to the action of the Voltaic spark, taken by means of platina wires, phosphoretted hydrogen in small quantities is produced from it; but there are no proofs that hydrogen is essential to its existence; and phosphorus in its common state, may contain a minute portion of the hydruret of phosphorus mixed with it; it would be very difficult to detect in phosphoric acid, the small quantity of water that this hydrogen would produce; and the red colour which phosphorus sometimes possesses, seems to be owing to an admixture of small quantities of oxide of phosphorus. There are some analogies that favour the idea of the compound nature of phosphorus, which will be discussed in the progress of this work; but in the arrangement of the facts of the science, it must be still regarded as an undecomposed body.

12. Phosphorus has not as yet been applied to any of the purposes of the common arts; but various preparations of it are employed for producing quick inflammation. One of the best, is a sulphuret of phosphorus, containing two of phosphorus to one of sulphur; a little of it applied to a common brimstone match, inflames when gently rubbed.

V. *Of Carbon or Charcoal, and the Diamond.*

1. The name carbon signifies the pure inflammable part of charcoal, lamp-black, and other similar substances. The purest known form in which it can be obtained, is by passing oils or spirit of wine through ignited tubes. It then appears as an impalpable black powder; it has no taste nor smell; it is a conductor of electricity; it is more than twice as heavy as water. For the common purposes of experiments, the charcoal of light wood, such as the alder, that has been exposed to boiling water, and afterwards ignited to whiteness, is sufficiently pure. Such charcoal, however, rapidly attracts moisture from the atmosphere, so as to increase in weight from 12 to 14 per cent., and when dry, absorbs several times its volume of any gas, to which it may be exposed, and it must therefore be employed immediately after ignition, and whilst yet warm.

Carbon, whether coherent in charcoal, or in powder, is infusible by any heat that has hitherto been applied. I have exposed it to the powers of intense ignition of different Voltaic batteries; that of Mr. Children, mentioned page 109, one of 40 double plates, of 18 inches square, and the battery of 2000 double plates of 4 inches, both in vacuo, and in compressed gases on which it had no power of chemical action. A little hydrogen was given off from it, and it slowly volatilized in these experiments, and the part remaining was much harder than before, so as in one case to scratch glass, and the lustre was greater; but its other properties were unaltered, and there was no appearance of fusion. Its capacity for heat, according to Dr. Crawford, is to that of water as .2631 to 1.

2. There are two distinct combinations of carbon and oxygen; viz., carbonic acid and carbonic oxide.

Carbonic acid is formed whenever charcoal or carbonaceous matter is burnt in air or oxygen, and it is evolved during fermentation, by the decomposition of animal or vegetable substances, and from limestones by ignition, or the action of acids.

The most expeditious mode of procuring it for chemical purposes, is by the action of weak solution of muriatic acid on coarsely powdered marble. It may be collected over water, or, for accurate experiments, over mercury.

Carbonic acid gas was the first elastic fluid certainly distinguished from air; the knowledge of its acid nature is owing to Dr. Black, who discovered it in 1755. M. Lavoisier, nearly thirty years afterwards, ascertained its composition.

Carbonic acid gas extinguishes flame, has a peculiar sharp taste, and a faint but agreeable smell. It is not respirable. Its specific gravity is to that of hydrogen as 20·7 to 1. 100 cubical inches weigh at the mean temperature and pressure, 47 grains.

Carbonic acid gas is absorbed by water, which unites to its own volume of the gas at 41°. By heat it is expelled from the water.

If carbonic acid gas be poured from one vessel into another vessel containing a lighted taper, the flame is extinguished.

It reddens litmus paper, and combines with alkalies, alkaline earths, and metallic oxides.

A synthetical proof of the composition of carbonic acid gas, has been already given, page 74.

Common charcoal, even when very well burnt, contains a little hydrogen, and affords a minute quantity of water in its combustion; but the charcoal from the

decomposition of oils gives carbonic acid gas alone. It burns when inflamed in dry oxygen with brilliant scintillations; there is no perceptible change in the volume of the gas; and when the process is complete, the oxygen is found converted into carbonic acid gas.

The proportions of the elements in carbonic acid gas are easily learned by the difference between its weight and that of oxygen. This difference proves, as has been stated before, that it must consist of 13 of charcoal to 34 of oxygen.

The constitution of carbonic acid gas is proved analytically by its action upon potassium. If this metal is strongly heated in a retort containing the gas, it takes fire, and burns with a red light. Charcoal in fine powder is deposited, the gas disappears, and oxygen is found added to the potassium.

3. The compound of carbon and oxygen, containing less oxygen than carbonic acid gas, may be formed in many modes; as by igniting chalk or any substance containing carbonic acid with charcoal, zinc filings, iron, or tin; or by igniting difficultly reducible metallic oxides with charcoal, or by passing carbonic acid gas over charcoal heated to whiteness, in a porcelain tube. In this last case, the composition of the gas is shewn by the circumstances of the experiment; charcoal disappears, and the carbonic acid becomes *carbonic oxide gas*, and there is a considerable expansion. The true nature of this elastic fluid was discovered by Mr. Cruikshank in March, 1801.

Carbonic oxide may be purified from the carbonic acid with which it is usually mixed, by washing in lime water.

It is combustible, and by the contact of an inflamed or ignited body, burns in the atmosphere with a lam-

bent blue flame. Its specific gravity, according to Cruikshank, is to that of hydrogen as 13·2 to 1. 100 cubical inches weigh about 30 grains.

Carbonic oxide may be taken into the lungs, but is fatal to animal life. I once took three inspirations of it mixed with about one-fourth of common air; the effect was a temporary loss of sensation, which was succeeded by giddiness, sickness, acute pains in different parts of the body, and extreme debility; some days elapsed before I entirely recovered.*

Water absorbs about $\frac{1}{50}$ of its bulk of carbonic oxide.

Chlorine has no immediate action on carbonic oxide, when they are exposed to each other in common daylight over dry mercury; not even when the electric spark is passed through them. MM. Gay Lussac, Thénard, and Mr. Murray, have asserted that they do not act on each other even when long exposed to the direct solar beams. But experiments which I have seen made by my brother, Mr. John Davy, prove the contrary; they rapidly combine under this circumstance, and when in equal volumes, are condensed to one-half; and form a peculiar gas, which he has discovered is possessed of very curious properties, approaching to an acid in its nature.

The nature of carbonic oxide, and the proportions of its elements, are easily demonstrated by analytical experiments. When two in volume of it are mixed with one in volume of oxygen, and an electrical spark passed through the mixture, an inflammation takes place, and two in volume of pure carbonic acid are formed, and there is no other product.

When potassium is strongly heated in it, combustion

* [These effects he experienced in breathing carburetted hydrogen. Vide Vol. III. p. 278.]

takes place, charcoal is deposited, no gas is disengaged, and oxygen is added to the potassium.

From the experiments on carbonic acid and carbonic oxide, it is evident that the number representing carbon is about 11·4; and carbonic acid is represented by 30 added to 11·4, or 41·4; and carbonic oxide by 15 added to 11·4, or by 26·4.

Some chemists have been perplexed to find a reason why carbonic oxide, which contains more carbon, is lighter than carbonic acid; but as Mr. Dalton has ingeniously and justly observed, there is no difficulty in this circumstance; carbon in the gaseous state, is probably considerably lighter than oxygen. The specific gravity of gases bears no relation to the density of the fluids or solids, from which they are formed; ether is lighter than water; but the vapour rising from it is much heavier than steam. If carbonic oxide be supposed to be constituted by equal volumes of gaseous carbon and oxygen, occupying the space of two in volume, then the specific gravity of gaseous carbon will be to that of oxygen as 13 to 17; or if the constitution of the carbonic oxide is similar to that of the nitrous oxide, it will be only as 6·5 to 17.

4. No compound of carbon and chlorine has been as yet discovered. They have no action on each other under any circumstances to which they have been exposed.*

5. There are two compounds of carbon and hydrogen, which are perfectly distinct and well characterized bodies.

One of them, which has been called *carburetted hydrogen*, is disengaged in certain natural operations, particularly during the decomposition of vegetable sub-

* [Carbon has since been combined with chlorine; four compounds are now known.]

stances; it is the gas evolved in stagnant waters. It may be procured by the distillation of coal that burns with flame, and by decomposing the salt called acetate of potash by a red heat; it should be washed with lime water to separate it from carbonic acid.

It burns with a bright yellowish flame. It has no taste, but a disagreeable empyreumatic smell. Water absorbs about $\frac{1}{30}$ of its volume. Its specific gravity, in its purest form, is to that of hydrogen as rather less than 8 to 1. 100 cubical inches weigh about 17 grains.

When one of this gas in volume is mixed with two of oxygen gas, and an electrical spark passed through them over mercury, water and about one in volume of carbonic acid are the products. Hence one in volume of carburetted hydrogen must contain two in volume of hydrogen gas, and as much carbon as will form a volume of carbonic acid. This likewise is shewn by the phenomena of its electrization. When points of platina are electrically ignited in it, or sparks passed through it, charcoal is deposited, and double its volume of hydrogen is produced. When it is mixed with twice its volume of chlorine over mercury, and acted on by the electrical spark, an inflammation takes place, charcoal is deposited, there is a considerable expansion, and about four volumes of muriatic acid gas are produced.

It is evident from these different results that carburetted hydrogen may be considered as composed of one proportion of carbon 11.4, and four of hydrogen 4, and the number representing it will be 15.4.

6. When a mixture of four parts of oil of vitriol and one part of strong spirit of wine, or alcohol is heated in a retort, a gas is generated, which, when washed by water, is found to be a peculiar gaseous compound of carbon and hydrogen; it has been called *olefiant gas*, and likewise *super-carburetted* hydrogen. It burns,

when kindled, with a beautiful white flame of intense splendour. According to Dalton, water absorbs $\frac{1}{8}$ of its volume of the gas. Its specific gravity is to that of hydrogen nearly as 13 to 1; 100 cubical inches of it weigh between 29 and 30 grains.

When it is mixed with an equal volume of chlorine, the two gases condense each other, and a peculiar fluid is formed, which has been supposed to be an oil; but which is a *peculiar compound*, not soluble in water, and composed of hydrogen, carbon, and chlorine. The nature of olefiant gas may be easily demonstrated, and likewise the proportion of its elements. If pure sulphur be sublimed in the gas in a glass tube over mercury, there is a great expansion; sulphuretted hydrogen is formed, and charcoal deposited; one volume of gas forms about two in volume of sulphuretted hydrogen: the sulphur must not be used in much larger quantity than is sufficient to unite to the hydrogen; for in this case, by the long application of heat, the volume is more than doubled; two grains of sulphur and a cubical inch of gas are proper proportions.

The gas is decomposed by electrical sparks; one volume of it expands to about two; charcoal is deposited, and the expanded gas is found to be hydrogen.

It detonates with great violence by the electrical spark, when mixed with three times its volume of oxygen; water and nearly two volumes of carbonic acid are formed in this process.

When it is detonated with an equal volume of oxygen, it expands greatly, and the two gases together become more than three volumes and a half. In this case only the $\frac{1}{8}$ or $\frac{1}{10}$ of a volume of carbonic acid gas is formed, but more than a volume and a half of carbonic oxide; a little hydrogen is consumed, but the

greatest part remains untouched and mixed with the carbonic oxide; and it may be separated by combustion with chlorine.

If an experiment of this kind could be made without the production of any carbonic acid, or the consumption of any hydrogen, the volume of the gases would be exactly doubled, and they would consist of equal parts of carbonic oxide and hydrogen.

It is evident from all these experiments that olefiant gas may be considered as constituted by two proportions of carbon 22·8 and 4, of hydrogen 4, and the number representing it is 26·8: and supposing a double volume of gaseous carbon in olefiant gas, its specific gravity will be found to be the same as from the data presented by carbonic oxide.

7. Most of the gases that form carbonic acid in burning were noticed by Dr. Priestley, who confounded them under the general name of heavy inflammable air. Olefiant gas was first described as a specific substance, in 1794, by Bondt, Deiman and a society of Dutch chemists. M. Berthollet and Mr. Murray suppose that there is a great variety of gases which consist of oxygen, hydrogen, and carbon, in different proportions; but the experiments of Mr. Dalton, Dr. Henry, and Dr. Thomson, are entirely opposed to these views; and the researches which I have made in conjunction with my brother Mr. John Davy, have convinced me of the correctness of Dr. Henry's opinion, that what have been called different *oxicarburetted hydrogen gases* are merely mixtures of olefiant gas, carburetted hydrogen, carbonic oxide, and hydrogen gases. We used chlorine for separating olefiant gas at common temperatures, and the same substance for separating hydrogen by explosion, or the action of light; and sulphur for decompos-

ing the carburetted hydrogen: and in these modes of analysis our results were unequivocal.*

8. Carbon and azote have no known action on each other.

9. Carbon and sulphur combine with each other and form a very peculiar compound, which was called by its discoverer Lampadius, alcohol of sulphur. It was first formed by Lampadius, during a process on the distillation of wood containing pyrites, and was afterwards produced by Messrs. Desormes and Clement, by passing sulphur in vapour over charcoal strongly ignited in a porcelain tube. MM. Desormes and Clement first ascertained its nature, and their researches have been confirmed by those of Berthollet, Thenard, and Vauquelin, and Berzelius and Marcet.

The alcohol of sulphur or sulphuret of carbon in its common state is yellow, from holding sulphur in solution, but when purified by distillation at a gentle heat, it is a limpid colourless fluid of specific gravity 1.272. Its refractive power is 1.645. It is an extremely volatile substance, and boils at the common atmospheric pressure at a temperature between 105 and 110. It is very inflammable, and the vapour exploded with oxygen gas in excess affords sulphurous and carbonic acid gases. From the experiments of Berzelius and Marcet, it appears that the alcohol of sulphur or sulphuret of carbon consists of 84.83 of sulphur and 15.17 of carbon, which agrees very nearly with 2 proportions of sulphur 60, and 1 of carbon 11.4.

10. Phosphorus has been supposed capable of uniting to carbon; but in cases when specimens of phosphorus afford charcoal it is most probably mixed with the

* [A detailed account of these experiments may be found in the *Edinburgh Journal of Science*, vol. vi. New Series.]

substance, or in triple combination with oxygen and hydrogen; and no distinct action of the two bodies, and no definite compound of them has as yet been described.

11. A number of forms of carbon are found in nature; one of the most interesting of them is the *diamond*; the properties of this stone are well known, it is the hardest of the gems, and is usually crystallized, often in the form of an octohedron: its specific gravity is about 3.5; it does not conduct electricity. Of all known bodies it has the greatest power of refracting light. When the diamond is strongly heated in air, it consumes away; and if it be exposed to oxygen continuously ignited by a burning glass, or by other means, it acts upon the oxygen nearly in the same manner as charcoal. The volume of the oxygen is not perceptibly changed, and it is found converted into carbonic acid. M. Lavoisier first determined that carbonic acid was formed from diamond; and Messrs. Tennant, Allen, and Pepys, have demonstrated by some refined experiments that it produces about the same quantity as an equal weight of charcoal. Hence it has been concluded, that the diamond is pure carbon, differing from charcoal merely in the arrangement of its parts. When it is considered, however, that charcoal is a conductor and diamond a non-conductor of electricity, and that their physical properties differ entirely, it is impossible to receive this conclusion without doubt. I found that diamond powder heated strongly with potassium became blackened; and an effect was produced on the metal similar to that which the absorption of a minute quantity of oxygen would occasion; this would lead to the suspicion that there may be a little oxygen in diamond; but new experiments are wanting to prove this, and the

quantity, if any, must be very minute, which does not harmonise with the doctrine of definite proportions. If it should be ultimately found that the diamond is merely pure carbon, it will be an argument in favour of the varieties of elementary forms being produced by different aggregations or arrangements of particles of the same matter; for it is scarcely possible to fix upon bodies less analogous than lamp black, and the most perfect and beautiful of the gems.*

12. Plumbago or black lead, and anthracite or stone coal, are both tolerably pure forms of the carbonaceous element. In plumbago the carbon is united either chemically or mechanically to about $\frac{1}{25}$ of iron; in anthracite with small quantities of earthy matter. In the anthracite of Kilkenny in Ireland, the texture is often fibrous, and the substance has all the characters of well burned charcoal. In flaming coal the carbonaceous element is united to bitumen.

13. Few substances are more important in civilized life than the different forms of carbon; in their various uses they are essential to the comforts and well being of society, and are necessary in almost all the useful arts and manufactures.

The inflammable gases produced by the distillation of pit-coal have already been successfully used in manufactories for the purpose of affording light, and the application is at once safe and economical.

In nature the carbonaceous element is constantly active in an important series of operations; it is evolved in fermentation and combustion in carbonic acid; it is separated from oxygen in the organs of plants, is a principal element in animal structures, and is found in

* [His later researches confirmed this conclusion.]

different forms in almost all the products of organized beings.

VI. *Of Boron,* or the Boracic basis.*

1. There is a white crystalline substance found native in volcanic districts called boracic acid. It may be procured artificially from borax by heating it in oil of vitriol diluted with eight times its weight of water: it is difficultly soluble in water, and may be separated by a filter of cloth or paper. When this substance, slightly moistened, is exposed between two surfaces of platina, electrified by a Voltaic battery of not less than 100 double plates; a dark-coloured substance separates on the plate negatively electrified. This substance is *boron*, or the basis of the boracic acid. In this way it can be procured only in very minute quantities, and to obtain it for the purposes of experiment, boracic acid that has been long exposed to a red heat, is powdered and strongly ignited with an equal weight of potassium, in a tube of iron or copper. The result is exposed to diluted muriatic acid, and washed with it till nothing remains but a dark powder, which, when dried at a red heat, is the substance in question.

2. I first discovered boron in October, 1807, by the electrical decomposition of boracic acid, and by potas-

* In my first paper on this substance I named it boracium, for I supposed that in its pure form it would be found to be metallic; subsequent experiments have not justified this conjecture. It is more analogous to carbon than to any other substance; and I venture to propose Boron as a more unexceptionable name; the termination in *um* having been long used as characteristic of a metal. MM. Gay Lussac and Thenard have proposed to call it Bore, a word that cannot with propriety be adopted in our language, though short and appropriate in the French nomenclature.

sium, in March, 1808; but not in sufficient quantities to examine its properties, or to ascertain its nature. MM. Gay Lussac and Thenard, in June, 1808, made the experiment of heating boracic acid and potassium together, but they did not describe the properties of boron till the middle of November; and in the beginning of the same month I had procured sufficient quantities of the substances to ascertain its chemical relations. MM. Gay Lussac and Thenard, I believe, recomposed the boracic acid before me, and our experiments were independent of each other; but in my first paper on potassium and sodium read at the Royal Society, in November, 1807, at a time when the French chemists had no idea of the existence of the alkaline metals, I pointed out the probable application of these bodies to the decomposition of the acids not decomposed.

3. Boron is an opaque, dark olive coloured powder, infusible, and not volatile at any temperature to which it has yet been exposed. When heated strongly in contact with air, it burns, and forms dry boracic acid. In oxygen gas it throws off bright scintillations, becomes coated with boracic acid, and the portion not converted into acid, is found darker coloured than before. When gently heated in chlorine it emits white fumes, but has no energetic action on the gas. It is a non-conductor of electricity, and insoluble in water.

4. That boron combines with oxygen is shewn by the phenomena of its combustion. *Boracic acid* is the only well known result of their combination: the preparation of boron proves that the boracic acid consists of this body united to oxygen, for oxygen is added to the potassium in the process. It is very difficult to ascertain the proportions of boron and oxygen in boracic acid, for the boracic acid formed in combustion prevents

the process from going on; and the black substance, which is probably an oxide of boron, is burnt only with great difficulty. From comparing the quantity of potassium required to decompose a given quantity of boracic acid, with the quantity of oxygen absorbed in the production of the acid, I am inclined to believe that boracic acid cannot contain much less than $\frac{2}{3}$ of its weight of oxygen: I have made a number of experiments on this subject, but have never gained perfectly satisfactory results. MM. Gay Lussac and Thenard conceive that boracic acid contains only $\frac{1}{3}$ of its weight of oxygen; but their conclusions were drawn from the action of boron on solution of nitric acid, and the evaporation of the products; and boracic acid forms volatile compounds both with water and nitric acid; for I find that dry nitre and boracic acid afford by distillation a fluid containing a considerable quantity of boracic acid. From the quantity of ammonia required to neutralize boracic acid, it appears that the number representing it is about 160: and to destroy the alkaline properties of 90 parts of potassa requires twice 160 of boracic acid, so that its acid powers are extremely feeble.

Boracic acid in its common form is in combination with water; it then appears as a series of thin white hexagonal scales; its taste is very slightly acid, it reddens vegetable blues. By a long continued white heat the water is driven off from it, and a part of the acid sublimes, the remaining acid is a transparent fixed glass, which rapidly attracts moisture from the air. The compound of boracic acid and water appears to contain, from my experiments, about 57 parts of acid to 43 of water. The specific gravity of the hydrat of boracic acid, as it may be called, is 1.479, that of the dry acid 1.803.

Boracic acid is very little soluble in water, even when boiling that fluid does not take up $\frac{1}{50}$ of its weight. It dissolves in alcohol, and gives it the power of burning with a green flame.

5. Much still remains to be known respecting the nature and properties of boron, and its combinations. Probably a combination of it with chlorine may be formed.* It seems to exert no action on any of the inflammable bodies except sulphur, which dissolves a little of it by a long continued heat, and gains a green tint.

It has hitherto been obtained in quantities too small to ascertain whether it will have any application to the arts.

DIVISION V.

OF METALS; THEIR PRIMARY COMBINATIONS WITH OTHER UNDECOMPOUNDED BODIES, AND WITH EACH OTHER.

I. *General Observations.*

1. The metals form a numerous and very important class of natural bodies; they are connected with each other by close analogies, and by remote analogies to the inflammable solids described in the preceding pages: the number of metals known, or the existence of which may be presumed, amounts to 38.† The characteristic properties of the metals are a high degree of lustre, opacity, combustibility, and the power of conducting

* [A gaseous compound of boron and chlorine has been obtained by M. Berzelius.]

† [The number at present known is 42; cadmium, lithium, thorium, and vanadium have been discovered since 1812.]

electricity. A considerable degree of specific gravity was formerly considered as an essential character of metallic substances; but I have discovered bodies lighter even than water, which agree in all other essential qualities with metals, and which consequently must be arranged with them. In the order of classification to be adopted in the following pages, the most inflammable metals will be the first considered: though of recent discovery they are the most important as agents of analytical chemistry, and have offered the means of reducing other substances to the metallic form. The most inflammable metals produce alkalies, alkaline earths, and earths in combustion. Other metals afford the substances called oxides, which are analogous to earths; and a few are converted into acids. The metals that produce alkalies are potassium and sodium: the alkaline earths are formed from metals, which have been called barium, strontium, calcium, and magnesium. The metals supposed to be contained in common earths are silicium, alumium, zirconium, ittrium, and glucium. The metals that produce oxides are manganese, zinc, tin, iron, lead, antimony, bismuth, tellurium, cobalt, copper, nickel, palladium, uranium, osmium, tungsten, titanium, columbium, cerium, iridium, rhodium, mercury, silver, gold, and platina. The metals that produce acids are arsenic, molybdenum, and chromium.

2. The metals differ considerably in their mechanical properties, in degrees of hardness, ductility, and tenacity; all of them that are fusible by common means assume regular crystalline forms by slow cooling, and these forms are usually cubical or octohedral. The common metals in consequence of their fusibility, malleability, hardness, and durability, have been the most important instruments of the arts; the uses of them have

been essential to the progress of civilization; and most of the comforts, and many of the luxuries and refinements of social life are connected with their applications.

II. *Of Potassium.*

1. There is a body usually called potash, or the vegetable alkali, which may be thus procured: quick lime is mixed with solution of wood-ashes, and boiled for some time with it. The liquor so obtained, after being passed through bibulous paper, is evaporated till a solid matter remains; this solid matter is heated with alcohol or pure spirit; the spirit is separated by distillation in a vessel of silver; a fusible solid mass is produced, which is the substance in question.

To form *potassium*: this substance, in a thin piece, is placed between two discs of platina connected with the extremities of a Voltaic apparatus of 200 double plates; it will soon undergo fusion, oxygen will separate at the positive surface, and small metallic globules will appear at the negative surface, which consist of potassium. I discovered this metal in the beginning of October, 1807.

2. It may be procured by chemical means without electricity. If iron turnings be heated to whiteness in a curved gun-barrel, and potash be melted and made slowly to come in contact with the turnings, air being excluded,* potassium will be formed, and may be collected in a cool part of the tube; this method of procuring it was discovered by MM. Gay Lussac and Thenard, in 1808. It may likewise be produced by

* See Plate VIII. fig. 25.

igniting potash with charcoal, as M. Curaudau shewed in the same year.

3. Potassium is possessed of very extraordinary properties; it is lighter than water, its specific gravity is between 8 and 9, water being 10. It is a solid at common temperatures; it is very soft, and easily moulded by the fingers. It fuses at about 150° Fahrenheit, and rises in vapour, in a heat a little below that of redness. It is perfectly opaque. Its colour is white, like that of silver when it is newly cut, but it rapidly tarnishes in the air; and to be preserved from change must be kept under naphtha. It is a conductor of electricity. When thrown upon water it acts with great violence, swims upon the surface, and burns with a beautiful light, which is white mixed with red and violet; the water in which it burns is found alkaline, and contains a solution of potassa. It inflames when gently heated in the air, burns with a red light, and throws off fumes, which are alkaline. It burns spontaneously in chlorine with intense brilliancy.

It acts upon all fluid bodies containing water, or much oxygen or chlorine; and in its general powers of chemical combination may be compared to the alkahest, or universal solvent imagined by the alchemists.

4. Potassium combines with oxygen in different proportions. When potassium is gently heated in common air or in oxygen, the result of its combustion is an orange-coloured fusible substance; and for every grain of potassium consumed, about a cubical inch and $\frac{7}{10}$ of oxygen disappear. To make the experiment accurately the metal should be burnt in a tray of platina, covered with a coating of common salt, which may be easily done by fusing it in contact with the platina. This salt

is one of the few substances that has no action on the oxides of potassium.

The substance procured by the combustion of potassium at a low temperature, I had observed in October, 1807, but I supposed it to be the oxide of potassium containing the smallest quantity of oxygen, for it effervesced in water; MM. Gay Lussac and Thenard, in 1810, first demonstrated its real nature, and shewed that it was the combination of oxygen and potassium containing the largest quantity of oxygen.

The gas produced by its effervescence with water is oxygen. When it is fused and brought in contact with combustible bodies, they burn with vividness. When it is heated in carbonic acid, oxygen gas is expelled, and it is converted into the compound called subcarbonate of potash.

When it is heated very strongly upon platina, oxygen gas is expelled from it, and there remains a difficultly fusible substance of a gray colour, vitreous in its fracture, and which dissolves in water without effervescence, but with much heat, and renders the water alkaline. This substance is pure potash or *potassa*, which was unknown in its uncombined state till I discovered potassium, but which has long been familiar to chemists combined with water in the substance which has been called pure potash; but which ought to be called the *hydrat of potassa*.

That the potash obtained by alcohol in the manner described in the beginning of this section, is a compound of potassa and water, is shewn by many experiments. If it be made to act upon iron turnings at a dull red heat, the iron becomes combined with oxygen, hydrogen is given off in abundance, and the alkali loses its easy fusibility, becomes harder, more opaque, and of

greater specific gravity. In producing potassium in the experiment of acting upon white hot iron turnings by common potash, hydrogen is disengaged in abundance from the decomposition of the water in the potash ; and I have procured this water by heating together the potash prepared by alcohol, and boracic acid ; 100 parts of potash treated in this way gave between 17 and 18 parts of pure water.

Potassa entirely free from water may be procured by other means besides the decomposition of the orange oxide of potassium, or the action of iron on common potash ; for instance, by acting on potassium by a small quantity of water, or by heating potassium with common potash. The proportion of oxygen in potassa is learned by the action of potassium upon water : 8 grains of potassium produce from water about 9 cubical inches and a half of hydrogen ; and for these there must be added to the metal 4 cubical inches and three quarters of oxygen.

5. It has been mentioned, page 82, that the number representing potassium is 75 ; and it appears from the experiments, that the orange oxide of potassium must consist of 1 proportion of potassium 75, and 3 of oxygen 45 ; and the number representing it is 120. Potassa must consist of 1 proportion of potassium 75, and 1 of oxygen 15, and the number representing it is 90. Hydrat of potassa, or the potash prepared by alcohol, must contain 1 proportion of potassa 90, and 1 of water 17.*

* In the few experiments that I have made on hydrat of potassa, there has been rather more water indicated, between 17 and 19 per cent. ; but the potash I used was, I doubt not, adulterated with a little soda, as no particular care was taken to purify it, and hydrat of soda contains more water in proportion ; and there is great reason to believe that 90 and 17 are the true estimation. MM. Gay Lussac and Thenard allow about $\frac{1}{3}$ of water in potash.

6. When potassium is heated strongly in a small quantity of common air, the oxygen of which is not sufficient for its conversion into potassa, a substance is formed of a grayish colour, which, when thrown into water, effervesces without inflaming. This substance is likewise generated in experiments on the production of potassium by iron and hydrate of potassa, when a little common air is admitted into the barrel. It is doubtful whether it be a mixture of potassa and potassium, or a combination of potassium with a smaller quantity of oxygen than exists in potassa, that is, a protoxide of potassium. If a protoxide of potassium, it probably contains 2 proportions of potassium and 1 of oxygen.

7. I have already referred to the action of potassium and chlorine; the inflammation produced when thin pieces of potassium are introduced into chlorine is very vivid: potassium separates chlorine from hydrogen and phosphorus with inflammation; and when potassium is made to act upon sulphurane [chloride of sulphur] there is a violent explosion. The attraction of chlorine for potassium is much stronger than the attraction of oxygen; potassa, and the orange oxide of potassium, are immediately decomposed by chlorine, the chlorine combines with the metal, and the oxygen is set free.

The combination of chlorine and potassium is the substance which has been improperly called muriate of potash, and which, in common cases, is formed by causing muriatic acid and solution of potassa to act upon each other, and by heating the mixture to redness; in which case the hydrogen of the acid, and the oxygen of the alkali are set free as water; and the metal of the alkali and the chlorine of the acid combine. From various analytical experiments it appears that muriate of potash, [chloride of potassium,] which may be called

potassane, consists of 75 of potassium and 67 of chlorine, and the number representing it is 142. Potassane is the only known combination of potassium and chlorine.

8. There appears to be a gaseous combination of potassium and hydrogen; for I found that when potassium is heated strongly in hydrogen the gas contracts in volume, and becomes spontaneously inflammable, and gives alkaline fumes in its combustion. MM. Gay Lussac and Thenard state that there is a solid compound of hydrogen and potassium, which may be obtained by heating the metal for a long while in the gas, at a temperature just below that of ignition. They describe it as a grayish solid, and state that it gives off its hydrogen by the action of mercury. As yet no experiments have been made on the proportions in which hydrogen and potassium combine.

9. Potassium and sulphur combine with great energy when they are heated together, producing much light and heat, even when the experiment is made out of the contact of air. The *sulphuret of potassium* is of a dark gray colour; acts with great energy upon water, producing sulphuretted hydrogen, and burns brilliantly when heated in the air, becoming the salt called sulphate of potash. From my experiments there is every reason to believe that this compound consists of one proportion of sulphur 30, and one of potassium 75, and the number representing it will be 105. Potassium has so strong an attraction for sulphur, that it rapidly separates it from hydrogen; and potassium heated in sulphuretted hydrogen takes fire, and burns with great brilliancy, and sulphuret of potassium is formed, and hydrogen set free.

10. Potassium and phosphorus enter into union producing light; but they act on each other with less

energy than potassium and sulphur. The *phosphoret of potassium* in its common form is a substance of a dark chocolate colour; but when heated with potassium in great excess it becomes of a deep gray colour, and of considerable lustre, so that it is likely that phosphorus and potassium are capable of combining in two proportions; probably the chocolate-coloured substance contains one proportion of each, and the dark gray substance two proportions of the metal.

The phosphoret of potassium burns with great brilliancy when exposed to air, and when thrown into water produces an explosion in consequence of the immediate disengagement of phosphoretted hydrogen.

11. When charcoal is present during the production of potassium, it usually contains a small quantity of carbonaceous matter; and charcoal that has been heated strongly in contact with potassium effervesces in water, and renders it alkaline, though previously exposed to a temperature at which potassium rises in vapour. These circumstances shew that there is an attraction, though feeble, between potassium and carbon; but as yet no compound of the two bodies of which the proportions can be assigned has been obtained.

12. Potassium, like all other metals, has resisted all attempts to resolve it into other forms of matter. Since I first discovered it, and announced it as an undecomposed substance, there has been much discussion respecting its nature. MM. Gay Lussac, Thenard, Ritter, and Dalton, supposed that it was a compound of hydrogen and potassa; but the first two chemists have allowed that the phenomena are incompatible with such an hypothesis; in this case potassium should form hydrate of potassa, or a substance containing water in combustion, which is not the case; nor has hydrogen

been in any instance obtained in experiments on potassium, except when substances known to contain hydrogen were present; and it would not be more absurd to say that phosphorus is a compound of hydrogen and phosphoric acid, than to say that potassium is a compound of hydrogen and potassa.

13. Potassium, of all known substances, is that which has the strongest attraction for oxygen; and it produces such a condensation of it that the oxides of potassium are heavier than the metal itself. Potassium may be used as a general agent for detecting the presence of oxygen in bodies; and a number of substances undecomposable by other chemical agents are readily decomposed by this substance.

The compounds of potassium are of great use in the arts; potassa enters into the composition of soft soap, and the salts having a basis of potassa are many of them used in medicine.

III. *Sodium.*

1. *Sodium* may be procured exactly in the same manner as potassium, by electrical or chemical decomposition, the mineral alkali, or the alkali from the ashes of marine plants being used instead of pearl ashes. Rather a higher degree of heat is necessary for its production by the action of iron.

I discovered sodium a few days after I discovered potassium, in the year 1807.

2. In many of its characters it resembles potassium; it is as white as silver, has great lustre, and is a conductor of electricity. It enters into fusion at about 200° Fahrenheit, and rises in vapour at a strong red heat. Its specific gravity is between 9 and 10. When

heated strongly in oxygen or chlorine, it burns with great brilliancy. When thrown upon water it effervesces violently, but does not inflame,* swims on the surface, gradually diminishes with great agitation, and renders the water a solution of soda. It acts upon most substances in a manner similar to potassium, but with less energy. It tarnishes in the air, but more slowly, and like potassium it is best preserved under naphtha.

3. Sodium forms two distinct definite combinations with oxygen: one is pure soda, which has long been known, combined with water, in the substance which has been called by chemists *Soda*, but which was not examined in its uncombined state, till I formed it from the metal; the other is the orange oxide of sodium, which I observed in 1807; but of which the true nature was pointed out in 1810 by MM. Gay Lussac and Thenard.

Pure soda may be made by burning sodium in a quantity of air containing no more oxygen than is sufficient for its conversion into the alkali, i. e. the metal must be in excess: a strong degree of heat must be applied.

Pure soda is of a gray colour, it is a non-conductor of electricity, of a vitreous fracture, requiring a strong red heat for its fusion. When a little water is added to it, there is a violent action between the two bodies; the soda becomes white, crystalline in its appearance, and much more fusible and volatile, and is then the substance which has been long known under the name of soda, but which may, with more propriety, be called *hydrate of soda*.

The *oxide of sodium* may be formed by burning sodium in oxygen gas in excess. It is of a deep orange colour, very fusible, and a non-conductor of electricity; when acted upon by water, it gives off oxygen gas, and

[* If thrown on hot water inflammation occurs.]

the water becomes a solution of soda; it deflagrates when strongly heated with combustible bodies.

The proportions of oxygen in soda, and the orange oxide or peroxide of sodium, are easily learned by the action of sodium on water, and on oxygen. If a given weight of sodium, in a little glass tube, be thrown by means of the finger, under a graduated inverted jar filled with water, the quantity of hydrogen evolved will indicate the quantity of oxygen combined with the metal, to form soda; and when sodium is burnt slowly in a tray of platina, lined with dry common salt in oxygen in great excess; from the quantity of oxygen absorbed, the composition of the peroxide may be learned. From experiments that I have made on this subject, compared with those made by MM. Gay Lussac and Thenard, it appears that the number representing sodium is 88, and that soda consists of one proportion of sodium, and two of oxygen, 88 and 30: the oxide of sodium, of one proportion of sodium, and 3 of oxygen, 88 and 45: and hydrate of soda (soda prepared by alcohol) contains one proportion of sodium, two of oxygen, and two of water, and the number representing it is 152.

When sodium is kept for some time in a small quantity of moist air, or when sodium in excess is heated with hydrate of soda, a dark grayish substance is formed, more inflammable than sodium, and which affords hydrogen by its action upon water. It is probable that this is sodium in its first degree of oxygenation, or the *protoxide of sodium*; but as yet no experiments have been made on its constitution. If the protoxide, it is likely that it consists of one proportion of sodium, and one of oxygen.

4. Only one combination of sodium and chlorine is

known: it is the important substance *common salt*. It may be formed directly by combustion, or by decomposing any compound of chlorine by sodium. Its properties are well known; it is a non-conductor of electricity, is fusible at a strong red heat, is volatile at a white heat, and crystallizes in cubes. Sodium has a much stronger attraction for chlorine than oxygen; and soda or hydrate of soda is decomposed by chlorine, oxygen being expelled from the first, and oxygen and water from the second.

Potassium has a stronger attraction for chlorine than sodium has; and one mode of procuring sodium easily, is by heating together to redness common salt and potassium. The compound of sodium and chlorine* has been called muriate of soda, in the French nomenclature; for it was falsely supposed to be composed of muriatic acid and soda; and it is a curious circumstance, that the progress of discovery should have shewn that it is a less compounded body than hydrate of soda, which six years ago was considered as a simple substance, and one of its elements. According to the nomenclature which I have ventured to propose, the chemical name for common salt will be *sodane*.

Common salt consists of one proportion of sodium, 88, and two of chlorine, 134; and the number representing it is 222: when the proper corrections are made, the most accurate analyses, particularly those of Dr. Marcet, are found to agree with this number.

5. There is no known action between sodium and hydrogen, or azote.

6. Sodium combines readily with sulphur, and with phosphorus, presenting similar phenomena to those presented by potassium. The sulphurets and phosphorets of sodium agree in their general properties with those

* [Chloride of sodium.]

of potassium, except that they are rather less inflammable. They form, by burning, compounds of sulphuric and phosphoric acid and soda, and therefore must contain two proportions of the inflammable substances, to one of sodium.

7. Sodium, when made from substances containing charcoal, usually affords charcoal by combustion; but as yet no definite combination of the two bodies has been obtained. No experiments have been made on the action of sodium on boron.

8. Potassium and sodium combine with great facility, and form peculiar compounds, which differ in their properties according to the proportions of their ingredients. By a small quantity of sodium, potassium is rendered fluid at common temperatures, and its specific gravity considerably diminished. Eight parts of potassium, and one of sodium, form a compound that swims in naphtha, and that is fluid at the common temperature of the air. Three parts of sodium, and one of potassium, make a compound, fluid at common temperatures. A little potassium destroys the ductility of sodium, and renders it very brittle, and very soft.

9. The compounds of sodium are of great importance in the common arts, and are subservient to many of the wants of life. Soda is the most important ingredient in the different species of glass, and in hard soaps. The glasses are composed of soda united to earths and oxides; the soaps consist of soda, united to oily substances. Common salt is found abundantly in nature; it exists in small quantities in almost all waters and all soils. It diminishes the tendency of animal or vegetable substances to decompose, and probably preserves the ocean in a state fitted for the purposes of animal life. It is a part of the nourishment of animals, and

though taken in very small quantities, seems to perform an important part in their economy.

9. The compounds formed by potassium and sodium, like the metals themselves, are possessed of strong resemblances; they may however be chemically distinguished by a very simple test; the diluted aqueous solutions of the compounds of potassium, render cloudy the nitro-muriatic solution of platina, which is not the case with similar solutions of the compounds of sodium. Most of the compounds of sodium differ from those of potassium, in containing double proportions of the other elements. Potassa contains one proportion of oxygen only; soda contains two, and the salts having potassa for their basis, contain only one proportion of acid, whilst those having soda for their basis, contain two. The attractions of potassium for all substances that have been examined, are stronger than those of sodium; and when sodium is procured from compounds, by the agency of potassium, 150 parts in weight of potassium, or two proportions, are required to produce 88 parts of sodium, or one proportion.

IV. *Barium.*

1. There is a mineral substance found in Cumberland, Yorkshire, and other parts of Britain, called Witherite, or carbonate of baryta. By dissolving this substance in dilute solution of nitric acid, evaporating the solution to dryness, and heating the salt obtained to whiteness, a light fawn-coloured powder is procured, which is baryta or barium combined with oxygen. To obtain *barium*, a quantity of this substance is made into a paste, with water, and placed on a plate of platina; a cavity is made in the paste to receive a globule of mercury; the mercury is rendered negative, the platina

positive, by means of a Voltaic battery, containing about 100 double plates.

In a short time an amalgam will be formed, consisting of mercury and barium. This amalgam must be introduced into a little tube made of glass free from lead, which must be bent in the form of a retort, filled with the vapour of naphtha, and hermetically sealed. Heat must be applied to the end of the tube containing the amalgam, till all the mercury has been driven off; there will remain a solid difficultly fusible metal, which is barium.

2. I first gained indications of the decomposition of baryta, in the end of October 1807, and I obtained an alloy of it with iron, in March, 1808. The process of electrifying mercury, in contact with the earth, was pointed out to me in the course of my inquiries, by MM. Berzelius and Pontin of Stockholm, in May 1808; and in the beginning of June in the same year, I obtained *the metal*.

3. Barium, as procured by heating the amalgam, appeared of a dark gray colour, with a lustre inferior to that of cast iron. It was considerably heavier than sulphuric acid, for though surrounded by globules of gas, it sunk immediately in that fluid. It instantly became covered with a crust of baryta, when exposed to air, and burnt with a deep red light, when gently heated. When thrown into water it effervesced violently, disappeared, and the water was found to be a solution of baryta.

Barium as yet has been obtained only in very minute quantities. I have never possessed enough of it to ascertain its general chemical and physical characters, and no experiments upon it have been published by any other person.

4. From some results that I have obtained, it seems

probable that barium may be procured by chemical, as well as electrical decomposition. When baryta, or the salt improperly called muriate of baryta, ignited to whiteness, was exposed to the agency of potassium, that metal being sent through it in vapour, a dark gray substance appeared, diffused through the baryta, or the muriate, not volatile, that effervesces copiously in water, and that lost its metallic appearance by exposure to air:—the potassium in this process was converted into potassa.

5. The only well-known combination of barium with oxygen is *baryta*, or *baria*. It is of a pale grayish green colour. Its specific gravity is about 4, that of water being 1. This substance is a non-conductor of electricity, has a strong caustic taste, reddens turmeric, and renders green vegetable blues. When acted upon by a small quantity of water, it heats violently, becomes white, unites to a proportion of water, and becomes a hydrate. The pure alkaline earth is infusible, except by an intense heat: the hydrate fuses at a strong red heat; a considerable part of its water is expelled by a still higher temperature. Baryta is soluble at 60° , in about 20 parts of water, and at 212° in about 2 parts. That baryta is composed of barium and oxygen, is proved by the combustion of barium in oxygen; in which, as I have found, oxygen is absorbed, and no product but baryta formed. It is likewise proved synthetically by the action of barium upon water, in which case hydrogen is evolved; and analytically it appears from the action of potassium, on the earth. From indirect experiments, I am inclined to consider baryta as composed of 89.7 of barium, and 10.3 of oxygen: and supposing the earth to consist of one proportion of metal and one of oxygen, the number representing

barium will be 130, and that representing the alkaline earth will be 145.

Barium, as would appear from the experiments of MM. Gay Lussac and Thenard, is capable of combining with more oxygen than exists in baryta. These able chemists state, that when baryta is gently heated by a spirit lamp, in a glass tube filled with oxygen gas, an absorption of the gas takes place. As yet no experiments have been made on the properties of this *oxide of barium*, or on the quantity of oxygen it contains; probably baryta may be easily combined with oxygen, by heating it with hyper-oxy muriate of baryta. *The hydrate of baryta*, if its composition be estimated from M. Berthollet's experiments, consists of one proportion of baryta, and one of water.

6. One combination only of barium and chlorine is known; it may be formed by heating baryta in muriatic acid gas, or in chlorine. In the first case, the oxygen of the baryta produces water by combining with the hydrogen of the acid; in the second it is expelled: and in an experiment made on purpose, I found that for every part in volume of chlorine absorbed, half a part of oxygen was given off from the alkaline earth. Hence it may be concluded that the compound of barium and chlorine contains one proportion of metal 130, and one of chlorine 67. This substance is fusible by a very strong heat, is very soluble in water; its taste is bitter, its colour white, it is crystalline and transparent. It is improperly called in the French nomenclature, muriate of baryta. According to the principles of nomenclature which I have proposed, its name will be *barane*.*

8. No other combinations of barium, except those with oxygen and chlorine, have been as yet examined;

* [Chloride of barium.]

there can, however, be little doubt that its powers of combination will be, in many respects, analogous to those of potassium and sodium, as, of all metallic substances, it is the nearest related to these bodies.

9. The compounds of barium have as yet been applied to the arts in very few cases. Baryta is employed in small quantities, in the manufacture of certain kinds of porcelain; most of the salts containing baryta as a basis, are poisonous. The combination of baryta and carbonic acid, made artificially by pouring a solution of carbonate of ammonia, into a solution of nitrate of baryta, forms a pigment of a very pure white colour.

V. *Strontium*.

1. *Strontium* may be procured precisely in the same manner as barium; carbonate of strontia, or strontianite, a mineral found at Strontian in Scotland, being used instead of witherite. I first procured this metal in 1808, but in quantities too small to make an accurate examination of its properties. It seemed very analogous to barium, had not a very high lustre, appeared fixed, difficultly fusible, and not volatile. It became converted into strontia by exposure to air, and when thrown into water, decomposed it with great violence, producing hydrogen gas, and making the water a solution of strontia.

2. One combination of strontium with oxygen only is at present known; it is *strontia*, or strontites, the substance procured by burning strontium.* It may be produced in large quantities by igniting strontianite intensely with charcoal powder, or by heating to white-

* [A peroxide may be obtained in the same manner as the peroxide of barium.]

ness the salt formed from this fossil, by the action of nitric acid. It appears of a light fawn colour, and agrees in many of its characters with baryta. It is fusible only by an intense heat. Its specific gravity is between three and four, water being one. It is soluble in about 200 parts of water, at common temperatures, and is much more soluble in hot than cold water; its taste is acrid and alkaline, it reddens paper tinged with turmeric. When acted upon by a small quantity of water it becomes hot, its colour changes to white, and it is converted into a hydrate, and then becomes fusible at a white heat. From indirect experiments, I am disposed to regard it as composed of about 86 of strontium and 14 oxygen; and supposing it to contain one proportion of metal and one of oxygen, the number representing strontium will be 90, and that representing the earth 105.

3. No experiments have as yet been made on the direct combination of strontium and chlorine; but a substance which appears to consist of these two bodies, and no other elements, may be made, by heating strontia strongly in chlorine, or muriatic acid gas, or by igniting to whiteness the salt formed by the solution of strontianite in muriatic acid. By the action of chlorine on strontia, oxygen is expelled: by the action of muriatic acid gas upon it, water is formed. The compound of chlorine and strontium, or *strontane*,* is a white substance, difficultly fusible, fixed in the fire, a non-conductor of electricity, and of a peculiar bitter taste; when brought in contact with the flame of wax, tallow, oil, or alcohol, it tinges it of a rose colour; and this is a distinctive character of the compounds of strontium; the salts formed from it give this tint to

* [Chloride of strontium.]

flame, those of baryta give a yellow tint. From direct experiments I ascertained that 50 parts of strontane consisted of about 29 parts of metal and 21 of chlorine ; so that it must be regarded as composed of one proportion of strontium, and one of chlorine, 90 and 67.

4. No experiments have as yet been made on the action of strontium, on any of the other elementary substances.

5. None of the compounds of this body have as yet been applied to any of the purposes of the arts, and its combinations are rare in nature.

VI. *Calcium.*

1. *Calcium* may be obtained by the same processes as barium and strontium. Mild calcareous earth, or chalk being used instead of witherite and strontianite ; or common well-burnt lime may be employed for making the paste, from which the mercurial amalgam is to be formed by Voltaic electricity.

I first procured calcium about the same time as barium and strontium, but only in very minute quantities, so that little can be said concerning its nature. It appeared brighter and whiter than these two metals, and burnt, when gently heated, producing dry lime. I have had no opportunity of examining its general physical and chemical qualities.

2. There is only one known combination of calcium and oxygen, which is the important substance, *lime* or *calcia*. The nature of this substance is proved by the phenomena of the combustion of calcium ; the metal becomes converted into the earth, with the absorption of oxygen gas. When the amalgam of calcium is thrown into water, hydrogen gas is disengaged, and the

water becomes a solution of lime; and from the quantity of hydrogen gas disengaged, compared with the quantity of lime formed in experiments of this kind, M. Berzelius has endeavoured to ascertain the proportion of oxygen in lime. The nature of lime may be also proved by analysis; when potassium in vapour is sent through the earth, ignited to whiteness, the potassium, I have found, becomes potassa, and a dark gray substance of metallic splendour, which is calcium either wholly or partly deprived of oxygen, is found embedded in the potassa, and it effervesces violently, and forms a solution of lime, by the action of water.

Lime is obtained for common purposes, from marble of the whitest kind, such as the Parian or Carara marble, by long exposure to a strong heat. It is a white soft substance, of specific gravity 2·3. It requires an intense degree of heat for its fusion, and has not yet been rendered volatile. Its taste is analogous to, but milder than that of baryta and strontia. It is soluble in about 450 parts of water, and seems to be more soluble in cold than in hot water. It acts upon vegetable colours in a manner similar to the other alkaline earths. When water, in small quantities, is added to it, a considerable heat is produced, a portion of the water combines with the lime, and it becomes a hydrate; but water does not adhere to it with the same degree of energy, as to baria and strontia, for it may be expelled by a strong red heat. From the experiments of M. Berzelius, and those which I have made, it appears that lime consists of about 20 of metal to 7·5 of oxygen, and the number representing calcium is 40, and that representing lime 55; and the *hydrate of lime* must consist of 55 lime and 17 water, which estimation agrees with the experiments of M. Lavoisier and Mr. Dalton.

I have attempted to combine lime with more oxygen, but without success.

3. When lime is heated strongly in contact with chlorine, oxygen is expelled, and chlorine absorbed; and, as happens in all the decompositions of metallic oxides, of which the metals combine with only one proportion of oxygen and chlorine, for every two in volume of chlorine absorbed, a volume of oxygen is expelled. The substance formed by the action of chlorine on lime, as the oxygen of the lime is expelled, must evidently consist of chlorine and calcium. It has been called dry muriate of lime; according to the true view of its composition, it may be called *calcane*.* It is a semi-transparent crystalline substance, fusible at a strong red heat, a non-conductor of electricity, has a very bitter taste, rapidly absorbs water from the atmosphere; and is extremely soluble in water: by the evaporation of its solution at a low heat, crystals may be obtained, which consist of calcane, combined with more than a third their weight of water. From my experiments, it appears that calcane consists of 31 chlorine and 19 of calcium, and hence it may be supposed to contain one proportion of the metal, and one of the gas, and the number representing it on this idea is 107; and it is evident, from the experiment on the action of chlorine on lime, that the proportion of oxygen in lime, and of chlorine in calcane, must be in the ratio of 15 to 67.

4. As yet no experiments have been made on the combinations of calcium with any of the inflammable, or acidiferous substances, or metals.

5. The compounds of calcium are found abundantly on the surface of the globe, and are of great importance

* [Chloride of calcium.]

in the economy of nature, and in the processes of art. Lime combined with carbonic acid is an essential part of fertile lands: a number of rocks are constituted by this substance. Gypsum or alabaster, is lime combined with sulphuric acid; and the earth of bones consists of lime united to phosphoric acid. There is no animal or vegetable substance that does not contain larger or smaller quantities of calcareous matter. The uses of lime in mortar are well known. Quicklime, employed as a manure, tends to decompose* and dissolve inert vegetable matter, and renders it proper for the nourishment of plants; and in this operation the lime is united to carbonic acid, and becomes a permanent part of the soil. In the process of tanning, lime is employed to remove the hair from the skins of animals, and it is used in certain operations of bleaching, dyeing, and other useful arts.

VII. *Magnesium.*

1. *Magnesium*† may be procured from the earth called magnesia, which is the same as the calcined magnesia of druggists, by processes similar to those referred to in the three preceding sections; but a much longer time is required to produce an amalgam of magnesium and quicksilver, by electrical powers, than to produce amalgams of the metals of the other alkaline earths.

* [According to the results of my experiments on the action of lime on animal and vegetable substances, it does not decompose, but preserves the majority of them.]

† In my first paper on the decomposition of the earths, published in 1808, I called the metal from magnesia, magnium, fearing lest, if called magnesium, it should be confounded with the name formerly applied to manganese. The candid criticisms of some philosophical friends have induced me to apply the termination in the usual manner.

I succeeded in decomposing magnesia likewise, in the following manner: I passed potassium in vapour through magnesia, heated to intense whiteness, in a tube of platinum, out of the contact of air; I then introduced a small quantity of mercury, and heated it gently for some time in the tube. An amalgam was obtained, which by distillation, out of the contact of the atmosphere, afforded a dark gray metallic film, which was infusible at the point at which plate glass softened, and which, in the process of distillation of the mercury, rendered the glass black at its point of contact with it. This film burnt when heated strongly, with a red light, and became converted into a white powder, which had the character of magnesia: when a portion of the metal was thrown into water, it sunk to the bottom, and effervesced slowly, becoming covered with a white powder; by adding a little muriatic acid to the water, the effervescence was violent; the metal rapidly disappeared, and the solution was found to contain magnesia.

I have made several experiments with the hope of obtaining larger quantities of magnesium, such as might have enabled me to examine its chemical and physical properties; but without success.* It is very difficult to procure a pure amalgam of magnesium by potassium and mercury; the heat must be intense; and at a high temperature, potassium acts with great energy upon platina, so that unless the tube is very solid, it is destroyed in the process, and when the heat is not very great, potassium remains in the tube, which is found afterwards in the amalgam. The potassium may, however, be separated by the action of water; which, even

* [M. Bussy has been successful, by heating together potassium and chloride of magnesium. This metal is not unlike silver in appearance; is malleable, and fusible at a red heat.]

in the amalgam, rapidly converts it into potassa, but which has a much feebler action on magnesium. When the amalgam contains potassium, it likewise usually contains platinum, which is very soluble in the compound of potassium and quicksilver.

2. There is only one known compound of magnesium and oxygen, which is the substance from which the metal is procured, *Magnesia*. That magnesia consists of magnesium and oxygen, is proved both by analysis and synthesis. In the production of magnesium by potassium, the potassium is found converted into potassa, and therefore must have gained oxygen from the magnesia; and in the formation of magnesia from magnesium, oxygen is absorbed. No experiments have as yet been made to determine the proportions of the elements in magnesia; but from experiments which I have made on the combinations of this substance with acids, assuming that they are single proportions, I am inclined to adopt 43 as the number representing it; and if it be supposed to be constituted by 1 proportion of metal, and 1 of oxygen, the number representing the metal will be 28.

Magnesia appears in its common form, as a white soft powder; its specific gravity is between 2 and 3. It is found in nature in the crystalline form; specimens have been brought from North America, which nearly resemble talc in their external characters. Magnesia has scarcely any taste, no smell; it reddens turmeric. It is infusible, except by the intense heat produced by the combustion of hydrogen gas in oxygen, or that generated by Voltaic electricity. It is scarcely soluble in water, but produces heat when water is mixed with it, and it absorbs a considerable portion of the fluid. When it is procured by the decomposition of a solution in which it

is combined with an acid, by means of solution of potassa or soda, it falls down in union with water, as a hydrate; but the water adheres to it with a very feeble attraction only, and is expelled entirely at a red heat. *Hydrate of magnesia* when dried at 212° , appears in coherent semitransparent masses, very brittle and soft; it contains about $\frac{1}{4}$ of its weight of water.

3. When magnesia is strongly heated in contact with chlorine, chlorine is absorbed, and oxygen expelled, and in the usual proportions as to volume. Hence it is evident that there exists a combination of magnesium and chlorine; though this body, which may be called *magnesane*,* has never been examined in a separate state. The salt called muriate of magnesia, is a compound of magnesane and water, and when it is acted on by a strong red heat, by far the greatest part of the chlorine unites to the hydrogen of the water, and rises in the form of muriatic acid gas, and the oxygen of the decomposed water combines with the magnesium to form magnesia; some magnesane is, however, found mixed with the magnesia, which affords crystals of muriate of magnesia by the action of water.

4. No experiments have as yet been made on the action of magnesium upon any of the inflammable or metallic substances.

5. The compounds of magnesium occur extensively diffused in nature. Magnesia exists in certain limestones which are found in different parts of Great Britain and Ireland, and which are less fitted for the general purposes of manure than common limestone. Magnesia, in its uncombined state, as appears from the experiments of Mr. Tennant, is injurious to plants, but united to carbonic acid, it seems to form an useful part

* [Chloride of magnesium.]

of the soil: the magnesian limestones are distinguished by their slow solution in acids; and they render weak solutions of nitric acid turbid by their action upon them. Magnesia, and some of its saline combinations are used in medicine; its application in bleaching has been referred to in an early part of this work.

VIII. *Aluminum.*

1. When a solution of ammonia or of potassa, not in excess, is thrown into a solution of alum, a substance falls down, which when well washed, and dried at a red heat, is alumina. This substance appears to contain a peculiar metal, but as yet Aluminum has not been obtained in a perfectly free state, though alloys of it with other metalline substances have been procured sufficiently distinct to indicate the probable nature of alumina.* Alumina cannot be decomposed by the electrization of mercury in contact with it, in the same manner as the alkaline earths. The first experiment by which I obtained evidences of its composition, was made in 1808, by fusing iron negatively electrified in contact with it; the earth was moist in this process, and a very high Voltaic power was applied. The globule of metal obtained was whiter than pure iron; effervesced slowly in water, becoming covered with a white powder, and the solution in muriatic acid decomposed by an alkali, afforded alumina and oxide of iron.

By passing potassium in vapour, through alumina heated to whiteness, the greatest part of the potassium became converted into potassa, which formed a cohe-

* [Pure aluminum has been obtained by Wöhler, by the action of potassium on the chloride. It requires a very high temperature for its fusion; it does not decompose cold water.]

rent mass with that part of the alumina not decomposed, and in this mass there were numerous gray particles, having the metallic lustre, and which became white when heated in the air, and which slowly effervesced in water. In a case in which a similar experiment was made, a strong red heat only being applied to the alumina, a mass was obtained, which took fire spontaneously by exposure to air, and which effervesced violently in water, and which probably contained the basis of alumina united to potassium.

2. That oxygen exists in alumina, cannot be doubted, when the conversion of potassium into potassa by its action upon it, is considered; and that it contains an inflammable substance united to oxygen, seems likewise evident; and that this is metalline in its nature appears extremely likely, both from the facts detailed, and from analogy; but this point cannot as yet be considered as demonstrated.

Alumina, in the form in which it is usually obtained, has no taste nor smell, adheres strongly to the tongue, has no action upon vegetable colours, is insoluble in water, is soluble in all the mineral acids and in hot solutions of fixed alkalies. When the precipitate from solutions of alum has been dried only at the temperature of the atmosphere, it is found combined with nearly an equal weight of water, and then appears as a white powder or a gelatinous substance. There is a native *hydrate of alumina* found in different parts of the world, crystallized and transparent, and which has been called Wavellite: from Mr. Gregor's experiments, and my own, it appears that this substance contains about 28 per cent. of water.*

No direct researches have been made on the quantity

* [Vide Vol. II. p. 303.]

of oxygen in alumina; but from some experiments that I have made on the quantity of ammonia required to decompose saturated solutions of alumina in acids, it would appear that the number representing alumina is about 48, and supposing it to consist of one proportion of aluminum, and one of oxygen, 33 will be the number representing aluminum.

3. No substance is known that can be regarded as a compound of chlorine and aluminum.* Alumina is soluble in solution of muriatic acid; but by heating the salt obtained, muriatic acid gas rises and alumina remains behind.

4. The compounds of alumina are found abundantly in the mineral kingdom, and many of them are of great importance in the common arts. Alumina forms a part of the greater number of rocks, and is found in larger or smaller quantities, in almost all soils. In its crystallized form coloured by small quantities of iron, it constitutes a beautiful class of gems, distinguished by the name *Telesia*, including the ruby, the sapphire, the oriental topaz, and other hard and brilliant stones.

Alumina combined with silica and other substances, forms the varieties of porcelain and china-ware. Its acid combinations are used to a great extent in dyeing and calico printing for fixing colours on stuffs.

IX. *Glucinum.*

1. There is an earth which was discovered by Vauquelin in 1798, called glucine, or glucina. It may be obtained from the beryl or the emerald, by the following process; the stone, in fine powder, must be ignited

* [Wöhler has obtained a chloride of aluminum by the action of chlorine on alumina under peculiar circumstances.]

for half an hour in a crucible of silver or platina, with three times its weight of hydrate of potassa or soda. The mass must be dissolved in solution of muriatic acid, and the compound obtained exposed to heat till it is dry. Water is then added to it, and the aqueous solution obtained acted on by solution of carbonate of potassa; a white powder is obtained which must be dissolved in diluted oil of vitriol not added in excess; a little of the salt called vitriolated tartar, or sulphate of potassa, must be mixed with the solution, and the whole evaporated till it begins to crystallise; crystals of alum will form. When no more can be obtained, the remaining liquor must be mixed with solution of carbonate of ammonia added in great excess; the mixed liquor must be passed through bibulous paper, evaporated to dryness, and the solid matter remaining heated to redness, it is then *glucina*.

2. There is great reason to believe that glucina is a compound of a peculiar metallic substance, which may be called *glucinum* and oxygen. The evidence that such is its composition, I have obtained, by heating it with potassium in the same manner as alumina; the potassium was for the most part converted into potassa, and dark coloured particles having a metallic appearance were found diffused through the mass, which regained the earthy character by being heated in the air, and by the action of water, and in this last case hydrogen was slowly disengaged.*

3. Glucina in its pure form appears as a white powder without taste or smell; it requires an intense degree of heat for its fusion; it is not soluble in water in any perceptible degree; it does not alter the colour of

* [Wöhler has obtained glucinum by decomposing its chloride by potassium;—in many of its properties it is analogous to aluminum.]

vegetable blues or yellows. When it is thrown down from an acid solution by an alkali, it exists in combination with water, as an hydrate. It forms sweet-tasted salts soluble in water, with the acids, and hence it gained its name, from *γλυκὺς*, sweet. From experiments on the quantity of ammonia necessary to decompose the muriate of glucina, I am inclined to adopt 54 as the number representing the earth, and supposing it a protoxide, 39 as the number representing the metal.

4. No compounds have been as yet examined in which glucinum can be supposed to exist, uncombined with oxygen. Glucina has not as yet been applied to any of the purposes of the arts, and its combinations in nature are very rare.

X. *Zirconum*.

1. There is a peculiar earth, which was discovered by M. Klaproth, in 1793, and which may be procured from a stone found in Ceylon, and called the jargon, or zircon, and likewise from the hyacinth, by the following process. The powder of these stones must be ignited for a long while with hydrate of potassa; the substance which is not dissolved by the hydrate of potassa, is principally zircon. The soluble matters must be separated by water, and the insoluble matter boiled in muriatic acid, and the solution so obtained, evaporated to dryness, and heated to the temperature of 212° . An aqueous solution of zircona in muriatic acid, is obtained by the action of water on the solid mass; and pure zircona is procured by decomposing this solution by solution of ammonia, and heating the powder obtained to redness.

2. There is the same evidence for believing that

zircona is a compound of a metal and oxygen, as that afforded by the action of potassium on the other earths. The alkaline metal when brought into contact with *zircona* ignited to whiteness, is for the most part converted into potassa, and dark particles, which when examined by a magnifying glass, appear metallic in some parts, of a chocolate brown in others, are found diffused through the potassa and the undecomposed earth.*

3. *Zircona* appears as a harsh whitish powder, without taste or smell; it possesses no action on vegetable colours, and is insoluble in water; its specific gravity is rather above 4. It is fusible at a lower temperature than any of the other earths; the heat of a good forge is sufficient to soften it. In a mass it is very hard and scratches rock crystal. *Zircona* is insoluble in water, but when precipitated from its solution, and dried at a low temperature, is found in the state of a hydrate; and has an appearance like that of resin or glue; its particles adhere together; the hydrate contains more than one-fifth of its weight of water. *Zircona* is soluble in the mineral acids and in solutions of alkaline carbonates. From experiments I have made on the comparative saturating powers of ammonia and *zircona*, I am disposed to give 85 as the number representing the earth, and 70 as the number representing the metal; supposing the earth to be a protoxide.

4. No substance has as yet been formed or examined, in which zirconum can be supposed to exist free from oxygen. It forms a crystallized muriate when dissolved

* [M. Berzelius has succeeded in obtaining zirconum pure by heating together potassium and dry fluoride of zirconum and potassium, and the after use of boiling water and of dilute muriatic acid. It is very similar to the two preceding metals.]

in muriatic acid; but the muriatic acid is expelled by heat, without any apparent union of chlorine and the metal. Its combinations, therefore, must be objects of future enquiry.

Zircona has not yet been found in sufficient quantities, to be applied to any of the purposes of the arts. It combines with the other earths, and forms compounds analogous to porcelain.

XI. *Silicum.*

1. Pure transparent quartz, or rock crystal, consists almost entirely of a peculiar earth called silex or silica: this earth may be procured from that stone, or from common flints, by igniting them in powder, with three or four times their weight of hydrate of potassa, or soda, in a silver crucible, making an aqueous solution of the substance so obtained, and adding to it any acid in quantities barely sufficient to neutralize the alkali, a gelatinous substance separates, which is silica combined with water; and the pure earth may be obtained by washing this substance well, and then igniting it to whiteness.

2. That *silica* consists of oxygen united to a peculiar inflammable basis, which is probably metallic, is shewn by many experiments. When iron is negatively electrified, and fused by the Voltaic battery in contact with hydrate of silica, the metalline globule procured contains a matter which affords silex during its solution; and when potassium is brought in contact with silica ignited to whiteness, a compound is formed consisting of silica and potassa, and black particles not unlike plumbago are found diffused through the compound. From some experiments I made, I am inclined to believe that these par-

ticles are conductors of electricity; they have little action upon water, unless it contain an acid, when they slowly dissolve in it with effervescence; they burn when strongly heated, and become converted into a white substance having the characters of silica; so that there can be little doubt, both from analysis and synthesis, of the nature of silica;* but no direct experiments have as yet been made upon the proportion of oxygen it contains.

3. Silica is a white powder, very analogous in its physical characters to the other earths; in its state of hydrate it is soluble in alkaline lixivia, and likewise in acids. It is separated from the common mineral acids by a very gentle heat, they rise from it in vapour, but it forms permanent compounds with boracic, phosphoric, and fluoric acids; its compound with phosphoric and boracic acids is a white powder, that with fluoric acid a permanent gas. From some experiments that I have made on the quantity of ammonia necessary to decompose the saturated solution of silica in muriatic acid, and from the composition of its gaseous fluoric combination, as ascertained by my brother Mr. John Davy, I estimate the number representing silica as 61; and as it seems to combine with double proportions of acids, I am inclined to regard it as a deutoxide, composed of 31 basis, and 30 oxygen.

4. Silica is one of the earths most generally diffused in nature. It forms, perhaps, the largest part of the solid surface of the globe. It is of great use in many of the arts; it is the basis of glass and porcelain, and

* [M. Berzelius has procured silicium by a method similar to that by which he succeeded in obtaining zirconum. In the trials he made of it, it did not appear to be a conductor of electricity, from whence he infers, it is not metallic. It combines with chlorine, and forms a volatile fluid.]

the art of manufacturing these substances depends upon the attraction of silica for other metallic oxides.

XII. *Ittrium*.

1. There is a mineral substance called Gadolinite, found at Itterby in Roslagen in Sweden, in which a peculiar earth was discovered in 1794, by Mr. Gadolin, and to which the name of *ittria* has been given. To procure this earth, the pulverized fossil must be digested for a considerable time in solution of muriatic acid. The solution obtained must be evaporated to dryness, redissolved in distilled water, and precipitated by caustic ammonia; the precipitate obtained must be digested with solution of hydrate of potassa; the remaining substance must be redissolved in solution of muriatic acid not used in excess, succinate of soda must be poured into this solution till all precipitation is complete; the filtered liquor must be decomposed by carbonate of soda, a white powder will fall down, which when ignited strongly, is pure *ittria*.

2. When ittria is treated with potassium in the same manner as the other earths, similar results are obtained; the potassium becomes potassa, and the earth gains appearances of metallization, so that it is scarcely to be doubted that ittria consists of inflammable matter, metallic in its nature, combined with oxygen.*

3. Ittria appears as a white powder, without taste, smell, or power of action on vegetable colours; it requires an intense degree of heat for its fusion; it is not soluble in water. No experiments have been made to ascertain whether it forms a hydrate with water, but this

* [This has been proved in a satisfactory manner by Wöhler, by the action of potassium on the chloride. Ittrium in many of its properties resembles glucinum and aluminum.]

is most probably the case ; its specific gravity is greater than that of any of the other earths, being more than 4.5. It combines with the acids, and forms sweet-tasted salts with those in which it is soluble. With acetic and sulphuric acids it forms crystals of an amethyst colour. It is not acted upon by solutions of caustic alkalies ; it is slightly soluble in solution of carbonate of ammonia.

4. It is probable that a compound of chlorine and ittria may be obtained ; but as yet no experiments have been made on any compounds of this substance, except such as contain oxygen ; the proportion of oxygen in ittria cannot be determined from any experiments hitherto published. According to Klaproth 55 parts of ittria combine with 18 parts of carbonic acid ; consequently, if it be supposed that the carbonate of ittria consists of one proportion of acid, and one of earth, the number representing ittria will be 126 ; and on the hypothesis that ittria consists of one proportion of metal, and one of oxygen, which is probable from all analogy, the number representing ittrium will be 111.

5. The compounds of ittria are very rare in nature, and as yet no application of this substance has been made to any of the purposes of the arts.

XIII. *Manganesum.*

1. The mineral called manganese has been referred to, page 167 ; it consists of a peculiar metal, *manganesum* united to oxygen. To procure the pure metal, solution of muriatic acid must be distilled from manganese in fine powder, the mixture strongly heated, and the process repeated till the washings in pure water give only a white precipitate with a solution of the salt called

prussiate of potassa and iron; an aqueous solution of potassa is then added to the mixture, so as to render it alkaline; the whole is then poured on a filtre, and the solid matter obtained well washed, dried, mixed with charcoal powder and oil, and intensely heated for half an hour in an infusible earthen crucible lined with charcoal powder; a number of small metallic globules will be obtained, which are globules of manganesum.

2. *Manganesum* was first procured in its pure form by Kaim and Gahn, between 1770 and 1775. It is of a grayish white colour; it has not much lustre; its hardness is nearly that of iron; the specific gravity is about 6.850. It is very brittle. It requires a higher degree of heat than iron for its fusion. It immediately tarnishes in the air, and becomes gray, brown, and at last black; when strongly heated in contact with oxygen, it burns with great brilliancy, throwing off vivid sparks; when heated in chlorine it takes fire and burns. It dissolves with effervescence in the mineral acids.

3. There are two definite combinations of manganesum and oxygen, one dark olive, and one brownish black. The first, or *olive oxide*, may be obtained by dissolving common manganese in nitrous acid, adding a little sugar, and precipitating by solution of potassa; a white powder is obtained, which must be heated to redness out of the contact of air, it is then the substance in question. The same body may be formed by precipitating the muriatic or sulphuric solutions of manganese by potassa, and treating the precipitate by heat; the white powder, when exposed to air, rapidly changes its colour to yellow, then puce colour, and lastly red brown: to be preserved it should be washed in *boiling* water, which contains little air, and the water driven off from it in a retort filled with hydrogen gas.

The *dark olive oxide of manganesum* in its pure form, when examined in large quantities, appears almost black, but when spread upon white paper its olive tint is apparent. It takes fire when gently heated, increases in weight, and gains a browner tint. It slowly absorbs oxygen from the air, even at common temperatures. It is the only known oxide of manganesum which dissolves in the acids without effervescence. The white powder produced by the action of acids on solutions of this oxide, is a compound of the oxide and water, or the *hydrated oxide of manganesum*; and the different tints that it assumes by exposure to air, seem to depend upon the formation of smaller, or larger quantities of the dark brown oxide, which probably retains the water contained in the white hydrate, and in this state is deep puce coloured; and when the water is expelled from it, it becomes dark brown, and then appears to be the same substance as the native oxide of manganesum, which may be called the *peroxide of manganesum*. The specific gravity of the peroxide is about 4; it is not capable of being combined with any of the acids; it gives off oxygen gas by a strong heat, as has been already stated, and by intense ignition it is partly or wholly converted into the first or olive oxide of manganesum. From some experiments that I have made on the two oxides of manganese, I conclude that the olive oxide consists of about 21 of oxygen to 79 of metal, and that the dark brown oxide contains nearly ten per cent. more of oxygen. According to these estimations, supposing the olive oxide a deutoxide, or an oxide containing two proportions of oxygen, the number representing manganesum will be 113; and the olive oxide will be represented by 143; and the dark brown oxide by 158, that is, it must be a tritoxide or an oxide con-

taining three proportions of oxygen. The white hydrated oxide of manganese appears from my experiments to contain about 24 per cent. of water. Hence it may be regarded as consisting of one proportion of olive oxide of manganese 143, and 34 of water, and the number representing it is 177. This hydrate is erroneously described in chemical books as the oxide of manganese containing the smallest quantity of oxygen; and there are many other cases in which hydrates have been confounded with oxides.

It has been supposed that there is a peculiar oxide of manganese, which may be procured by alkalis from the sulphuric solution; but when this solution is concentrated, the precipitate is the pale hydrate, having a very slight tint only of puce colour, apparently from the formation of a little of the other oxide, in consequence of the absorption of oxygen from the air dissolved in the alkaline solution used for its precipitation. It has been likewise conceived that there is a green oxide of manganese; the olive oxide becomes green by the action of potassa; but in this case a combination takes place between the alkali and the oxide. I have made many experiments on the native dark oxide by exposing it at different intervals for several hours to intense heat: in these cases it passed through different shades of brown and olive brown, and finally became dark olive. Colour is too indefinite a property to found a definite species upon; a mere change of temperature, without any evident change in composition, alters the colours of many bodies; and it is very probable that the different shades of colour of different precipitates from solutions of manganese, depend upon mixtures of the white hydrate with the puce coloured hydrate formed at the time of precipitation by the absorption of

oxygen from the air in the fluid, and the white hydrate seems to be always the result of the action of alkali on solutions, in cases when there can be no interference from the influence of free or loosely combined oxygen.

4. A compound of chlorine and manganesum may be obtained by combustion of the metal in chlorine, or by heating strongly the substance obtained by the solution of manganese in muriatic acid. When made in this last way, it appears as a pale pink coloured substance, and semi-transparent, and in brilliant scales. This compound has been described by Mr. John Davy as consisting of chlorine and manganese, and from his experiments may be considered as consisting of one proportion of the metal 113, and two of chlorine 134. It is probable that another compound consisting of one proportion of the metal and three of chlorine may be formed.*

5. Hydrogen, azote, sulphur, and charcoal, have no distinct chemical action on manganesum.

6. Phosphorus has been combined with manganesum by Pelletier; the *phosphoret* is a substance possessing metallic lustre and very combustible; its constitution has not yet been ascertained.

7. The action of boron, and the metals of the alkalis, and earths, on manganesum, has not yet been tried.

8. Manganesum in its oxidated form is of considerable use in certain arts. Its application for the production of chlorine has been already described. It is employed in glass-making, for depriving glass of colour: when in its state of full oxidation it gives a purple tint to glass, which is destroyed by thrusting a piece of

* [A perchloride has been formed by M. Dumas; a liquid at low temperatures, a gas at ordinary ones.]

wood into the melted glass, the inflammable matter of which seizes upon a part of its oxygen.

It is in some cases used to give colours to enamels in the manufacture of porcelain. The changes of colour of glass containing oxide of manganese according to its different states of oxidation are easily exemplified by adding a little dark oxide to powdered glass mixed with borax, and fusing them by the blow pipe: as long as the globule is preserved within the blue flame, where there is combustible matter, it remains colourless; but when it is exposed to air at the extreme point of the flame it becomes purple. I am inclined to believe that the deutoxide is the only oxide which enters into combination with vitrifiable substances; and that the peroxide when formed is mechanically diffused through the glass, and being produced only in very minute quantities is transparent and coloured. There is great reason to believe that the colouring matters of many gems are merely oxides finely divided in a state of mechanical diffusion through their substance.

XIV. *Zinc or Zincum.*

1. Zinc is procured for the purposes of commerce from various ores known by the names of calamine and blende or black jack. To obtain the metal from calamine, which is a combination of zinc with oxygen and carbonic acid, the ore is strongly ignited with charcoal or carbonaceous substances; the zinc, which is volatile, rises at a strong red heat, and becomes condensed in the cool part of the furnace or retort in which the process is carried on. Zinc is procured from blende by a similar operation, but the blende must be previously roasted, that is, exposed for a long while to a dull red

heat in a state of minute division. The zinc of commerce is seldom quite pure; to obtain it in this state white vitriol or sulphate of zinc is dissolved in pure water and exposed to the action of a plate of common zinc; this will separate any volatile metals that may happen to exist in the solution, which is then to be precipitated by subcarbonate of potassa: the white precipitate ignited with charcoal powder affords the metal.

2. Zinc is of a bluish white colour. Its hardness is nearly equal to that of copper. Its specific gravity varies from 6.8 to rather more than 7. When hammered it is 7.2. Its fusing point is 680° Fahrenheit, at a red heat in close vessels it volatilizes; and at this temperature in the atmosphere it burns with a brilliant bluish white flame. It has a certain degree of ductility, and when heated a little above 212° Fahrenheit, it is malleable, and when annealed it may be passed through rollers, and obtained in small thin sheets or leaves; it may be drawn into wire, the tenacity of which, according to Muschenbrock, is such, that a wire of $\frac{1}{10}$ of an inch in diameter will support a weight of about 26lbs. Its capacity for heat, according to Wilcke, is 0.102.

The atmosphere has but little effect on zinc at common temperatures; by exposure to the air for some time it acquires a grayish colour on the surface, which is owing to a partial oxidation. Zinc filings very slowly decompose water, hydrogen gas is evolved, and oxygen combines with the metal. The effect is rapidly produced when steam is passed over zinc at elevated temperatures. Zinc in thin leaves introduced into chlorine takes fire, and burns with a white light; even when in thick wire it may be made to burn in this gas by a gentle heat.

3. There is one well known combination of zinc with oxygen; it is obtained by the combustion of zinc in the atmosphere, or by the precipitation of solutions of zinc in acids by alkalies and subsequent ignition. When examined in a state of minute division, such as it appears when obtained by combustion, it is white, and similar to cotton in its appearance; but when examined in mass it has a tint of pale yellow. It becomes fluid at a white heat, and is capable of being volatilized by an intense white heat.* It is soluble in most of the acids, and in aqueous solutions of the fixed alkalies; when precipitated from its acid solutions by alkalies, it is in the state of combination with water, and a strong red heat is required for the expulsion of the water with which it is combined. From my experiments, and those made by my brother Mr. John Davy, it appears that the *white oxide of zinc* contains about 82 parts of metal and 18 of oxygen. M. Proust gives 80 to 20, which is not a wide difference. On the estimation of 18 per cent., supposing that the oxide of zinc consists of one proportion of oxygen and one of metal, the number representing zinc is 66 taking away the fractional part; the oxide of zinc is represented by 81: and the hydrate, supposing it to contain one proportion of water, will be denoted by 17 added to 81; but as yet no experiments have been made to shew that this is the composition of the hydrate. It has been supposed that there is a gray oxide of zinc produced by keeping zinc melted in the open air; and a yellow oxide formed by fusing the white powder produced by precipitation from acids, both containing less oxygen than the oxide just

* [These effects of a high temperature are doubtful; and on the Author's copy the passage is marked by a query; the oxide is commonly said to be fixed in the fire.]

described; but there are no facts to warrant the idea that these bodies are distinct compounds of zinc and oxygen. The gray powder formed upon the surface of melted zinc, I am inclined to consider as a mixture of the white oxide with small particles of unburnt zinc, and the yellow oxide the same as the oxide produced by combustion free from water.

4. When zinc is burned in chlorine a solid substance is formed of a whitish gray colour and semi-transparent. This is the only compound known of zinc and chlorine. It may likewise be made by heating together zinc filings and corrosive sublimate; it is as soft as wax, fuses at a temperature a little above 212° , and rises in the gaseous form at a heat much below the red heat. Its taste is intensely acrid, and it corrodes the skin; it acts upon water and dissolves in it, producing much heat, and its solution decomposed by an alkali affords the white hydrated oxide of zinc. The compound of zinc and chlorine has been called butter of zinc and muriate of zinc; following the nomenclature already proposed its name will be *zincane*.* From the experiments of my brother, Mr. John Davy, it consists of nearly equal parts by weight of zinc and chlorine; consequently it contains one proportion of metal and one of gas 66 and 67, and the number representing it will be 133.

5. It is not easy to combine zinc and sulphur. When a solution of sulphuretted hydrogen and an alkali is dropped into an acid solution of zinc, a whitish powder falls down, which has been supposed to be a *sulphuret of zinc*. When zinc and sulphur are heated together in close vessels the sulphur rises in vapour without uniting to the zinc; but it is stated by Mr. E. Davy, that in some experiments made in the laboratory of the Royal

* [Chloride of zinc.]

Institution, in which sulphur in vapour was passed over melted zinc, they united, and formed a white crystalline substance, analogous to the substance found in nature, and called phosphorescent blende. The proportions of the elements in the blendes, or supposed sulphurets of zinc, have not yet been ascertained with accuracy; but if some experiments on record can be depended upon, they must contain two proportions of metal to one of sulphur.

6. Zinc combines with phosphorus, when the metal is fused and the phosphorus brought in contact with it. The *phosphoret of zinc* was discovered by Pelletier; it is possessed of metallic splendour, and is of a dull gray colour analogous to lead; when hammered or filed it emits the odour of phosphorus. From experiments made on its composition in the laboratory of the Royal Institution by Mr. E. Davy, it is probable that it consists of one proportion of phosphorus and one of metal.

7. Zinc has not been combined with hydrogen, azote, or boron: zinc sometimes during its solution in acids leaves a residuum having the characters of carbonaceous matter: but no definite compound of zinc and carbon has as yet been described.

8. Zinc readily enters into union with the metals of the fixed alkalies; great heat is produced during the process, and metallic compounds or alloys are obtained, which rapidly decompose water and tarnish in the atmosphere.

9. Zinc is applied to a number of important uses: it is particularly employed in the manufacture of brass and tombacs; which consist of this metal combined with different proportions of copper. It is used by the Chinese in various alloys: some of its combinations are employed in medicine.

XV. *Tin or Stannum.*

1. Tin is procured from the native combinations of this metal with oxygen, known by the names of tin stones, or oxides of tin, by ignition with charcoal or carbonaceous substances. The metal obtained from these ores is not pure. To obtain it in a state of purity, metallic tin should be boiled for some time in solution of nitric acid, the white powder formed, should be well washed in pure water, and heated strongly in contact with about one-fourth of its weight of charcoal powder in a covered crucible for about half an hour; a button of pure tin will be found at the bottom of the crucible.

2. Tin has been known since the earliest periods of civilization; it was used in the time of Moses: it is mentioned by Homer, and was brought from Cornwall by the Phoenicians and Greeks, some centuries before the Christian era; it is called by Aristotle *Κασσιτερον Κελτικον*, or Celtic tin. The colour of tin is white, and resembles that of silver. Its hardness is greater than that of lead, and less than that of zinc. Its specific gravity is 7.291, and it is somewhat increased by hammering; it is very malleable, and may be extended into extremely thin leaves. Tin foil is about the $\frac{1}{1000}$ part of an inch thick; it has comparatively little ductility or tenacity. Tin is flexible, and when bent produces a crackling noise. It has a slight taste, and when rubbed emits a peculiar smell. It fuses at 442° Fahrenheit, but requires an intense degree of heat for its evaporation. Its capacity for heat, according to Dalton, is .07. It acquires a slight tarnish by exposure to the atmosphere, but undergoes no further change. It is not affected by water at common temperatures, but when steam is

passed over red hot tin it is decomposed, oxide of tin is formed, and hydrogen gas is evolved. When heated strongly in air, it takes fire and burns with a pale white light; when burnt upon charcoal by a stream of oxygen gas, the colour of the flame is white, edged with violet. Tin foil burns when very gently heated in chlorine.

3. There are two definite combinations of tin and oxygen: the first, which may be called the *protoxide*, is gray; the second, which may be called the *peroxide*, is white; the first is formed by heating tin in the air, or by dissolving tin in muriatic acid, and precipitating the solution whilst recent, and before it has been exposed to air by solution of hydrate of potassa, not added in excess. This substance, after being heated to whiteness, is the protoxide of tin; and it is converted into the peroxide by being boiled with diluted nitric acid, dried by evaporation and heated to redness. From experiments which I have made, it appears that the protoxide of tin contains about 13·5 per cent. of oxygen, and from experiments made by Mr. John Davy, the peroxide is composed of about 24 of oxygen and 76 of metal. These oxides are difficultly fusible bodies, insoluble in water, soluble in diluted oil of vitriol, and in fixed alkaline solutions. Computing from their composition, and supposing one to consist of 1 proportion of tin and 1 of oxygen, and the other of 1 of tin and 2 of oxygen, the number representing tin will be 110, and the number standing for the protoxide will be 125, and that standing for the deutoxide, or white oxide, 140. Both these oxides appear capable of combining with water to form hydrates; and when precipitated from their acid solutions, they always contain water, but experiments are wanting to determine the quantity: both are insoluble in water.

4. As there are two combinations of tin with oxygen, so there are two which it forms with chlorine. When tin is burnt in chlorine a very volatile clear liquor is formed, a non-conductor of electricity, and which, when mixed with a little water, becomes a solid crystalline substance, a true muriate of tin containing the peroxide of tin. This liquor, which has been called Libavius's liquor, from its discoverer Libavius, may be likewise procured by heating together tin filings and corrosive sublimate, or an amalgam of tin and corrosive sublimate. It consists, according to the analysis of Mr. John Davy, of 2 proportions of chlorine 134, and 1 of tin 110; and according to the proposed principles of nomenclature, its name will be *stannanea*.* The other compound of tin and chlorine is a gray semi-transparent crystalline solid; it may be procured by heating together an amalgam of tin and calomel: it dissolves in water and forms a solution which rapidly absorbs oxygen from the air, depositing oxide of tin. This compound of chlorine and tin, it appears from the experiments of Mr. John Davy, who first described it, consists of 1 proportion of tin 110, and 1 of chlorine 67; it may be called *stannane*.†

5. There are 2 *sulphurets of tin*, one may be made by fusing tin and sulphur together; it is of a bluish colour and lamellated structure; and from the experiments of my brother consists of 1 proportion of tin and 1 of sulphur 110 and 30. The other sulphuret of tin, or the supersulphuret, is made by heating together the peroxide of tin and sulphur; it is of a beautiful gold colour, and appears in fine flakes; it was formerly called *aurum musivum*. It has been supposed by Pelletier and

* [Perchloride of tin.]

† [Chloride of tin.]

Proust to contain tin in an oxydated state, but Mr. John Davy has shewn that this is not the case; and that it consists merely of 1 proportion of metallic tin combined with 2 proportions of sulphur, 110 and 60; so that the number representing it is 170.

6. Tin combines with phosphorus when the two substances are heated together. As yet only one compound is known, which appears from my experiments to consist of 17 per cent. of phosphorus, and therefore may be regarded as composed of one proportion of tin and one of phosphorus, 110 and 20. The *phosphoret of tin* has a metallic appearance, is so soft that it may be cut with a knife; the phosphorus burns when it is gently heated in the air.

7. Tin has not been combined with hydrogen, azote, carbon, or boron; it readily unites to the metals of the fixed alkalies, and forms alloys which speedily tarnish in the air, and which effervesce in water. It unites with zinc by fusion; the alloy is harder than zinc and stronger than tin.

8. Tin is a metal of great use, and of various application; it is an important ingredient in pewter, bell-metal, and bronze; it is employed to cover culinary vessels, as tin plate; some of its acid compounds are used in dyeing. Tin is almost always found in nature in the oxydated state, and in the crystalline form; and it appears from the analysis of Klaproth that the native oxide or tin stone of Cornwall must contain one proportion of tin and two of oxygen. All the well known combinations of this metal are such as they ought to be according to the theory of definite proportions, and its compounds with oxygen, sulphur, and chlorine, afford similar results, which correspond with the numbers gained from its simplest combination.

XVI. *Iron or Ferrum.*

1. The iron of commerce is obtained from various ores of that metal, in which it exists combined with oxygen, by intense ignition with carbonaceous substances: the purest iron is made from an ore called hæmatites by ignition with charcoal; and the metal is hammered whilst in a soft state, exposed to air, till it becomes ductile. Iron was known in the time of Moses, and used for the manufacture of swords, knives, and axes. It is referred to in the Iliad and the Odyssey of Homer. A ball of iron was one of the prizes said by that poet to have been offered by Achilles at the funeral rites of Patroclus; and the effect of the burning brand feigned to have been thrust by Ulysses and his companions into the eye of Polyphemus, is compared by the poet to that of the hot iron plunged into water by the smith. The soft iron employed in the useful arts is free from any alloy,* and therefore may be used for the purposes of chemistry.

2. The colour of iron is well known, and its other sensible properties; its specific gravity is about 7·7. Its malleability, though considerable, is inferior to that of gold, silver, and copper. Its ductility and tenacity are, however, greater; it may be drawn into extremely fine wire, and a wire of 0·078 of an inch in diameter is capable of supporting 549·25 lbs. It requires the highest heat of a wind furnace for its perfect fusion; it is attracted by the magnet, and is capable of acquiring magnetism, though in its unalloyed state it retains it only for a very short time. When iron is exposed to the atmosphere it slowly combines with oxygen and carbonic acid, and its surface becomes covered with a yellowish substance well known by the name of *rust*. It burns with great splendour in oxygen gas, as has

* [It contains carbon; according to Berzelius ·5 per cent.]

been stated page 169. At common temperatures it slowly decomposes water.* Hydrogen gas is evolved, and oxygen combines with the metal. The effect is rapidly produced when the vapour of water is passed over red hot iron. When gently heated in chlorine it takes fire and burns with a deep red light.

3. The combinations of iron and oxygen have been referred to, page 80. The *black* and the *red-brown oxides* are the only oxides of this metal known; these substances, which have been considered as green and white oxides, are in fact hydrated oxides. The black oxide of iron, which may be regarded as a deutoxide, as containing two proportions of oxygen, is formed by the rapid combination of iron in oxygen. The red-brown oxide, which must be considered as a tritoxide, may be produced from the black by keeping its powder red hot for a considerable time in contact with the atmosphere often changing the surface. Reasoning on the composition of these oxides, 103 must be the number representing iron; and the black oxide, or the deutoxide, consists of one proportion of iron 103, and two of oxygen 30; and the brown-red oxide, or tritoxide of 103 metal, and 45 oxygen. Both these oxides are soluble in the common acids. The black produces pale green solutions; the brown-red, deep yellow solutions: the solutions of triple prussiate of potassa, precipitate the solutions of the black oxide white; those of the red bright blue. When solutions of these oxides are acted upon by solutions of pure alkalies, a white precipitate, having a tint of green or olive, is thrown down from the solution containing the black oxide; and an orange coloured precipitate from the solution con-

*[This, at common temperatures, or even at 212° , has not been confirmed by later experiments.]

taining the red-brown oxide; and both these precipitates, I find, are the oxides combined with water, or hydrates. The pale hydrate, when exposed to air, rapidly changes in colour, first becomes pale olive, then dark olive, then olive brown, and last of all orange; so that there is strong reason to conclude that the colours of different precipitates depend upon their being composed of mixtures of the two hydrates; and solutions of the black oxide cannot be exposed to air for a moment without being changed by the absorption of oxygen. I have made no experiments to ascertain the composition of the two hydrates; probably the white contains two proportions of water. It would seem from the experiments of Dr. Thomson that there is a black hydrate formed by passing steam over iron, and which probably consists of one proportion of iron, two of oxygen, and one of water. Mr. Daubuisson has described a native hydrated oxide of iron.

4. There are two compounds of iron and chlorine. The one containing the largest proportion of chlorine is formed by burning iron wire in the gas. It is a very beautiful substance, of a bright yellowish brown colour. It has a high degree of splendour, and is very volatile, rising in the gaseous state at a temperature a little above that of boiling water, and crystallizing in small iridescent plates. It acts with violence upon water, and forms a solution of red muriate of iron. I have called it *ferranea*, and I find by analysis that it consists of one proportion of iron 103, and three proportions of chlorine 201.

The other compound of chlorine and iron has been analysed by my brother, Mr. John Davy; it contains proportions which agree nearly with one of iron 103, and two of chlorine 134. It is a dark gray opaque substance, fusible at a red heat, and not rising in vapour

at the point of fusion of glass. It forms a solution of green muriate of iron by its action upon water; it may be named *ferrane*.

5. No combinations of iron with hydrogen or azote are known; but this metal readily combines with sulphur. There are two well known sulphurets of iron; one is formed by heating iron filings and sulphur together out of the contact of air; they combine with great energy, producing the effect of ignition. The *sulphuret of iron* formed in this way is of metallic splendour, and a dull yellow colour. This compound is found in nature, and has been analyzed by Mr. Hatchett. It is magnetic, and has been called magnetic pyrites. The other sulphuret of iron, which may be called the *supersulphuret*, has not yet been made artificially, but it is found abundantly in metallic veins: it is of a bright yellow colour, and often crystallized in cubes. According to Hatchett and Proust, the sulphuret of iron consists of about 63 of iron to 37 sulphur, and the supersulphuret of about 46 to 54; so that the quantity of iron remaining the same, the last sulphuret contains nearly double as much sulphur as the first; and iron being represented by 103, the proportions are not very remote from two of sulphur 60 in the sulphuret, and four of sulphur 120 in the hypersulphuret.

6. Iron is capable of combining with phosphorus; but the proportions of the elements of *phosphoret of iron* have not been ascertained; nor is it known whether more than one compound of this kind exists. The phosphoret may be made by passing phosphorus in vapour over ignited iron. It is very brittle, of a dark steel gray colour, and of the specific gravity of 6·7. This substance, which was first found in the peculiar iron called cold short iron, because it is brittle when

cold, was once supposed by Bergman and Meyer to be a peculiar metal; but Klaproth discovered its real nature. It may be formed likewise by heating together phosphoric acid, iron, and charcoal.

7. Iron is capable of combining with carbon; and *steel*, perhaps the most important substance employed in the useful arts, is one of the results of their combination. Steel is usually made by a process called *cementation*, which consists in keeping bars of iron in contact with powdered charcoal in a state of ignition for ten or twelve days, in earthen troughs, or crucibles, the mouths of which are closed with clay. Cemented steel is made into the substance called *cast steel* by being fused in a close crucible with a mixture of powdered glass and charcoal. Steel is possessed of the power of receiving very different degrees of hardness by different applications of heat or cold. When it is heated to redness, and suffered to cool slowly, it is found very soft; but if plunged into cold mercury or water, it acquires extreme hardness; and by heating hard steel to different degrees, it receives different degrees of temper, from that which renders it proper for files, to that which fits it for watch springs. In the process of tempering, the steel changes colour when exposed to the air. Between 430° and 450° Fahrenheit, according to Mr. Stoddart, it assumes a pale yellowish tinge; at 460° the colour is a straw yellow, and the metal is of the temper necessary for pen-knives, razors, and fine edged tools. The colour gradually deepens as the temperature rises higher, and it passes through brown, red, and purple, to 580 , when it becomes of an uniform deep blue. These changes of colour seem to depend upon the formation of a plate of oxide, which increases in thickness the higher the temperature. Steel is of greater specific

gravity than iron; when the metal is hammered it is about 7·8. When it is acted upon by an acid, such as diluted nitric acid, a black spot appears upon it from the separation of the carbonaceous matter. Steel is attracted by the magnet, and is capable of receiving permanent magnetism. It is not easy to determine the exact quantity of carbon in steel, but it consists of several proportions of iron to one of carbonaceous matter. Different specimens of steel are said, on the authority of Bergman, Vauquelin, and Mushet, to contain only from $\frac{1}{140}$ to $\frac{1}{60}$ of carbon.

Iron has been converted into steel by cementation with diamond by Morveu and Sir George Mackenzie.

Plumbago, or *black lead*, as has been mentioned page 231, is a compound of carbon, with $\frac{1}{25}$ its weight of iron. There is a substance formed in iron foundries called *kish*, of a brilliant appearance, usually in thin scales, analogous to plates of polished steel. It consists chiefly of carbonaceous matter united to iron, and a little manganese.

8. When iron and charcoal are strongly ignited with boracic acid, the iron produces, during its solution, boracic acid, as M. Descotils has shewn. Hence it is probable, as MM. Gay Lussac and Thenard have supposed, that iron is capable of combining with boron.

9. Iron is capable of combining with potassium and sodium; these alloys are more fusible and whiter than iron, and effervesce copiously in water. There is great reason to believe that alloys may be formed of iron and the metals of the earths. *Cast iron*, which is produced by fusing iron ores with pitcoal, during its conversion into malleable iron, affords about one-fourth of its weight of a glass, which consists of silex, alumine, lime, oxide of iron, and oxide of manganese. In the process for

reducing cast iron into malleable iron called *blooming*, the iron, after being fused in a forge by a fire of charcoal, is hammered, whilst in a soft state, on an anvil by a large hammer worked by water; a vivid combustion, which seems to be connected with the formation of the glass and the oxides, takes place on the surface of the mass: that the earths are formed by the oxidation of metals combined in the cast iron, seems probable from the circumstance of the combustion; and the idea is confirmed by the distinct metallic character of cast iron; it is white, crystallized, and has all the appearances of a perfect alloy. Specimens of cast iron usually contain likewise sulphur and carbon.

10. Manganese forms very readily binary combinations with iron; the alloys have a white colour, and are very brittle. Iron likewise combines with tin. By fusing the two metals together Bergman obtained two alloys; the first containing 21 parts of tin, and one part of iron: the second two parts of iron and one of tin. The first was very malleable, harder than tin, and not so brilliant; the second scarcely malleable, and very hard. The formation of *tin plate* depends upon the chemical attraction between the two metals. Tin plate is formed by dipping thin plates of iron into melted tin; the iron must be very clean. It is usual to add $\frac{1}{10}$ of copper to the tin to prevent it from forming too thick a coat.

11. To describe the uses of iron would require volumes; as it is the most generally diffused metal, so it is likewise the most important in its applications to the purposes of society. By means of it the earth has been cultivated and subdued. Without iron, houses, cities and ships could not be built. It is subservient both to the common and the refined arts; it forms the

machinery by which the most important mechanical powers are generated and applied. Its uses have awakened human industry, and made it more efficacious, and have offered an infinite variety of resources to ingenuity and talent.

XVII. *Lead or Plumbum.*

1. The lead of commerce is chiefly procured from certain ores in which it is combined with sulphur. The sulphur is expelled or burnt by a long continued heat in a reverberatory furnace, and the metal is obtained by fusion. To procure pure lead, a solution of the lead of commerce in nitric acid, largely diluted with water, may be precipitated by zinc; or a solution of acetate of lead, i. e. sugar of lead may be used. The arborescent brilliant metallic substance produced from solution of sugar of lead by zinc, is generally *pure lead*.

2. Lead was known in a very early age of the world. It is often mentioned by Moses; and is described by Homer as in common use at the period of the Trojan war,

Lead is of a bluish white colour, but soon tarnishes by exposure to the air. It is the softest of the common metals. Its specific gravity is 11·352, and is not increased by hammering. It is very malleable, but not very ductile. Its tenacity is such, that a wire of $\frac{1}{126}$ of an inch in diameter supports only 18·4 pounds. Its point of fusion is 612° , but an intense degree of heat is required for its evaporation. It combines with oxygen slowly, at the temperature of its fusion, and burns when strongly ignited in the atmosphere; when a current of oxygen gas is thrown upon it in this state, the flame it

emits is of a brilliant whiteness, and it sends off a dense smoke. When heated in chlorine it unites to it, but does not inflame.

3. Lead combines with oxygen in different proportions; and three of its combinations with this substance appear to be well defined and distinct bodies. Two of the oxides of lead may be formed by heat with accession of air; the one is *massicot*, the other is *minium*, or red lead. When lead is heated in contact with the atmosphere, it soon becomes of a dirty yellow, or yellowish green colour, and at length of a pure yellow colour. This oxide is massicot, and is the oxide existing in the different salts of lead; when precipitated from these salts by caustic alkalies, it falls down in combination with water, and appears as a white *hydrated oxide of lead*; the water may be expelled from it by a strong red heat. From the experiments of Vauquelin and Klaproth, it may be concluded that this oxide of lead contains about 7 per cent, of oxygen. Litharge is this oxide of lead, according to Dr. Thomson, mixed with a little carbonate of lead: litharge is formed during the extraction of silver from lead by the calcination of the lead, and the carbonic acid is acquired from the carbonaceous matter burnt in the flame of the furnace.

Massicot is fusible at a strong red heat, and appears, when fused, as a yellow glass, insoluble in water, without taste or smell, and of great specific gravity.

The first oxide of lead by being heated moderately in contact with air, for a considerable time, combines with an additional quantity of oxygen, and then becomes of a beautiful red colour, in which state it is called minium, or red lead. 100 parts of lead carefully and slowly converted into minium become between 110 and 111 parts; so that there is strong reason to believe

that the quantity of lead being the same, the oxygen in minium is to that in massicot, as 3 to 2. Minium exposed to a strong red heat, gives off from 3 to 4 per cent. of oxygen gas, and becomes massicot.

When nitric acid is digested upon minium, a part is dissolved, but a puce-coloured powder remains, which contains more oxygen than minium, and the formation of which seems to be owing to the circumstance that the oxide which dissolves during its solution, becomes massicot, and affords oxygen to the undissolved portion, so as to convert it into a new substance. The *puce coloured oxide of lead*, long dried at 212° , loses from 6 to 7 parts per cent. during its conversion into massicot by fusion; so that it may be considered as containing twice as much oxygen as that oxide, the proportion of lead being considered as the same. On these views massicot will be a deutoxide of lead, minium a tritoxide, and the puce-coloured oxide a tetroxide; and the number representing lead will be 398; and the oxides will be composed respectively of 398 of metal, and 30, 45, and 60, of oxygen.

4. One combination only of lead with chlorine is known; it may be obtained directly by heating lead in chlorine, or by decomposing the oxides of lead by chlorine, in which case oxygen is expelled, or by acting on oxides of lead by muriatic acid gas, when water is formed. The combination of chlorine and lead is a dull whitish semi-transparent substance, fusible at a heat below redness, and volatile at an intense heat. This substance has a sweetish taste, and is soluble in 22 parts of cold water. It was called horn lead by the old chemists, and improperly muriate of lead by modern chemists. The name proposed for it is *plumbane*.*

* [Chloride of lead.]

According to my experiments made on the absorption of chlorine by lead, it contains 401 of lead to 131 of chlorine, which agrees very nearly indeed with one proportion of metal, and two of chlorine; and this compound decomposed by alkalies affords the oxide containing two proportions of oxygen.

5. Sulphur is easily made to unite with lead by a gentle heat. One combination only of these bodies is certainly known, and it is the same as the substance found in nature, referred to in the beginning of this section, and called *galena*. It is very brittle, brilliant, and of a deep bluish gray colour. It is less fusible than lead, and crystallizes in cubes. 100 parts of lead, in becoming the sulphuret, unite to about 15 parts of sulphur; which gives the *sulphuret*, as consisting of one proportion of metal and two proportions of sulphur.

6. A compound of lead with phosphorus may be formed by fusing together equal parts of filings of lead and phosphoric acid. It is of a silver white colour with a shade of blue; may be cut with a knife, but is brittle under the hammer. The same substance may be formed by bringing phosphorus in contact with melted lead. According to Pelletier it consists of 88 parts lead and 12 of phosphorus, which gives nearly 3 proportions of phosphorus 60, to one of lead 398.

7. There are no known combinations of lead with hydrogen, azote, carbon, or boron.

8. Lead unites by fusion with the metals of the fixed alkalies, and forms compounds which tarnish in the air, and are readily decomposed by the agency of water.

9. Lead combines with zinc, tin, and iron. Its alloy with iron is made with great difficulty, and has not been accurately examined. The alloys of zinc and lead may be easily made by fusion. These alloys are

harder than zinc, and ductile. In whatever proportions the metals are melted together, the mass, on cooling, is found to contain them in a state of chemical union or intimate mixture. Lead and tin combine in a similar manner; this alloy is harder and more tenacious than tin. It is said by Muschenbroeck that these qualities exist in the highest degree in the alloy, when it is composed of 3 parts of tin, and one of lead; which quantities nearly correspond with single proportions of each of the two metals. This mixture is often employed to cover copper vessels; and, as appears from the experiments of M. Proust, is difficultly acted upon by vegetable acids, and when acted upon, the tin is dissolved, and not the lead; so that such vessels may be safely employed for culinary purposes.

10. Lead is very extensively used both in the common and refined arts. Its oxides, and some of its saline combinations, are extensively applied in painting; white lead is the deutoxide combined with carbonic acid. Both massicot and minium are common pigments. The deutoxide combined with chromic acid forms the most beautiful and permanent deep yellow known. Lead is used as an ingredient in various solders; it is applied for covering houses and churches. It will be unnecessary to dwell upon its still more familiar applications. Its oxide forms an important part of flint glass, and is used in various enamels and pastes.

XVIII. *Antimony or Antimonium.*

The ancients were acquainted with certain ores of antimony; the most common of them, the sulphuret of antimony, was employed by the ladies of the oriental

countries to tinge the extremity of the eyelid black, for the purpose of giving greater brilliancy of effect to the pupil. Basil Valentine is the first chemist who has described the process of extracting antimony from the sulphuret, though it does not appear that he was the inventor of this process. He published his *Currus Triumphalis Antimonii* towards the end of the fifteenth century.

To procure antimony, the common antimony sold by druggists, which appears as a series of crystals, like needles, possessing the metallic brilliancy, and which are composed of the metal and sulphur, are ignited with half their weight of iron filings, and a quarter of their weight of nitre added when they are in fusion; the antimony will be found in the bottom of the vessel in which the experiment is made. To obtain it quite pure, it may be dissolved in aqua regia, water is added to the solution, a white powder will fall down; this is to be ignited for about twenty minutes, with twice its weight of tartar, when the metal will be produced.

2. Antimony is of a brilliant white colour with a shade of blue, its hardness is equal to that of zinc. Its specific gravity is about 6.8. It is very brittle, and may be easily pulverised. It has little tenacity. It fuses at about 810° Fahrenheit. On cooling it crystallizes, and its laminated structure is owing to the new arrangement of its parts. It is but little affected by exposure to the air or water at common temperatures; but when the vapour of water is passed over red hot antimony, it acts so powerfully upon the water, as to decompose it with explosions.

3. Two combinations of antimony with oxygen are known; one, the *fusible oxide*, is obtained by dissolving antimony in muriatic acid by heat, and adding water to

the concentrated solution, a white powder falls down, which, when washed with a solution of subcarbonate of potassa, and afterwards with distilled water, is a combination of the fusible oxide with water, and by fusion at a red heat it becomes the pure oxide. This substance is of a dirty yellowish white colour. It crystallizes by slow cooling after fusion. By being strongly heated in contact with the atmosphere, in volatilizing, it combines with more oxygen, and condenses in white crystals of a silvery lustre; and this substance is the *peroxide* or antimony saturated with oxygen.* This oxide is much less fusible and volatile than the other, and is more difficultly combined with acids. The fusible oxide, in its combination with water, was for a long while called the powder of Algaroth, from its discoverer Algarotti. Antimony burns, when heated strongly in the air, with a faint white light, and produces the second or the volatile oxide, which rises from it in the form of a dense white smoke. From experiments made on these two oxides by Mr. John Davy and myself, it appears that the fusible oxide contains $\frac{2}{3}$ as much oxygen as the fixed infusible oxide, supposing the metal to be the same in both; and calculating from his experiments on the fusible oxide, the number representing antimony is 170; the fusible oxide may be considered as consisting of 170 metal, and 30 of oxygen, and the peroxide of 170 metal, and 45 oxygen.

4. Antimony burns spontaneously when powdered and thrown into chlorine. In this way the only known compound of antimony and chlorine, *antimonane*,† or

* [Another oxide, containing a larger proportion of oxygen, has since been discovered by Berzelius, called, from its acid properties, Antimonic acid.]

† [Chloride of antimony.]

butter of antimony, is formed.* It is a soft semitransparent substance, of a yellowish white colour, very fusible, volatile at a moderate degree of heat. It crystallizes in parallelopipeds. It is a very caustic and corrosive substance; it acts with great energy upon water: with a small quantity of water it heats violently, and forms a solution; a large quantity precipitates the fusible oxide of antimony, and muriatic acid is found in the solution. Antimonane may be likewise formed by the distillation of a mixture of powdered corrosive sublimate and antimony. From the experiments of Mr. John Davy, it appears to contain 44 per cent. of chlorine, and therefore may be regarded as consisting of 1 proportion of metal and 2 of chlorine.†

5. Sulphur and antimony are readily combined by fusing them together, when they form a compound of metallic appearance, similar to the natural sulphuret, and which is much more fusible than antimony: according to Proust it contains about 25 per cent. of sulphur, and may therefore be considered as consisting of 1 proportion of metal, and 2 proportions of sulphur, 170 and 60.

6. Antimony has not yet been combined with hydrogen, azote, carbon, or boron.

7. Antimony combines with phosphorus by fusion. According to the experiments of M. Pelletier, the *phosphoret* is white, brittle, and has the metallic lustre; its composition has not been determined.

8. Potassium and sodium may be combined with antimony by fusion; they form alloys very similar to

* [Two other chlorides have since been formed, a bichloride and a perchloride;—the latter a volatile liquid, resembling in many of its properties the liquor of Libavius.]

† [According to Berzelius, it is a sesquichloride.]

those they form with lead and tin in their obvious properties. Antimony may be combined with all the other metals which have been described. The alloy of tin and antimony is employed in the arts, particularly for making music plates. Antimony very much impairs the magnetic properties of iron. The alloy of lead and antimony is used for printers' types; and for this purpose it is formed of 16 parts of lead and 1 of antimony. The oxides of antimony are used for giving a yellow colour to glass. Various combinations of antimony are employed in medicine.

XIX. *Bismuth or Bismuthium.*

1. The bismuth of commerce is procured from ores which usually contain it in the metallic state, or combined with sulphur, by roasting, and ignition with charcoal. The metal may be obtained in a state of purity by dissolving the ore in strong nitric acid, and adding water to the solution, a white precipitate will appear; it is to be washed, dried, and heated to a dull red for about 20 minutes, with a little oil, and some black flux, (a substance made by heating together nitre and tartar;) a globule of metal will thus be procured.

2. The ores of bismuth were first described by Agricola before 1530; the properties of the pure metal were not known before the middle of the last century. The colour of bismuth is white with a slight tint of red. It is nearly of the same hardness as copper. Its specific gravity is 9.822, and it is increased by hammering. It is brittle; it cannot be drawn into wire. Its tenacity is such that a rod $\frac{1}{10}$ of an inch in diameter sustains a weight of about 29lbs. It fuses at about 476° Fahrenheit, and if slowly cooled crystallizes in tubes: if it is

exposed to a strong heat in close vessels it sublimes unaltered. Bismuth acquires a superficial tarnish by exposure to the air; it is not affected by water.

3. One combination only of bismuth and oxygen is certainly known.* When bismuth is kept at a dull red heat in open vessels, its surface soon becomes tarnished; and by exposing fresh surfaces the whole may be converted into an oxide. When heated more intensely in the atmosphere, or in oxygen gas, it burns with a bluish flame, and a yellow oxide is formed, which fuses at an elevated temperature. The oxides formed by slow or rapid combustion are of the same kind. When in powder, they are yellow; when fused, they form a yellowish green vitreous mass. The oxide of bismuth requires an intense degree of heat for its volatilization. When this oxide is precipitated from its solutions in acids by water or alkalies, it appears as a white powder, which probably is a compound of the oxide and water. Klaproth has shewn that 100 parts of bismuth, by treatment with nitric acid and water, produce about 123 parts of the white powder. This powder has been called magistery of bismuth. Geoffroy found 100 parts of bismuth become 110 parts by exposure to heat and air; but in his experiments probably some of the metal escaped oxydation. John Davy has found the yellow oxide to contain in 100 parts 90 parts of metal, and 10 of oxygen, and this estimation is very near that of Bucholz; and supposing the oxide to consist of 1 proportion of metal, and 1 of oxygen, the number representing bismuth will be 135.

4. Bismuth, when thrown in fine powder into chlorine, takes fire, and burns with a pale blue light; in this case

* [Another oxide, a peroxide of a brown colour, has since been obtained, analogous to the puce-coloured oxide of lead.]

the only known compound, bismuth and chloride, is formed. It has been called *butter of bismuth*. It may be called *bismuthane*.^{*} It is an easily fusible substance, volatile at a moderate heat; its colour is grayish. It corrodes the skin, and is readily decomposed by water; the bismuth combines with the oxygen of the water; the chlorine with its hydrogen. From the experiments of Mr. J. Davy, it appears that bismuthane contains 33·6 per cent. of chlorine, and therefore may be considered as consisting of 1 proportion of metal, and 1 of the gas, 135 and 67.

5. There are no known combinations of bismuth with hydrogen, azote, carbon, or boron.

6. Bismuth combines with sulphur when they are fused together; the *sulphuret* is of a bluish gray colour, and has metallic lustre. According to Mr. J. Davy's experiments it contains about 18 per cent. of sulphur. By this estimation, the sulphuret of bismuth must contain about one proportion of metal to one of sulphur.

7. Bismuth appears to have little affinity for phosphorus, the attempts hitherto made to form this compound have been unsuccessful.

8. The action of the metals of the fixed alkalies on bismuth is similar to that which they exert on other easily fusible metals.

9. Bismuth forms alloys with all the metals which have been described, except zinc: these alloys have been little examined. It sometimes enters into the composition of pewter; and it forms a principal part of Newton's fusible metal. This alloy is composed of 8 parts of bismuth, 5 of lead, and 3 of tin, and melts at a temperature below that at which water boils.

^{*} [Chloride of bismuth.]

10. Bismuth is not of much use in the arts. The white hydrate is employed as a pigment, but is not very permanent, becoming yellow by the action of light. It is probable that the Roman ladies used a preparation of bismuth for whitening the skin; for Martial, in speaking of a lady who made too free a use of cosmetics, describes her as afraid of the sun.

This metal is sometimes employed in alloys to make easily fusible solders. The white hydrate has been lately employed in medicine, as a remedy in spasmodic affections of the stomach.

XX. *Tellurium.*

1. Tellurium was discovered by Klaproth, in 1798, and was procured by him from an ore found near Zalethna in Transylvania, in which it exists, in alloy with gold, lead, and silver. The process for obtaining the metal is very simple; the ore is dissolved in aqua regia made of a mixture of 1 part strong nitric acid and two parts muriatic acid. When the solution is saturated, water is to be added, a white powder falls down, which when dried and heated in a retort of glass, with $\frac{1}{10}$ of its weight of charcoal powder, will afford pure *tellurium*.

2. Tellurium is of a colour nearly the same as that of antimony. It easily fuses, and rises in vapour at a strong red heat. It burns, when heated in the air, with a vivid bluish green flame, sending off a dense white smoke. Its powder takes fire in chlorine. Its specific gravity is 6.115.

3. One *oxide of tellurium* only is known, the substance formed by combustion of the metal;* it is white with a

* [Another compound of the metal and oxygen has since been formed,—the telluric acid.]

tint of yellow, when examined in the mass. It fuses by a strong heat, and requires a very high temperature for its volatilization. When precipitated from its acid solutions, it is found in union with water, as a white hydrate. According to Klaproth, 178 grains of oxide of tellurium afford 148 grains of metal: supposing the oxide to consist of one proportion of oxygen, and one of metal, the number representing tellurium will be 74.

4. When tellurium is burnt in chlorine an easily fusible substance is formed, which rises in vapour at a strong heat, and crystallizes. Its colour is white: it is semitransparent; when decomposed by water, it affords the white hydrated oxide. From my own experiments it appears this compound, or *tellurane*, consists of 2 in weight of metal to 1.83 of chlorine: it may therefore be regarded as composed of one proportion of metal 74, and of chlorine 67.

5. Tellurium and hydrogen are capable of being combined. To make this combination, hydrate of potassa and oxide of tellurium are ignited with charcoal, and the mixture acted upon, by diluted sulphuric acid, in a retort connected with a mercurial pneumatic apparatus; an elastic fluid will be generated, which consists of hydrogen combined with tellurium. It is possessed of very singular properties. It is soluble in water, and forms a claret coloured solution. It combines with the alkalies. It burns with a bluish flame, depositing oxide of tellurium. Its smell is very strong and peculiar, not unlike that of sulphuretted hydrogen. I discovered this elastic fluid in August, 1809. When tellurium is made the electrical negative surface in water in the Voltaic circuit, a brown powder is formed, which appears to be a solid combination of hydrogen and tellurium, and which was first observed by M. Ritter, in

1808 : a claret coloured solution of the gas is likewise formed when the water is free from air. The composition of *telluretted hydrogen* gas, and of the solid *hydruret of tellurium* has not been yet ascertained.

6. Tellurium has not been combined with azote, carbon, or boron. No experiments are on record as to its action on phosphorus.

7. It unites to nearly its own weight of sulphur by fusion; the result is a lead coloured striated mass. It seems probable that the sulphuret contains two proportions of sulphur.

8. Very few experiments have been made upon the alloys of tellurium. It combines readily by fusion with potassium and sodium, producing heat and light; and forms with them difficultly fusible alloys, which, when thrown in water, produce purple solutions consisting of the alkalies united to telluretted hydrogen.

Tellurium has as yet been found in quantities too small to render it applicable to any of the purposes of the common arts.

XXI. *Cobalt or Cobaltum.*

1. Cobalt is procured from its ores, which are for the most part combinations of this substance with other metals; or of its oxide with arsenic or sulphuric acids. It is difficult to obtain the metal in a state of complete purity. The pure oxide may be procured by dissolving the ore known by the name of *arsenical cobalt* in nitric acid. The sulphur either remains on the surface, or is acidified with the arsenic, when both are to be separated by nitrate of lead. The excess of lead is to be removed by a little sulphuric acid, and the copper, if any, precipitated by a bar of iron. The clear solution is to be pre-

precipitated by carbonate of potassa, and the precipitate digested with solution of ammonia. The ammoniuret is to be evaporated till it does not tinge turmeric, and then acted on by solution of potassa: the precipitate, if any, is to be separated, and the clear fluid evaporated to dryness. The solid matter obtained by evaporation will afford *cobalt* when mixed with a little very fine charcoal powder, and exposed to an intense heat for about half an hour in a covered Hessian crucible.

Metallic cobalt was first procured by Brandt in 1733.

2. Cobalt is of a light gray colour, with a tint of red; its hardness appears to be greater than that of copper; its specific gravity is about 7.7. It is rather brittle; its fusing point is very high, not much inferior to that of iron, on the scale of Wedgwood as 130 to 158; it suffers little change from being exposed to air or water at common temperatures. Like iron, it is attracted by the magnet, and is capable of being rendered permanently magnetic. When fused on charcoal, and acted on by a stream of oxygen gas it burns brilliantly, throwing off bright sparks.

3. Cobalt combines with oxygen; when kept red hot for some time it becomes covered with a dark powder, and by being long exposed to air in a state of intense ignition, it is entirely oxidated, and in this process, according to Klaproth, 100 grains of the metal became 118 grains of oxide. This oxide though it appears black is in fact deep blue, and gives this tint to glass. It seems to be cobalt in its first state of oxygenation, and when dissolved by acids and thrown down by fixed alkalies, forms the basis of a *hydrate* of a bright blue colour. By heating the *hydrate gently in the air*, it gradually becomes black, loses its water, and absorbs oxygen. This black powder has the property of de-

composing muriatic acid; its excess of oxygen combines with the hydrogen, and the chlorine, is set free. From a rude experiment, I am inclined to conclude that the oxygen in the dark powder from cobalt is to that in the blue oxide as 3 to 2. If Klaproth's experiments be made the ground work of calculation, and the blue oxide be considered as a deutoxide, then the number representing cobalt will be 166, and the blue oxide will consist of 166 of cobalt and 30 of oxygen; and the black oxide of 166 and 45. M. Thenard states that there is an olive oxide of cobalt, produced by exposing the blue powder to air at common temperatures, which must be regarded as a hydrate. And Mr. Proust has stated that there is a red hydrated oxide of cobalt; it is probable that this last is a compound of the black oxide and water; and the substance supposed by M. Thenard to be an olive oxide, a mixture of the two hydrates.

4. Cobalt combines with chlorine: the compound may be obtained by introducing chlorine into an exhausted retort containing the metal in fine powder and gently heating it; a combustion takes place, but the results of this combustion have not yet been accurately examined.

5. Cobalt is not known to enter into combination with hydrogen, azote, carbon, or boron.

6. Cobalt combines with sulphur and phosphorus, but with considerable difficulty. The sulphuret is formed by acting on oxide of cobalt in a state of ignition by sulphur, and according to Proust, it consists of 71.5 parts cobalt and 28.5 of sulphur; which indicate nearly one proportion of metal 166, and two of sulphur 60. The *phosphoret* is made by dropping phosphorus upon ignited cobalt; it has not been minutely examined, nor its composition ascertained.

7. The action of the metals of the alkalies and earths on cobalt has not been examined.

8. There are no accurate experiments on the combinations of pure cobalt with the common metals. Lead unites with it, as Gmelin has shewn, and forms an alloy less malleable in proportion as it contains more cobalt.

9. Cobalt in its metallic state is not employed in the useful arts. In its state of combination with oxygen, it is used to give glass, porcelain, &c. a rich blue colour. One grain of the pure oxide will give a deep tint of blue to 240 grains of glass. The solution of the oxide of cobalt in muriatic acid, forms one of the most beautiful of the sympathetic inks. This solution when concentrated is pale rose-coloured in the cold, but becomes blue green when heated; letters or figures traced by it upon paper are invisible in the cold, but become blue green when held before the fire.

XXII. *Copper, Cuprum.*

1. The copper of commerce is procured from the native combinations of this metal with sulphur, oxygen, and certain acids, by roasting and fusion either alone or with lime and carbonaceous substances; it is not however quite pure. To obtain it in a state of purity—dissolve copper in strong muriatic acid, dilute the solution with water, and put into it a polished plate of iron, the copper will be precipitated in its metallic state: it should be washed in dilute muriatic acid and then with pure water.

2. Copper is of a red colour, it is a little harder than silver, its specific gravity is about 8.89. When rubbed it emits an unpleasant smell, and it has a disagreeable taste. It is very malleable, has considerable ductility,

and in tenacity is superior to all the metals except iron; it fuses at a low white heat. By exposure to the air copper becomes tarnished, and after some time is coated with a green crust, which consists of the metal in union with oxygen and carbonic acid. Copper is not affected by being kept in water, nor does it decompose this fluid at any temperature.—It burns with a red flame edged with green, when fused and acted upon by oxygen, when in thin leaves it inflames spontaneously in chlorine.

3. Two compounds are known, consisting of oxygen and copper. One is found native, and is the *ruby copper ore*. It occurs in octohedrons of a considerable lustre, its powder is dull orange red. According to Mr. Chenevix it contains 11·5 per cent. of oxygen; according to Mr. J. Davy about 11 per cent. It is soluble in strong muriatic acid; and when this solution is precipitated, a pale orange coloured powder falls down, which is this oxide united to water. The other oxide of copper is formed in the combustion of copper, or by heating the precipitate from a nitrous solution of copper by potassa to redness. It is a black powder, and appears, from various experiments, to contain about 20 per cent. of oxygen; when it is precipitated from acids by potassa it is in combination with water, and is then pale *blue*; and, as I have found, contains 10 per cent. of water. If the red oxide of copper be considered as a protoxide and the black a deutoxide, the number representing copper will be 120, and this number will be found to correspond accurately with that gained from the analysis of the other combinations of the metal. The number representing the protoxide is 135, and that representing the deutoxide 150, and that representing the blue hydrate 167.

4. As there are two combinations of copper and oxygen, so there are likewise two combinations of this metal with chlorine; both are produced at the same time, by the combustion of the metal in chlorine; one is a fixed easily fusible substance, like rosin in its exterior characters, the other is a yellowish sublimate. The first of these, as appears from the analysis of Mr. John Davy, consists of 36 chlorine and 64 copper; and the second of 53 chlorine and 47 copper; the first may be called *cuprane*, the second *cupraneæ*.* Cuprane may be formed likewise by heating strongly together a mixture of one part of copper filings, and two parts of corrosive sublimate; and it was in this way first produced by Boyle, who appears to have been its discoverer. Cuprane is converted into cupraneæ by being heated in chlorine. Cuprane may be regarded as consisting of one proportion of copper 120, and one of chlorine 67; and cupraneæ of one of copper 120, and two of chlorine 134. Cuprane is not soluble in water, but slowly becomes green by the action of the atmosphere; when introduced into the flame of a candle, it produces a most beautiful light possessing almost all the prismatic colours. *Cupraneæ* dissolves in water and gives it a greenish colour; it is decomposed by a strong heat, and converted into cuprane by the expulsion of oxygen.

5. Copper readily combines with sulphur, producing ignition when they are fused together; and they form together a substance more fusible than copper, brittle, and of a deep gray colour. This substance is likewise found native, and according to the analysis of Mr. Chenevix, contains about 19 per cent. of sulphur. The artificial *sulphuret* in some synthetical experiments which I made upon it, appeared to contain from 21 to

* [Chloride and perchloride.]

19 per cent. It may therefore be regarded as composed of one proportion of copper 120, and one of sulphur 30. It is probable that a super-sulphuret of copper may exist: some of the golden coloured copper pyrites, which contain as much as 41 per cent. of copper, afford from 35 to 45 of sulphur: but they likewise contain iron. No *supersulphuret* has as yet been made artificially.

6. Copper combines with phosphorus, by fusion. The *phosphoret* is of a white colour, and very brittle, its specific gravity is 7.1220. It was first formed by Margraff. According to Pelletier, it contains 20 per cent. of phosphorus; and allowing this estimation, it must be composed of 2 proportions of metal, and 3 of phosphorus, 240 and 60.

7. Copper has not yet been combined with hydrogen, azote, carbon, or boron.

8. It unites to the fixed alkaline metals, and to all the common metals that have been described. Some of its alloys with the common metals are well known. Copper is rendered yellowish white by alloy with a small quantity of manganesum. United to zinc, copper produces brass, Dutch gold, Rupert's metal, and pinchbeck: from a third to a twelfth of zinc is used; the paler the alloy required, the larger the quantity of zinc.

Copper, with a fourth of its weight of lead, forms pot-metal. Copper alloyed with from $\frac{1}{12}$ to $\frac{1}{5}$ of tin forms the different species of bronze and bell-metal. The best composition for the mirrors of reflecting telescopes is a combination of 32 parts copper, 15 parts tin, 1 part brass, 1 silver, and 1 of arsenic. Tutenag, according to Keir, is a white alloy of copper, zinc, and iron.

9. The account of the alloys of copper proves its im-

portance in the arts. It is used unalloyed likewise for various important purposes, such as sheathing of ships, forming vessels when united with other metals, for culinary purposes, &c. In early ages the alloys of copper formed the principal arms offensive and defensive. I have examined an ancient Attican helmet, which consisted of an alloy of copper and tin. The swords and spear-heads of the early inhabitants of Greece and Italy seem to have been composed of the same materials.

All the saline combinations of copper are poisonous.

XXIII. *Nickel or Nickolum.*

Nickel was discovered by Cronstedt in 1751, and examined in its pure state by Bergman, in 1775. Nickel exists in an ore called kupfer-nickel combined chiefly with sulphur, or in nickel ochre, in which it is united to oxygen. The metal may be obtained from these ores by roasting and ignition with charcoal; but in this case it is far from being pure. *Pure nickel*, or at least nickel free from any other metallic substance, may be procured by nearly the same process as cobalt. The precipitate from the ammoniacal solution by solution of potassa, contains the nickel, and this precipitate must be intensely heated with charcoal powder.

2. Nickel is of a white colour, and possesses considerable lustre; its hardness is little inferior to that of iron: its specific gravity is about 8.38, but when forged it increases to 8.82. It is ductile and malleable. It requires a stronger heat for its fusion than iron—like that metal, it is attracted by the magnet. When in fusion it burns like iron under a stream of oxygen gas.

3. Nickel when intensely ignited, exposed to air, becomes a dark brown powder, which is still attractable

by the magnet. Its solution in nitric acid decomposed by potassa affords a pale grass green hydrated oxide, which contains more than a fourth of its weight of water, and which when heated to dull redness becomes an oxide of a pale ash gray colour, and which, according to Tupputi, is composed of 21·2 parts of oxygen, and 78·8 of metal: this oxide by stronger ignition becomes darker coloured, but when pure cannot be reduced to the metallic state by heat alone. Another oxide of nickel has been described by M. Thenard containing more oxygen. It may be procured by acting on the hydrate of nickel by the salts called hyperoxymuriates; it is of a black colour; its composition has not been accurately ascertained. From the nature of the compounds of nickel and sulphur, to be discussed immediately, it seems probable that the gray oxide is a deutoxide; and if this be allowed, the number representing nickel will be 111, and the gray oxide will be represented by two proportions of oxygen 30 added to one proportion of metal.

4. Nickel, when strongly heated in chlorine, smokes and produces an olive-coloured substance; but the composition of this substance has not been ascertained, nor its properties examined. When muriate of nickel is decomposed by heat, some white brilliant scales are formed, which, as appears from experiments made by Mr. E. Davy, consist of nickel and chlorine; but their composition has not been exactly ascertained.

5. Sulphur combines with nickel by fusion, and forms a bright gray sulphuret, possessing metallic lustre. From experiments made on this *sulphuret* by Mr. E. Davy, it appears to contain about 34 per cent. of sulphur, which gives proportions corresponding nearly to one proportion of metal 111, and two of sulphur 60,

and harmonizes with the supposition that the olive oxide is a deutoxide. The same enquirer states that there is a supersulphuret, which may be formed by heating the gray oxide with sulphur, and which contains about 56·5 of nickel to 43·5 of sulphur, and which agrees nearly with one proportion of metal to three of sulphur.

6. *Phosphuret of nickel* may be formed by causing phosphorus in vapour to act on metallic nickel in ignition; the phosphuret is almost black, and has metallic lustre. Its composition has not as yet been accurately determined.

7. Some specimens of nickel afford carbon during their solution in acids; but no definite compound of carbon and the metal has been obtained. Nickel has not yet been combined with hydrogen, azote, or boron, nor have any accurate experiments been made on its action on the metals of the fixed alkalies.

8. It forms alloys with some of the common metals; but few of these compounds have been examined with attention; with tin it produces a white brittle compound. Its alloy with copper is less ductile than pure copper, and is slightly magnetic. Its combination with iron is the most interesting of these compounds; these metals seem to unite in all proportions; the colour of the alloy approaches nearer to that of silver in proportion as the nickel prevails, and the iron retains its malleability. In all the meteoric stones that have been examined it is remarkable that the iron is alloyed by from 1·5 to 17 per cent. of nickel. The masses of iron found in Siberia and South America contain nickel, and there is the strongest probability that they are likewise of meteoric origin. The alloy of iron and nickel is much less liable to rust than common iron, and it is

likely may be advantageously applied in the useful arts. The oxide of nickel is employed to give colours to enamels and porcelain; in different mixtures it produces brown red, and grass green tints.

XXIV. *Uranium.*

1. *Uranium* was discovered by Klaproth in 1789. It may be procured from the ores called Pechblende, and Uranochre, by the following process. Let the ores be boiled in moderately diluted nitric acid, and solution of sulphate of soda added, and the precipitate, if any, separated. The clear solution is to be acted on by solution of potassa, the precipitate digested in ammonia, and the residuum heated with strong nitric acid, and evaporated to dryness. The clear solution obtained from the mass by pure water when precipitated by solution of potassa will afford a yellow powder, and this made into a ball with wax, and intensely ignited in a crucible of charcoal, affords metallic *uranium*.

2. *Uranium* is of an iron gray colour, and has considerable lustre; it is hard and brittle. Its specific gravity, according to Klaproth, is 8.1; its fusing point is higher than that of manganesum. It undergoes no change by exposure to air, but when heated strongly, burns, combines with oxygen, and assumes a blackish colour.

3. Two compounds of uranium and oxygen have been examined by Klaproth; the precipitate thrown down from the solution of uranium in nitric acid when heated to dull redness is still yellow, and this treated with oil, and incinerated slightly, so as to burn off the oil, becomes a *black oxide*. It appears probable from Bucholz's experiments, that the oxygen in the black

oxide is to that in the yellow as 1 to 3, and that the yellow oxide contains about 80 of metal and 20 of oxygen. I found that 8 parts of potassa precipitated 8·2 parts of yellow oxide of uranium from the saturated nitrous solution; and from this experiment, if potassa and the yellow oxide of uranium be supposed analogous in composition, 76·8 will be the number for the metal, and the black oxide must be a compound of three proportions of metal with one of oxygen. The substance that has been called the native oxide, and which is crystallized in quadrangular plates, is, I find, a *hydrated oxide*. Bucholz supposes that there are several different oxides of uranium; but he founds his opinion upon the different colours of precipitates, which may be mixtures of hydrates of the two oxides.

4. No experiments have as yet been made on the action of uranium on chlorine, hydrogen, azote, boron, the metals of the fixed alkalies, or the common metals.

5. A sulphuret of uranium may be procured by igniting the oxide with sulphur. It is a black heavy powder: its composition has not been ascertained.

6. Uranium has been hitherto found in quantities too small to render it applicable to the purposes of the arts. Its oxides give bright colours to glass, which, according to the proportions are brown, apple green, or emerald green.

XXV. *Osmium*.

1. This metallic substance was discovered by Mr. Tennant in 1804. It may be obtained from crude platina, by dissolving all the soluble parts in aqua regia, and distilling the black powder that remains with nitre, at a heat under redness; a sublimate rises, which is

soluble in water. When mercury is shaken with the solution, an amalgam is formed, and by distilling off the mercury, pure osmium is obtained.

2. This metal has been procured only in very minute quantities; it is of a dark blue colour, has not been fused, nor does it undergo any change at the most intense heat, unless in contact with air, when it is converted into a volatile oxide.

3. The composition of the *oxide of osmium* has not been ascertained;* it is a solid semi-transparent substance, having a sweet taste and a strong smell: it is soluble in water, combines with potassa, and makes with it an orange solution in water. It tinges the skin of a dark colour; and produces a purple with solution of galls.

4. No combinations of osmium with any of the undecomposed substances described in the foregoing sections except that with oxygen have been examined. The metal is not soluble in any of the acids; but when fused with the hydrate of potassa becomes oxidated and combines with the potassa. It is a metal very easily reduced, being precipitated from the aqueous solution of the oxide by ether or alcohol.

XXVI. *Tungsten or Tungstenum.*

1. Tungstenum is obtained from a mineral known by the name of *wolfram*; it contains the oxides of tungsten, iron, and manganese, with earthy matter. To procure the metal pure, boil finely pulverised wolfram in strong muriatic acid for some time; separate the solution; the

* [According to Berzelius, five different oxides of this metal may be obtained, and four different chlorides and sulphurets; he has determined the proportion of oxygen, chlorine, and sulphur in each.]

residuum contains a yellow powder; it is to be washed, dissolved in ammonia, evaporated to dryness, then mixed with a little fine charcoal powder, and exposed to a very intense heat for about 20 minutes in a covered hessian crucible. Small grains of pure tungstenum will be found at the bottom of the crucible.

Tungstenum in its metallic form was first procured by Messrs. D'Elhuyars, in 1782.

2. *Tungstenum* is of a grayish white colour, and has considerable lustre. It is hard, and rather brittle. Its specific gravity is about 17.3. It requires the strongest heat of a forge for its fusion. It is scarcely affected by exposure to air at common temperatures. At a temperature below redness its surface exhibits iridescent colours like iron.

3. Tungstenum combines with oxygen. When the metal in fine powder is heated to redness, it soon acquires a yellow colour, and is gradually converted into a *yellow oxide*, which is not soluble in water. Its specific gravity is 612, water being 100. It is very difficultly fusible. According to Messrs. D'Elhuyars, 100 grains of metal by calcination form 124 grains of yellow oxide. Supposing the oxide a deutoxide, this result would give the number representing tungstenum as 125, and the number representing the yellow oxide as 155. Taking Klaproth's analysis of the combination of tungstic acid and lime,* as the basis of calculation, and supposing double proportions of the oxide to one proportion of lime, the number representing the yellow oxide will be 124. From this it is probable that in D'Elhuyars' experiments the metal was not entirely converted into oxide: and that the number is about 94; but new experiments are wanting to elucidate this point. When

* 17 of lime, 77 of tungsten oxide.

the yellow oxide of tungstenum is digested with solution of tin in muriatic acid, it becomes blue from losing oxygen. It is probable that in this state it is a protoxide, but no accurate researches have been made on this blue substance.

4. Tungstenum, I have found, burns with a deep red light when heated in chlorine, and forms an orange-coloured volatile substance, which affords the yellow oxide of tungstenum, and muriatic acid, when decomposed by water. I have made no experiments, nor are any, I believe, on record, on the composition of *tungstenane*.*

5. Sulphur and phosphorus are both capable of being combined with tungstenum by being made to act upon it in a state of ignition; but the properties of the *sulphuret* and *phosphuret* have not been examined accurately.

6. This substance has not been combined with hydrogen, azote, carbon, boron, or the metals of the fixed alkalies. From the experiments of Messrs. D'Elhuyars, it appears to unite with most of the common metals; but its alloys have been only rudely examined.

7. Tungstenum and its oxides have as yet been applied to no uses: it was stated by Guyton de Morveau that the yellow oxide formed a mordant useful in dyeing; that the red juices of fruits were fixed by it, so as to make permanent and beautiful lakes. The dyers who have tried the experiments in this country have not, however, given a favourable report of the results.

XXVII. *Titanium*.

1. Titanium is obtained from a mineral long known

* [Chloride of tungsten.]

by the name of *red schorl*, or titanite. The mineral in powder is to be fused with five or six times its weight of subcarbonate of potassa; the mass is to be fully exposed to the action of water, and the solid matter remaining digested and boiled with muriatic acid. The white powder not dissolved, when mixed with oil, and intensely heated in a crucible with charcoal, affords titanium. The oxide of titanium was discovered by Mr. Gregor, in 1781, in an ore found in the valley of Menachan in Cornwall; but metallic titanium was not produced till 1796, by Vauquelin and Hecht.

2. Little is known concerning the physical and chemical properties of *titanium*; it has only been procured in minute quantities, and in an imperfectly reduced state. Its colour resembles that of copper. It has much lustre. It is brittle. It tarnishes by exposure to air; and requires the most intense heat of a forge even for its imperfect fusion.

3. Titanium combines with oxygen when it is exposed to heat in the atmosphere, and acquires a blue colour. The *red oxide*, which contains a larger proportion of oxygen, is found in the mineral kingdom. There is a powder of a white colour procured by fusing the red oxide with potash. This has been supposed to be a peroxide, but is probably an hydrated oxide; no precise experiments have been made on the composition of these bodies. This white powder is insoluble in acid and alkaline solutions: and becomes yellow by being heated.

4. The combination of titanium with chlorine has not yet been made.*

* [It has since been formed—it is a colourless volatile liquid;—a sulphuret also has been obtained.]

5. Titanium is not known to combine with hydrogen, azote, carbon, or boron.

6. Titanium has not yet been combined with sulphur. It enters into union with phosphorus. The phosphuret is brittle, and has metallic lustre. Its composition has not been determined.

7. The agency of the metals of the alkalies on titanium has not been examined. It has not been combined with any of the common metals except iron; and this alloy is not characterised by any remarkable properties.

8. Titanium has not been employed in the arts except for one purpose; its oxide has been used at Sevres in the manufacture of porcelain, to impart a brown colour.

XXVIII. *Columbium.*

1. Columbium exists in an ore brought from North America, of a black colour; and likewise in two substances found in Sweden, called tantalite and ittrotantalite. The metal may be procured by igniting the ores with hydrated fixed alkali; and saturating the alkali with nitric acid, a white powder falls down. This powder was first obtained by Mr. Hatchett from the American specimen, in 1802: and soon after the same substance was procured by Ekeberg from the Swedish mineral, and considered by him as a new substance. Dr. Wollaston, in 1810, demonstrated the identity of the two bodies.

2. The white powder combines with alkalies and metallic oxides, and reddens litmus paper. Hence Mr. Hatchett named it *columbic acid*. Attempts were made to reduce it by ignition with charcoal, in the same

manner as the oxide of titanium, but without success; it became black, but did not acquire the metallic lustre. By passing potassium in vapour through the white powder, heated to redness, I found that the potassium became converted into potassa, and a dark coloured brilliant powder, like plumbago, was produced. This is probably the metallic basis of the substance, or pure columbium.

3. No experiments have been made upon the combinations of this substance.* The white powder is soluble in boiling sulphuric acid; and it is precipitated from its solution of an olive colour by triple prussiate of potash, and of a bright orange by solution of galls.

XXIX. *Cerium.*

1. There is a mineral found at Ridderhytta in Sweden, very like tungsten, of a reddish colour, and which has been called cerite. From this substance, Hisinger and Berzelius, in 1804, extracted a brown powder, having the characters of a metallic oxide, and which they named *oxide of cerium*. To procure this powder, the ore is digested with solution of nitro-muriatic acid; and the solution obtained evaporated to dryness, and heated with a little muriatic acid; the solution so procured is to be precipitated by solution of ammonia; the precipitate well washed is to be redissolved in muriatic acid, and oxalic acid is to be added; a white insoluble oxalate of cerium is thus formed, which when heated to redness affords the *brown oxide of this metal*.

2. *Cerium* had not been obtained in the metallic form till I succeeded in reducing some oxide sent me

* [It has since been combined with chlorine, fluorine, and with sulphur.]

by M. Berzelius, by means of potassium; potassa was formed, and a deep gray metallic powder, which became brown by oxidation.

3. When the brown oxide of cerium is digested in the mineral acids, it becomes dissolved; and is thrown down from those solutions by alkalies, as a white powder: it is supposed that this powder contains less oxygen than the brown oxide; but it is probably a *hydrate*. As yet, no experiments have been made either on the composition of the brown oxide, or the white powder.

4. No researches have as yet been made on the combinations of cerium with the other undecomposed bodies.* The solutions of cerium are not precipitated by solutions of galls; they give a white precipitate with the triple prussiate of potassa.

XXX. *Palladium*.

1. *Palladium* was discovered by Dr. Wollaston, in 1803. It exists in the ores of platinum, both those from Peru and the Brazils. It may be procured by dissolving crude platina in aqua regia, and precipitating the saturated solution by solution of prussiate of mercury. The precipitate, washed, dried, and exposed to a strong heat, is converted into palladium.

2. The colour of palladium is white, resembling that of platinum. Its hardness is rather greater than that of bar iron. Its specific gravity varies from 11.3 to 11.8. It is very malleable, but has little ductility. It fuses at a high temperature; but the precise point has not been determined. It is not affected by air or

* [Besides an additional oxide, two chlorides, and a sulphuret of cerium have since been formed.]

water at common temperatures. When it is heated strongly, its surface acquires a blue colour.

3. Palladium combines with oxygen and chlorine by heat; but neither its oxides nor its combinations with chlorine have as yet been examined. Dissolved in nitric acid it forms a beautiful red solution, from which the alkalies throw down an orange coloured powder, which probably is a *hydrated oxide*.*

4. Palladium readily combines with sulphur when they are heated together in a glass tube. The *sulphuret* is rather paler than the metal, and very brittle: in an experiment that I made, 6·7 grains of palladium gained 1·5 grains, by being converted into the sulphuret; and supposing the sulphuret to consist of one proportion of metal, and one of sulphur, the number representing palladium will be 134.

5. Palladium seems to have no action on hydrogen, azote, or carbon. Its relations of attraction to boron, and the metals of the fixed alkalies, have not yet been examined. It forms alloys with most of the common metals; but the properties of these compounds have not been examined with attention.

Palladium has not as yet been found in sufficient quantities to be applied to the purposes of the arts.

XXXI. *Iridium*.

1. Iridium was discovered in 1803, by Mr. Tennant; before Mr. Tennant had published his experiments, it was likewise discovered by M. Descotils.

Iridium exists in minute quantity in the crude ore of

* [It combines with two different proportions of oxygen and chlorine: from the analytical results of Berzelius the number representing this metal appears to be about 99.]

platina. To obtain it, the black powder (remaining after the solution of the ore of platina in nitro-muriatic acid) is to be mixed with about five times its weight of pure soda, and heated to redness in a silver crucible for about thirty minutes. The dry mass is to be dissolved in diluted muriatic acid, and the undissolved residuum is to be alternately treated with alkali and acid; by which means it will be all taken up. The solution, containing an excess of muriatic acid, is to be evaporated to dryness, redissolved in pure water, and slowly evaporated, so long as any octohedral crystals form. These crystals are muriate of iridium, and are reduced to the metallic state by exposing them for a short time to an elevated temperature in a platinum crucible.

2. Little is known concerning the properties of this metal. Iridium is of a white colour. It is brittle: and requires for its fusion a most intense heat: it is probable that its specific gravity is higher than that of platinum. It is not acted upon by oxygen even when heated to whiteness. From its relations to muriatic acid, which dissolves it, it seems that it is capable of uniting to chlorine.*

3. Iridium has not been combined with hydrogen, azote, sulphur, phosphorus, carbon, or boron, or the metals of the alkalies. It unites to lead; and forms a malleable alloy with copper. Dr. Wollaston has found amongst the grains of crude platina small white particles of specific gravity 19·25; which consist of iridium alloyed with osmium, and no other metallic substance. The osmium may be oxidated by the water in hydrate of potassa, and united with the potassa; and the iridium

* [Four oxides of this metal have been obtained and examined by Berzelius, and as many chlorides, and it has been combined with sulphur and carbon.]

combined with chlorine by treatment with muriatic acid, and thus dissolved.

XXXII. *Rhodium.*

1. Rhodium was obtained by Dr. Wollaston, in 1804, from the ore of platina, by the following process. The ore is dissolved in dilute aqua regia; a solution of sal-ammoniac is added; the clear liquor, separated from the precipitate, is acted on by a rod of zinc. By the zinc, a black powder is thrown down, which is washed with very diluted nitric acid. This black powder is redissolved in dilute aqua regia; to this solution some common salt is added; the whole is then evaporated to dryness, and washed by alcohol, till it has dissolved all the soluble matter; there remains behind a deep red substance, which, when dissolved in water, and acted on by a rod of zinc, affords a metallic powder, which, intensely ignited with borax, gives a metallic button of rhodium.

2. The specific gravity of rhodium exceeds 11. Its colour approaches to that of silver, with a tint of yellow. It is not acted upon by nitric or sulphuric acids. It is not known whether it combines with oxygen; but solution of potassa throws down a yellow coloured powder from the red crystals obtained by dissolving in water the powder left after the washing by alcohol, and evaporating, so as to permit crystallization. From the action of nitro-muriatic acid on rhodium, it is probable that it combines with chlorine.*

3. Rhodium unites with sulphur, and is rendered easily fusible by it. It likewise combines with lead,

* [Two oxides of this metal have since been formed, and two chlorides.]

copper, and bismuth; and its alloys are easily soluble in nitro-muriatic acid.

XXXIII. *Mercury, or Mercurium.*

1. Mercury has been known from the earliest ages of the world. It is found native in the mines of Idria, Spain, and Peru; and likewise combined with sulphur in cinnabar, from which it is separated by distillation with quicklime.

2. Mercury is of the specific gravity 13.56. Its colour is a brilliant white. It is fluid at the common temperature of the air; and becomes a solid at 39° below 0 of Fahrenheit's scale: at about 660° it boils. It is capable of being burnt when the Voltaic flame is made to act upon it, and it produces a brilliant greenish light; its capacity for heat has been referred to page 77.

3. There are two well known compounds of mercury and oxygen, one black and the other red; the first may be made by keeping mercury long in agitation, as, for instance, by fastening it in a bottle containing much air, to a mill wheel.* The second is produced by keeping it heated in the atmosphere for a long while, nearly at its boiling point. The black oxide of mercury may be likewise made by acting on calomel by solution of potassa; and the red oxide by acting on corrosive sublimate by the same substance, and gently heating the precipitates after they are washed. The black oxide changes when gently heated in the air, and becomes the red oxide. Both the black and the red oxides of mercury are decomposed by a strong heat; the mer-

* [The black powder thus formed is commonly a mixture of oxide of mercury and of the metal in a very finely divided state.]

cury revived and oxygen liberated: from experiments on their decomposition I ascertained that the quantity of mercury being the same, the red oxide contained exactly twice as much oxygen as the black; and my results give the number representing mercury about 380, the protoxide or black oxide as composed of 380 mercury and 15 oxygen, and the red oxide or deutoxide as constituted by 380 metal and 30 oxygen. It has not yet been ascertained whether these oxides can be combined with water, so as to form hydrates.

4. Mercury combines with chlorine: when the metal is heated in the gas, it burns with a pale red flame, and the substance called *corrosive sublimate* is formed. When corrosive sublimate is long rubbed with mercury till they are incorporated together, and the mass sublimed, the substance called calomel is formed. I have found by a minute analysis, that the quantity of chlorine in *calomel* is to that in corrosive sublimate as 1 to 2, the quantity of mercury being the same in both. Calomel is tasteless, corrosive sublimate has an acrid burning taste; calomel is insoluble, corrosive sublimate soluble in water. Calomel, according to my analysis, consists of one proportion of mercury 380, and one proportion of chlorine 67. Corrosive sublimate of 380 metal, and 134 chlorine. The names *mercurane* and *mercurana*, which may be adopted to signify the relations of their composition, are too similar to each other to be safely used as familiar appellations for the two substances, as corrosive sublimate is a powerful poison, calomel an excellent medicine.*

5. Sulphur and mercury readily combine by fusion; three parts of mercury and one of sulphur melted together, heated to redness, and then sublimed out of

* [The same remark applies to chloride and perchloride, or to protochloride and bichloride.]

the contact of air, afford a cake of a fine red colour, called *cinnabar*, and known in commerce under the name of *vermilion*. It seems from the experiments made on this substance, that it contains one proportion of mercury, and two of sulphur. When sulphur and mercury are heated strongly together, but not to sublimation, a black mass is obtained, which has been called *Ethiops mineral*.* It is probable that this substance contains a larger quantity of sulphur than cinnabar, but its composition has never been ascertained, and it is always converted into cinnabar by sublimation. The specific gravity of cinnabar is about 10, that of *Ethiops* is less: both these substances are easily decomposed by any metal having a stronger affinity for sulphur; when heated with iron filings, for instance, the sulphur combines with the iron, the mercury rises in vapour and condenses.

6. I have made a combination of phosphorus and mercury, by strongly heating together phosphorus and calomel. It is of a chocolate colour, and not fusible at the boiling point of mercury. I have made no experiments on its composition.

7. No combinations have as yet been effected of mercury with hydrogen, azote, charcoal, or boron.

8. Mercury unites readily with potassium and sodium, and forms solid alloys; the combination is attended with much heat: 1 part of potassium renders solid at common temperatures 70 parts of mercury. These *amalgams*, for so the metallic combinations of mercury are called, are of the colour of silver; the mercury rises

* [*Ethiops mineral*, according to the experiments of Mr. Brande, is a mixture of sulphur and bisulphuret of mercury.—A black protosulphuret it would appear may be formed by the action of sulphuretted hydrogen on calomel suspended in water.]

from them at a heat below redness. The alkaline metals are rapidly separated from the mercury by the oxygen of air or water.

9. Mercury combines with most of the common metals described in the preceding pages, and forms *amalgams* with them. It unites most readily with the easily fusible metals, but few researches have been made on its union with the difficultly fusible metals, as its volatility renders it difficult to make the experiments under favourable circumstances.

10. Mercury is a very important and useful metal. It is employed for extracting gold and silver from their ores. It is used in amalgamation with tin for covering mirrors. The sulphuret forms the most perfect red pigment as yet discovered. Its oxides and combinations with chlorine constitute some of the most important substances employed in pharmacy.

XXXIV. *Silver, or Argentum.*

1. Silver is found native, or is procured from ores, which are principally combinations of silver with other metals or with sulphur, but the silver of commerce is not pure. To obtain it pure: the metal must be dissolved in nitric acid, and the solution mixed with solution of common salt until no further precipitate takes place. The precipitate must be washed and ignited strongly with about three times its weight of subcarbonate of potassa, mixed with a little charcoal in powder, for half an hour; a button of pure silver will be procured.

2. Silver is of a brilliant white colour. It has no taste or smell; it has great lustre. Its hardness is inferior to that of copper: its specific gravity is about

10·40, and it is slightly increased by being hammered. It yields to none of the metals except gold in malleability. It is very ductile, and may be readily drawn out into extremely fine wire. Its tenacity is considerable. A wire of 0·078 of an inch in diameter will support 187·13 lbs. weight. Its fusing point is about 1000° Fahrenheit. It tarnishes slowly in the air; and this tarnish is owing to the presence of fumes containing sulphur.

3. Silver enters into combination with oxygen; it absorbs this gaseous principle when kept intensely ignited in an open vessel.* It burns with a fine green flame, and is converted into an oxide when acted upon by a powerful Voltaic instrument.

The olive oxide of silver is likewise obtained by dissolving the metal in nitrous acid, precipitating by aqueous solution of baryta, and heating the precipitate to dull redness. From my experiments I conclude that 100 of silver absorb about 7·3 parts of oxygen to become the brown oxide; and supposing this oxide to be composed of one proportion of metal and one of oxygen, the number representing silver will be 205. No other oxide of silver except the brown is certainly known.

4. Silver combines with chlorine when the metal is heated in contact with the gas. The compound, which may be called *argentane*, has been long known by the name of *hornsilver*. It is a whitish semi-transparent substance, cuts like horn, is fusible at a red heat, and is insoluble in water. It contains about 24·5 per cent. of chlorine, and may be considered as consisting of one proportion of silver 205, and one of chlorine 67.

* [M. Lucas has determined the curious fact, that the oxygen absorbed by silver in fusion, is disengaged on cooling in the act of becoming solid.]

5. Silver is not known to combine with hydrogen, azote, carbon, or boron.

6. Silver and sulphur combine. This combination is effected, according to M. Proust, when silver is exposed for a considerable time to the atmosphere; it is readily formed by heating to redness thin plates of silver and sulphur. It is of a black colour, is brittle, and has the metallic lustre. According to the experiments of Wenzel, 100 parts of silver by fusion combines with 14·7 parts of sulphur. The sulphuret of silver may therefore be regarded as constituted by one proportion of silver 205, and one of sulphur 30.

7. Silver combines with phosphorus. This compound may be made, according to M. Pelletier, by heating to redness a mixture of silver, phosphoric glass, and charcoal powder. It is a white brittle substance; its composition has not been determined with precision.

8. The action of the metals of the alkalies and earths on silver has not been examined. Silver forms alloys with most of the other metals, but the greater number of them have not been examined with much attention, or been applied to the purposes of the arts. The alloy of silver and copper is employed in coins. It is harder than pure silver, and better adapted to receive a fine impression.

9. Silver is employed for a great variety of purposes in the useful and ornamental arts. It is largely used for silvering copper, brass, and sometimes iron. In the common form in which it is applied, it is alloyed with $\frac{1}{12}$ of copper, which gives to it hardness, without impairing its colour or its lustre.

XXXV. *Gold, or Aurum.*

1. Gold is found native, alloyed with copper or silver. To obtain it in a state of purity: gold is dissolved in nitro-muriatic acid, the silver will remain an insoluble muriate, and must be separated; to the clear solution a solution of green sulphate of iron must be added; the gold will be precipitated in the state of a fine powder, and after being well washed in diluted muriatic acid, and then in distilled water, may be fused into a mass.

2. Gold is of a fine light yellow colour; its hardness is scarcely superior to that of tin. Its specific gravity is about 19·277, and it is somewhat increased by hammering. In malleability and ductility it is superior to all the other metals. It has a considerable degree of tenacity; a wire of 0·078 inch in diameter will support a weight of 150 lbs. It fuses at about 1300 Fahrenheit. It is not altered by exposure to air or water.

3. There are no accurate experiments recorded on the combinations of gold with oxygen. A purple oxide is formed when gold leaf is burnt by electricity, or gold wire by the Voltaic battery, but its composition has not been ascertained. No dependence can be put on the statements of chemists relative to pure oxides of gold, said to be obtained by treating solution of gold, with potassa, lime, and other substances, for in these instances, as far as my experience has gone, triple compounds always appear to be formed.

4. Gold combines with chlorine when the metal, in a minute state of division, is heated in chlorine, or when the nitro-muriate of gold is partially decomposed by heat, treated with muriatic acid and evaporated to

dryness. It is a brown substance, is very deliquescent, and readily decomposes the water in the atmosphere, forming a muriate of gold. It has not been examined with precision.

5. There are no known combinations of gold with hydrogen, azote, carbon or boron.

6. There has been no distinct combination made of gold and sulphur.

7. Gold combines with phosphorus: this compound has been recently made in the laboratory of the Royal Institution by Mr. E. Davy, by heating gold, in a minute state of division, with phosphorus in an exhausted tube. It is of a gray colour, and has the metallic lustre. It is readily decomposed by the heat of a spirit lamp; and contains about 14 per cent. of phosphorus.

8. The metals of the alkalies combine with gold; but the alloys have not been minutely examined.

9. Gold forms alloys with the other metals; many of them are brittle, as those of bismuth, antimony, and lead. Others are malleable, as those of silver, copper, and platina. The alloy of gold and copper is employed in coin.

The applications of gold to the useful and ornamental arts, &c. are too well known to need particular detail. The purple oxide of gold is employed for colouring glass and porcelain.

XXXVI. *Platinum.*

1. The ores of *platinum* are very rare; they have been found only in South America and in Spain.* The ores from South America consist of small roundish flat-

* [It has since been discovered in the Uralian Mountains.]

tened grains. The ore found in Spain is in a vein principally consisting of silver. The only places in South America in which grains of the ore of platina have been discovered are at Choco in Peru, Santa Fe near Carthagena, and a district in the Brazils.

Platinum is procured from the South American ore by dissolving it in aqua regia, and dropping into it a solution of sal-ammoniac; a yellow powder falls down, which must be redissolved in nitromuriatic acid, and again precipitated by sal-ammoniac; and after this second process, when ignited to whiteness, it is pure platinum. The particles may be made to unite into one mass by hammering them in a state of ignition.

2. Platinum was first described as a peculiar metal by Dr. Lewis, in 1754.

Platinum is of a white colour, but much less brilliant than silver; it is not quite so hard as malleable iron; its specific gravity after being hammered is 21·3, that of water being 1. It is very ductile, and malleable; may be easily drawn into wires about the $\frac{1}{2000}$ of an inch in diameter,* and hammered into very thin plates: its tenacity is such, that a wire 0·078 of an inch in diameter is capable of supporting a weight of 274·31 lbs. avoirdupois without breaking. It is not fusible by the heat of a forge; and requires either the intense heat of the concentrated solar rays, of Voltaic electricity, or of a flame produced by the agency of oxygen gas.

3. Platinum combines with oxygen only with great difficulty. When intensely ignited by Voltaic electricity, it fuses, and throws off sparks, and a fume rises from it, which is probably the oxide of platinum.

When solutions of platinum are precipitated by alka-

* [By Dr. Wollaston's method, vide Phil. Trans. for 1813. He obtained wire of the diameter of $\frac{1}{30000}$.]

lies, or alkaline earths, the precipitate always appears to be a compound of platinum, oxygen, and the earth or alkali employed; yet Mr. Chenevix has stated, that by dissolving the precipitate from the nitromuriatic solution by lime water in nitric acid, and driving off the acid by heat, a brown powder is formed, which is an oxide of platinum, and which contains 13 per cent. of oxygen. The same ingenious chemist states that there is another oxide of platinum of a green colour, made by heating the brown oxide, and which he believes contains 7 per cent. of oxygen. I have seen several experiments made by Mr. E. Davy, in which no precipitate was produced by the action of lime water on the nitromuriatic solution: the lime water was used in various proportions, but without success.

When an alloy of potassium and platinum is heated, exposed to air, both metals burn, and a yellow powder is formed, which gives off oxygen gas by ignition; but this powder, after being long washed, reddens turmeric, so that it is not pure oxide of platinum.*

4. A bright brown powder may be obtained by evaporating to dryness the nitromuriatic solution of platinum. This powder, when heated to whiteness, is resolved into platinum and chlorine, and the chlorine gas may be collected in a proper apparatus. From some experiments made on this powder, at my request, by Mr. E. Davy, it appears to contain about 24 per cent. of chlorine; but this estimate can be considered only as an approximation, for there are many difficulties in gaining accurate results on a substance so easily decomposed.

5. Sulphur combines with platinum when they are

* [Two oxides of this metal have been procured, and the same number of chlorides and of sulphurets.]

heated together in exhausted tubes. The *sulphuret* is an infusible black powder, decomposable by a white heat. According to Mr. E. Davy, who first made it in the laboratory of the Royal Institution, it contains about 16 per cent. of sulphur. He supposes that there is another combination of sulphur and platina, which may be made by heating the precipitate from the nitro-muriatic solution by sal-ammoniac and sulphur together, and which contains 28 per cent. of sulphur.

6. Phosphorus and platinum combine with great energy, when the phosphorus is made to act in vapour, in exhausted tubes, on platina, heated to dull redness; the combination is so violent that the mass becomes vividly ignited. The *phosphuret* of platina is an infusible bluish gray powder with little lustre. According to Mr. E. Davy, it contains more than 17 per cent. of phosphorus. He believes that there is a *superphosphuret* of platina containing 30 per cent. of phosphorus, made by heating the yellow powder procured by sal-ammoniac with phosphorus; but new experiments are wanting on this result, as well as on the results of the action of sulphur on platinum. The quantities do not correspond with the theory of definite proportions; and as neither the metal nor the compounds made can be fused under the circumstances of the experiment, it is not possible to say that the combination is perfect; and as all such combinations are decomposable by a strong heat, part of the compound first formed may be decomposed before other parts of the mass enter into union.

7. From the experiments of M. Descotils, it is probable that platinum is capable of combining with boron; it has never been united to hydrogen, azote, or carbon.

8. Platinum readily unites to potassium and sodium;

their combination takes place with ignition, and a bright brittle mass is obtained, from which the alkaline metals are readily separated by the action of air or water. Platinum combines with most of the other metals; but the properties of its alloys have been very little studied. To the fusible metals it communicates difficult fusibility. It amalgamates with mercury when heated with that metal in a finely divided state. It combines with gold, and renders its colour pale; even $\frac{1}{16}$ of platinum can be detected in union in gold, from the colour.

9. Platinum is a most valuable metal; as it is not oxidable, nor fusible under common circumstances, and only difficultly combinable with sulphur, and not acted upon by common acids; it is admirably adapted for the uses of the philosophical chemist, and may be advantageously employed in all cases where gold is applied, unless the use is connected with the colour or malleability of the metal. The general application of platinum as a manufactured metal to the purposes of the laboratory is one of the many benefits which chemistry and the useful arts owe to Dr. Wollaston.*

XXXVII. *Arsenic or Arsenicum.*

1. Arsenic may be easily procured by heating the substance known by the name of white arsenic in powder with charcoal, in a Florence flask, or a glass tube; before the mixture becomes red hot, a metal sublimes, and condenses in the upper part of the vessel, which is arsenic.

* [Vide Phil. Trans. for 1829, for Dr. Wollaston's directions on this subject, contained in his "Bakerian Lecture on a Method of rendering Platina malleable."]

2. Arsenic is of a bluish white colour, not unlike that of steel. Its specific gravity is 8.31. It is very brittle: its point of fusion has not been ascertained; but it is the most volatile of all the metals, rising in vapour at about 356° of Fahrenheit's scale. When a part of it is suddenly ignited, it burns with a pale bluish light, sending off dense white fumes. It burns spontaneously in chlorine.

3. There are two known combinations of arsenic and oxygen; both of which are possessed of several of the properties of acids. The first is the substance formed by combustion, and this contains the smallest quantity of oxygen; the compound containing the largest quantity of oxygen may be formed by distilling nitrous acid, mixed with $\frac{1}{16}$ of its weight of muriatic acid from the other compound. The compound formed by combustion has been called *arsenious acid*, and likewise white oxide of arsenic. When procured by precipitation from acid solutions, it exists as a *hydrate*: it is fusible by a strong heat suddenly applied, but sublimes slowly at 383 Fahrenheit; after fusion it appears as a white vitreous substance, of specific gravity 5. It is soluble in 80 parts of water at 60°, and in 15 parts at 212°. Its taste is acrid, leaving an impression of sweetness. When heated its smell is like that of garlic.* The compound of arsenic with the largest proportion of oxygen is called *arsenic acid*. It is much more fixed in the fire than arsenious acid, is very soluble in water, and has an intense sour taste. From experiments on the quantity of oxygen absorbed by arsenic during its conversion

* [The vapour of metallic arsenic has a smell like that of garlic;—the oxide, when heated on glass or porcelain, has a slight peculiar smell, and produces a peculiar sensation in the throat, even when its vapour is inhaled in an extremely diluted state.]

into these two compounds, made by Proust and by myself, I conclude that the arsenious acid consists of about 25 of oxygen, and 75 of metal; and the arsenic acid of 33 of oxygen and 67 of metal. Hence it appears that the quantity of metal being the same, the oxygen in the arsenic acid is to that in the arsenious nearly as 3 to 2; and if the arsenious acid be supposed to contain two proportions of oxygen, the number representing arsenic will be 90; and those representing arsenious and arsenic acids will be 120 and 135.

4. The only compound of chlorine and arsenic known, is made by the combustion of the metal in chlorine, or by distilling a mixture of arsenic and corrosive sublimate together: it is a heavy limpid fluid, capable of being easily rendered gaseous, forming a liquid muriate of arsenic by the action of a small quantity of water, and affording a precipitate of arsenious acid merely by the action of a large quantity. From my experiments it appears that the compound of chlorine and arsenic, which may be called *arsenicane*, consists of 40 metal to 60 of chlorine; therefore it may be regarded as composed of two proportions of chlorine, and one of metal.

5. Arsenic combines with hydrogen. The best known substance containing the two bodies is *arseniu-retted hydrogen* gas. This elastic fluid, which was discovered by Scheele, may be procured by dissolving an alloy of 14 parts of tin and 1 of arsenic in muriatic acid. This substance has an extremely fetid smell; it burns when brought near an inflamed taper in the atmosphere; its flame is blue; and if the vessel in which it is burnt have a narrow neck, it deposits arsenic. It inflames spontaneously when acted upon

by chlorine : it is soluble only to a very slight extent in water. It is probable that the gas called *arseniuretted* hydrogen is always a mixture of a true gaseous compound of arsenic and hydrogen, with common hydrogen. Its specific gravity varies, as I have found, from 5 to 8, that of hydrogen being 1. When decomposed by attracting the arsenic from it, by the action of ignited metals, there is an expansion of volume. M. Gay Lussac and Thenard found 100 parts of it decomposed by tin become 140 parts. M. Stromeyer states that he analysed a gas consisting of 106 parts arsenic, and 2·19 hydrogen. This would agree with the idea that the pure gas is composed of 2 proportions of hydrogen, and 1 of metal; but such a gas must be more than twice as heavy as any specimen that has been weighed; and this circumstance leads me to doubt of the correctness of M. Stromeyer's results.

There is likewise a solid compound of hydrogen and arsenic : it may be procured by acting on water by an alloy of potassium and arsenic in great excess; it is a brown powder, which burns, when gently heated in the atmosphere, and which gives off *arseniuretted* hydrogen, when heated in close vessels. The same substance is procured when arsenic is made the negative surface in contact with water in Voltaic combinations. No experiments have as yet been made on the proportions of its elements.

6. Sulphur and arsenic readily unite by fusion, and form a red vitreous semitransparent mass. The same substance is found native in different parts of Europe, and is called *realgar*; it is often crystallized in transparent prisms: its specific gravity is 3·225. If Thenard's account of its composition be considered as accurate, it must consist of two proportions of arsenic, and three of sulphur, 180

and 90.* If sulphuretted hydrogen gas be made to act upon a solution of arsenious acid in muriatic acid, a fine yellow powder falls to the bottom. This powder is usually called *orpiment*. It may be formed likewise by subliming arsenic and sulphur together in a heat not sufficient to produce a fusion of the mass. It is composed of thin plates, which have a considerable degree of flexibility. According to Thenard, it contains more sulphur than realgar; but M. Proust states that by fusion it becomes realgar.

7. Arsenic readily combines with phosphorus, and they form together a black powder; the proportions of its elements have not yet been accurately ascertained.

8. Arsenic has not been combined with azote, boron or carbon.

9. Arsenic combines with most metallic substances. It renders the metals of the fixed alkalies less fusible by uniting to them, but its alloys with the common metals are usually very fusible. It renders gold and platina brittle, and gives whiteness to copper: none of the alloys containing it in any considerable quantity are malleable.

10. Arsenic is not much used in the arts. Realgar and orpiment are employed as pigments. The solutions containing arsenious, or arsenic acid, are extremely poisonous. The arsenious acid in very small doses, has been employed in medicine, particularly for the cure of intermittent fevers.

XXXVIII. *Molybdenum*.

1. There is an ore found in different parts of Europe,

* [From the latest analyses realgar appears to be a protosulphuret, and orpiment a sesquisulphuret.]

particularly in Sweden, not unlike plumbago, from which Scheele, in 1778, procured a white powder, and from this powder, Hielm, in 1782, obtained a metal, which he called *molybdenum*.

Pure molybdenum may be obtained either from the ore like plumbago, or from another ore found in Carinthia, called the molybdate of lead; by digesting them in powder in nitric acid, and boiling the mass in sulphuric acid; by lixiviation with water a liquor is obtained, which when acted on by solution of ammonia, passed through a filter and mixed with nitric acid deposits a white powder, and this powder intensely ignited in a charcoal crucible mixed with linseed oil, affords the metal.

2. Molybdenum, according to the observations of Bucholz, is brittle; its specific gravity is 8.611. Its colour is white. It burns when placed on ignited charcoal, and acted on by a current of oxygen gas, and gives off a white smoke which collects in small needle-formed crystals.

3. There are two well-defined combinations of molybdenum and oxygen. One is blue, the other is pale yellow; they both possess acid properties, and therefore may be distinguished by the names *molybdous* and *molybdic* acids. The *molybdic acid* is easily obtained from the ores of the metal by treatment with acids and ammonia; the white powder described in the last page is this substance combined with water, and it may be procured pure by ignition. Its specific gravity is 3.4; its taste is sour; it is fusible in a strong heat, and volatilized by intense ignition: it is soluble in about 1000 times its weight of water.

The blue acid, or the *molybdous acid*, is formed by triturating together in boiling water, one part of molyb-

denum in powder, and two parts of molybdic acid. The solution is to be passed through a filtre, and evaporated in a temperature not exceeding 120° ; the blue acid remains in the state of a fine powder. This acid is more soluble in water than the molybdic acid, and acts more intensely on vegetable blues, converting them to red. According to Bucholz, the blue acid consists of 100 parts of metal to 34 of oxygen, and the yellow acid of 100 parts of metal to 50 of oxygen. On these data it seems probable that molybdous acid consists of 2 proportions of oxygen and one of metal; and molybdic acid of 3 proportions of oxygen and 1 of metal; and assuming the composition of the molybdous acid as the foundation of calculation, the number representing molybdenum will be 88.2. Mr. Bucholz supposes that there are *oxides* of molybdenum, containing smaller quantities of oxygen than the two acids. It is probable that there is a brown oxide containing a single proportion of oxygen obtained by exposing the metal to a red heat; but what Bucholz considers as a violet brown oxide produced by heating the brown oxide, is probably only a mixture of the brown oxide and the blue acid.

4. No direct experiments have, I believe, been made on the action of chlorine on molybdenum; but when the molybdic acid is dissolved in muriatic acid, and the residue heated to redness, chlorine rises, and the blue acid remains behind; but a grayish sublimate is likewise formed, in which chlorine is indicated by the action of nitrate of silver.

5. Molybdenum combines readily with sulphur by fusion, or by heating strongly together molybdic acid and sulphur. The sulphuret of molybdenum is a black shining powder, the same as the native mineral from which Scheele first procured the acid. According to

Bucholz, it contains 60 of metal and 40 of sulphur per cent. and therefore may be considered as consisting of one proportion of metal and two of sulphur.*

6. Phosphorus combines with molybdenum; but the properties and constitution of the *phosphuret* have not been investigated.

7. Hydrogen, azote, carbon, and boron, have not been combined with molybdenum.

8. It unites to several of the metals; one of the most perfect of its alloys is with iron. With lead it forms an alloy somewhat malleable. Most of its other alloys break under the hammer; from the difficult fusibility of the metal it is not easy to make them uniform in their constitution.

Molybdenum has not yet been applied to any of the purposes of the arts.

XXXIX. *Chromium.*

1. There are two ores from which chromium may be procured; one is the red lead ore of Siberia, the chromate of lead; the other is the chromate of iron, which has been found in France and in North America.

Chromium was discovered by Vauquelin, in 1797. To obtain chromium, chromate of lead in fine powder is to be digested with moderately strong muriatic acid, till its power of action is exhausted. The fluid produced is to be passed through a filtre, and a little oxide of silver, such as is procured by precipitation from nitric acid by potassa, very gradually added to it till the whole solution becomes of a red tint, and no longer. This liquor by slow evaporation deposits small ruby red crystals, which, when intensely ignited, mixed with a little

* [Three distinct combinations of this metal with sulphur have been obtained, and the same number with chlorine and oxygen.]

charcoal powder, afford *chromium*. Chromate of lead may be procured from chromate of iron, by decomposing it by hydrate of potassa, making a solution in nitric acid, and adding solution of nitrate of lead; the chromate of lead falls down as a beautiful orange powder.

2. Chromium is a white brittle metal, requiring an intense heat for its fusion; it is very difficultly acted on by acids. It does not readily enter into combustion. Its specific gravity is 5.9.

3. Very few experiments have been made on the combinations of chromium. The red crystals procured from chromate of lead by muriatic acid appear to be a *hydrated acid*; they are soluble in water, have a sour taste, and combine with alkalies.

The red crystals strongly heated become a green powder, which is considered as an *oxide of chromium*. It is said that from 100 parts of the red crystals, 67 parts of metal may be procured. The acid of chromium, when combined with alkalies, precipitates most of the metallic solutions. In solutions of mercury it produces a vermilion red precipitate; in those of silver, a carmine red; in those of tin, a green. The name chromium has been given to the metal from its remarkable colouring powers.*

4. The artificial chromate of lead forms a beautiful and permanent pigment. I have found the orange colour most pure when the nitrate of lead used for the precipitation contained an excess of acid. The oxide of chromium has been employed for giving an emerald green colour to glass and enamel. Chromic acid is the colouring matter of the spinelle ruby; and oxide of chromium gives its beautiful colour to the emerald. The oxide of chromium has been lately found in some meteoric stones.

* From *Χρῶμα*.

DIVISION VI.

OF SOME SUBSTANCES, THE NATURE OF WHICH IS NOT
YET CERTAINLY KNOWN.

I. *Preliminary Observations.*

THE bodies to be examined in this division have been arranged into a distinct class, because they present some extraordinary and anomalous results, and because as yet the knowledge obtained respecting their nature is imperfect; many of the facts ascertained respecting them harmonize with the general doctrines of the science, and some of them offer new views respecting the arrangements and properties of matter; they are therefore amongst the most interesting objects of chemical inquiry.

II. *Of the Fluoric Principle.*

1. There is a substance found abundantly in nature called *fluor spar*; it is usually either blue, green, yellow, or white, transparent, and crystallized in cubes. It is a common product of the mines in Derbyshire.

2. When this substance, in fine powder, is mixed with oil of vitriol and distilled in retorts of silver or lead, connected with receivers of the same metal artificially cooled; an intensely active fluid is produced. It has the appearance of sulphuric acid, but is much more volatile, and sends off white fumes when exposed to air. It must be examined with great caution, for when

applied to the skin it instantly disorganizes it, and produces very painful wounds. When potassium is introduced into it, it acts with intense energy upon it, and produces hydrogen gas, and a neutral salt: when lime is made to act upon it, there is a violent heat produced, water is given off, and the same substance as fluor spar is produced. When it is dropped into water a hissing noise is produced with much heat, and an acid fluid not disagreeable to the taste is formed if the water be in sufficient quantity. It instantly corrodes and dissolves glass.

3. If, instead of being distilled in metallic vessels, the mixture of fluor spar and oil of vitriol be distilled in glass vessels, little of the corrosive liquid will be obtained; but the glass will be acted upon, and a peculiar gaseous substance will be produced, which must be collected over mercury. The best mode of procuring this gaseous body is to mix the fluor spar with powdered glass or powdered quartz, and in this case the retort may be preserved from corrosion, and the gas obtained in greater quantities. This gas, which is called *silicated fluoric gas*, is possessed of very extraordinary properties.

It is very heavy; 100 cubical inches of it weigh 110.77 grains, and hence its specific gravity is to that of hydrogen nearly as 48 to 1. When it is brought in contact with water it instantly deposits a white gelatinous substance, which is hydrate of silica, and the water becomes an acid solution of silica; it produces white fumes when suffered to pass into the atmosphere. It is not affected by any of the common combustible bodies, but when potassium is strongly heated in it, it takes fire and burns with a deep red light; the gas is absorbed, and a fawn coloured substance is formed, which yields

alkali to water with slight effervescence, and contains a combustible body, and the washings afford potassa and a salt, from which the strong acid fluid mentioned in the last section may be procured by sulphuric acid.

4. If instead of glass or silica the fluor spar be mixed with dry vitreous boracic acid, and distilled in a glass vessel with sulphuric acid, the proportions being one part boracic acid, two fluor spar, and 12 oil of vitriol: the gaseous substance formed is of a different kind, and is called *the fluoboric gas*. 100 cubical inches of it weigh 73·5 grains, so that its specific gravity is rather more than 32 times that of hydrogen, When a little of it is suffered to pass into the atmosphere it produces fumes much more dense than those produced by the gas described in the last section, and which appear white, and almost opaque. It is absorbed rapidly by water, and forms with it a dense fluid like sulphuric acid in appearance and consistence; and when water is saturated with it at 50° Fahrenheit, it contains 700 times its volume of gas, and is of specific gravity 1·77. When potassium is heated in this gas, it takes fire, and burns with a red light; the gas is destroyed, if the metal be in sufficient quantity, and an olive coloured substance is produced, which seems to be principally boron, and a neutral salt, which by the action of oil of vitriol, affords the dense fluoric acid, and sulphate of potassa.

5. It appears extremely probable, from all the facts known respecting the fluoric combinations, that fluor spar contains a peculiar acid matter; and that this acid matter is united to lime in the spar, seems evident from the circumstance that gypsum, or sulphate of lime, is the residuum of the distillation of fluor spar and sulphuric acid: the results of experiments on the decom-

position of fluor spar, have been differently stated by different chemists; the maximum of sulphate of lime obtained from 100 grains in some experiments made in the laboratory of the Royal Institution, was 174·2 grains, and from this result fluor spar may be supposed to consist of 55 lime, and about 20·7 fluoric acid, and this last number will represent the acid.

6. The dense acid fluid described in 2. must, on the same idea, be supposed to be a compound of an acid unknown in a separate state, and water; and may be called *hydrofluoric* acid; and supposing all the water in oil of vitriol transferred to it, it will consist of 20·7 fluoric acid, and 17 water.

7. The gas formed by the action of hydro-sulphuric acid on a mixture containing silica and fluor spar, silicated fluoric gas must be regarded as a kind of neutro-saline gaseous compound consisting of fluoric acid and silica, and it has been found to afford, when decomposed by solutions of ammonia, 61·4 per cent. of silica; it may therefore be supposed to consist of two proportions of acid 41·4, and one of silica 61. According to this view of its composition, the number representing it is about 102; 1 volume of it condenses 2 volumes of ammonia, and they form together a peculiar saline substance which is decomposed by water. The composition of this salt is easily reconciled to the numbers above given, as representing silica and fluoric acid, on the supposition that it contains one proportion of ammonia, and one of silicated fluoric acid; and calculating the number of silicated fluoric acid on this supposition it would be about 99.

There is great reason to believe that, when potassium burns in this gas, it is the acid matter which is decomposed, and that it consists of oxygen united to an in-

flammable basis; for if it were the silica alone which is decomposed, or if a mere combination were formed between the potassium and the acid gas, the same quantity of fluete of lime or fluor spar ought to be formed from equal quantities of silicated fluoric acid, acted upon by potassium, and afterwards exposed to solution of ammonia, and the other absorbed by water, and acted on by solution of ammonia; which I have found is not the case, for in the first instance there is considerably less produced. In the experiment, it seems likely that the potassium acquires oxygen principally from acid matter combined with the silica, and that the inflammable basis of the acid partly combines with the potassa, and partly with the silica, or with silicum; and forms with the first a compound that effervesces, and is partly decomposed by water; and with the second an insoluble substance, which affords silicated fluoric acid by absorption of oxygen.

8. It is extremely likely that fluoboric acid gas is composed of the peculiar acid which is supposed to consist of oxygen and an inflammable basis, and boracic acid; but it appears that in the combustion of potassium in this gas it is the boracic acid alone that is decomposed, and that the fluoric acid combines with the potassa formed.

9. It is a peculiar circumstance with respect to the fluoric principle, that silicated fluoric gas, and fluoboric gas combine with bodies without decomposition. Thus they both form peculiar compounds with the alkalies; and though silica is deposited by the action of silicated fluoric gas on water, and on other oxidated bodies, yet the new compound formed always appears to contain part of the earth, which is supposed to be a constituent of the gas. In general, silica and boracic acid can only

be procured from the two gases by the intervention of bodies that contain water or oxygen: this circumstance, if it were not opposed by the results of the experiments on the action of potassium on silicated fluoric gas, which, however, ought to be repeated, might lead to the suspicion, that the fluoric gases are compounds of a principle unknown in the separate state, but analogous to chlorine, with silicum and boron; that the hydrofluoric acid is a compound of the same principle with hydrogen and water, and fluor spar a compound of the same principle with calcium.

10. If 20·7 be really the number representing fluoric acid, it can be supposed to contain only one proportion of oxygen, and the fluoric basis will be represented by 5·7, and it will be the only known acid so constituted.*

11. Silicated fluoric gas, when absorbed by water, affords an acid fluid, which, when acted upon by ammonia, deposits silica; and in glass vessels pure hydrofluoric acid cannot be obtained. Silicated fluoric gas seems to form only one combination with ammonia, which deposits silica by the action of water; but fluoroboric gas forms three combinations with ammonia, one volume of it condenses 1, 2, and 3 volumes of ammonia. The saline compound containing the least quantity of ammonia is solid, the other two compounds are fluids at the common temperature of the atmosphere.

12. The only use to which the fluoric combinations have as yet been applied is for etching on glass; for this purpose the hydrofluoric acid, or the fluuate of am-

* [From after and more extended inquiry, which will be given in a subsequent volume, he concluded that the fluoric principle is not an inflammable substance, but analogous to chlorine, and that the three acids above mentioned are compounded of it and of hydrogen, boron, and silicum respectively, as hinted at in the preceding paragraph.]

monia, should be used; the gases have no action on glass.

13. Silicated fluoric gas, and diluted hydrofluoric acid were discovered by Scheele, in 1771. Margraff, three years before, had pointed out some of the results of the action of acids on fluor spar; concentrated hydrofluoric acid, and fluoboric gas were made known by some elaborate researches of Gay Lussac and Thenard, in 1809. My brother, Mr. John Davy, in 1810 and 1811, extended the knowledge of the properties of these bodies, and the modes of procuring them pure, ascertained the specific gravity of fluoboric and silicated fluoric gases, and the proportions in their ammoniacal combinations. The action of potassium on silicated fluoric gas, and fluoboric gas was investigated by MM. Gay Lussac and Thenard, in 1809; and I made a number of experiments on the same subject about the same time.

III. *Of the Amalgam procured from Ammoniacal Compounds.*

1. When a globule of pure mercury is negatively electrified by a Voltaic apparatus of 100 pair of plates, it being in contact with solution of ammonia in a cavity made in a piece of muriate of ammonia, or any ammoniacal salt, moistened in such a manner, and so placed on a disc of platina, that the circuit is completed; the globule rapidly increases in volume, the quicksilver loses its fluidity, and at length becomes of the consistence of soft butter, and arborescent crystallizations shoot from it, which are quite solid. The amalgam so formed has perfectly metallic characters. It effervesces copiously when thrown into water, hydrogen gas is

given off, and a solution of ammonia is found in the water. When exposed to the air it gradually loses its consistence; it emits a strong odour of ammonia, and reddens paper tinged with turmeric held above it; and at last is found merely quicksilver.

This curious experiment was made about the same time by Dr. Zeebeck of Jena, and by MM. Hissinger and Berzelius of Stockholm, before the middle of the year 1808; and they were led to make it in consequence of my experiments on potassa and soda.

2. I found a still more easy mode of making the amalgam by employing mercury combined with a minute quantity of potassium, sodium, or barium. When a compound of this kind is placed in contact with a solution of ammonia, or any moistened ammoniacal salt, it enlarges to eight or ten times its bulk, and becomes a soft solid, and may be preserved for a much longer time than the amalgam formed by electrical powers; it changes very slowly even under water.

3. Different opinions have been formed, and may still be formed, concerning the nature of this extraordinary substance. M. Berzelius supposes that ammonia consists of a peculiar metal combined with oxygen, and of which metal hydrogen and azote are both peculiar oxides; this idea was one that I started likewise soon after the discovery of the amalgam.

4. Another view of the subject is, that the amalgam consists of mercury united to azote and hydrogen, the hydrogen being in larger proportion than in ammonia; and this view has been embraced and defended by MM. Gay Lussac and Thenard; but the subject is still obscure and mysterious, and the true theory of the experiment can only be developed in consequence of new facts.

5. Soon after the discovery of the amalgam, I attempted to procure a peculiar metallic substance from it by distillation out of the contact of air, but without success; whether I used the amalgam formed by electricity, or that procured by the intervention of the alkaline metals; on the application of heat, hydrogen and ammonia were always evolved, and the mercury recovered its former state. On the idea of the basis of ammonia being a peculiar metal, of which azote and hydrogen are oxides, these results can only be explained by supposing that the amalgam being formed from moist substances, sufficient water adheres to it, to afford oxygen, and to produce the gaseous matter; and the most perfect amalgam does not yield a quantity of gaseous matter equal to more than $\frac{1}{800}$ of its weight.

I procured ammonia and hydrogen by heating the amalgam, however, in cases in which it was carefully wiped with bibulous paper, and in which there was no appearance of adhering moisture; and similar results have been obtained by MM. Gay Lussac and Thenard.

In the most accurate experiments the proportions of ammonia and hydrogen were two to one in volume.

6. There is no instance known of mercury retaining its metallic characters in combination with any other substance than a metal; and it seems very probable that, if the matter existing in the amalgam from ammonia could be procured in its perfect form, and could be exhibited as a solid under pressure, and at a very low temperature, it would appear as an extremely light metallic substance. On the idea of its being a compound of azote and hydrogen it will consist of one proportion of azote 26, and 8 of hydrogen 8; and the number representing it will be 34.

It is very difficult, but not however altogether impos-

sible to reconcile the idea of the substance in the amalgam being elementary, with analogies belonging to the general series of definite proportions. On such a supposition azote must necessarily contain more than four times as much oxygen as hydrogen; and if 1 of basis to 5 of oxygen, be supposed in hydrogen, then there will be 1 to 25 in azote, and 1 to 40 in nitrous oxide, 1 to 55 in nitrous gas, 1 to 85 in nitrous acid, and 1 to 15 in ammonia; and 5, 15, 25, 40, 55, and 85, form a series of numbers having definite relations to each other.

If the hypothesis of the elementary nature of the substance in the amalgam be adopted, water must be supposed to be constituted by 1 basis, and 50 of oxygen.

It is extremely unlikely that such proportions should exist, and the general tenor of our knowledge of chemistry, as well as the results of the experiments, render it much more probable that the amalgam is composed of quicksilver, azote, and hydrogen.

DIVISION VII.

ON THE ANALOGIES BETWEEN THE UNDECOMPOUNDED SUBSTANCES; SPECULATIONS RESPECTING THEIR NATURE; ON THE MODES OF SEPARATING THEM, AND ON THE RELATIONS OF THEIR COMPOUNDS.

I. *Of the Analogies between the Undecomposed Substances; Ideas respecting their Nature.*

1. The undecomposed substances most analogous to each other, are certainly to be found amongst the metals; some of these are so similar that it requires

refined observation, and sometimes experiment to distinguish them. There is likewise a chain of gradations of resemblance which may be traced throughout the whole series of metallic bodies, at the same time that certain similar and characteristic properties are found to belong to metals in other respects most unlike each other.

Silver and palladium, antimony and tellurium, agree in a great number of qualities. Potassium and platinum, if we except their lustre, colour, and power of conducting electricity, are bodies extremely dissimilar; yet, by arranging the metals in the order of their natural resemblances, these two substances may be made parts of one chain of natural bodies: potassium, sodium, and barium are very like each other; barium approaches to manganesum, zinc, iron, tin, antimony. Platinum is analogous to gold, silver, and palladium; and palladium is connected by distinct analogies with tin, zinc, iron, and manganesum. Arsenic and chromium, though amongst the most dissimilar of the metals in other respects, agree in the property of forming acid matter by combination with oxygen.

Amongst the inflammable bodies not metallic there are analogies, but not a similar series. Sulphur and phosphorus agree in many respects; carbon and boron are likewise analogous, and are connected by distinct relations with the metallic substances. Azote, whilst it agrees with the other combustible bodies that have been named in forming an acid by saturation with oxygen, is analogous to carbon in its incapacity of uniting to chlorine.*

Chlorine and oxygen are separated from the inflammable bodies by a number of marked distinctions; yet

* [This remark does not now apply; vide p. 196, and note p. 225.]

sulphur agrees with chlorine in forming an acid by combining with hydrogen; and has a weak attraction for chlorine, and a strong attraction for metallic substances.

2. As far as our knowledge of the nature of compound bodies has extended, analogy of properties is connected with analogy of composition; if one of the inflammable solids or metals is proved to be compound, there would be strong evidence for supposing that the others were likewise compounded. It has been already mentioned that sulphur and phosphorus, when Voltaic electrical sparks are taken in them in a state of fusion, afford hydrogen gas. I found likewise that when an alloy of tellurium and potassium was acted upon by melted sulphur, telluretted and sulphuretted hydrogen equal to at least 80 times the volume of the sulphur were disengaged. I have made many experiments of this kind with similar results, the sulphur being recently sublimed in azote, and moisture being excluded with the greatest care. In the experiments of Voltaic electrization, it might be supposed that the hydrogen being only in a very small quantity might belong to an accidental admixture in the sulphur and the phosphorus; but the proportion is too large in the experiments on the action of tellurium, potassium, and sulphur, to allow of a similar inference, and it seems more probable that it arises either from the decomposition of the sulphur, or of the metals, or all of these bodies.

3. We know nothing of the true elements belonging to nature; but as far as we can reason from the relations of the properties of matter, hydrogen is the substance which approaches nearest to what the elements may be supposed to be. It has energetic powers of combination, its parts are highly repulsive as to each other, and

attractive of the particles of other matter; it enters into combination in a quantity very much smaller than any other substance, and in this respect it is approached by no known body.

After hydrogen, oxygen partakes most of the elementary character; it has perhaps a greater energy of attraction, and next to hydrogen is the body that enters into combination in the smallest proportion.

4. I have already hinted at the idea that all inflammable matters may be similarly constituted, and may contain hydrogen. And on this supposition they may be conceived to owe their powers of combining both with oxygen and chlorine, to the attractive energies of their combined hydrogen.

On the most probable view of the nature of the amalgam from ammonia, as I have mentioned, it must be supposed to be composed of hydrogen, azote, and quicksilver; and it may be regarded as a kind of type of the composition of the metals; and by supposing them and the inflammable bodies different combinations of hydrogen with another principle as yet unknown in the separate form; all the phenomena may be easily accounted for, and will be found in harmony with the theory of definite proportions.

The metal of ammonia or *ammonium* must be supposed to be constituted by 8 of hydrogen, and 26 of azote; and as azote unites to 5 proportions, 5 times 15 of oxygen, it may be supposed to contain 10 proportions of hydrogen; and its constitution may be thus expressed, 10 proportions of hydrogen and 16 proportions of an unknown basis. Ammonium, on the same hypothesis, will consist of 16 unknown basis, and 18 hydrogen. Potassium, the number representing which is 75; as it combines with 3 proportions of

oxygen, may be supposed to consist of 69 unknown basis, and 6 hydrogen. Sodium, which is represented by 88, and which likewise combines with 3 proportions of oxygen, may be considered as consisting of 82 basis, and 6 hydrogen. Tin, the number of which is 110, and which combines with 2 proportions of oxygen, may be supposed to be constituted by 106 of basis and 4 hydrogen; and silver, which is represented by 205, of 203 of basis, and 2 hydrogen. Amongst the acidifiable bodies, sulphur, which is represented by 30, may be supposed to consist of 6 hydrogen, and 24 basis; phosphorus of 4 hydrogen, and 16 basis; and charcoal of 4 hydrogen and 7.4 basis. It will be unnecessary to supply any more of these estimations, the principles of which are obvious; and in an elementary book it would be improper to dwell upon matters of mere speculation; even these transient views have been developed merely for the sake of pointing out a promising path of enquiry.

5. In supposing the quantity of hydrogen in the inflammable solids and metals denoted by the quantity of oxygen or of chlorine they absorb, it is taken for granted that the hydrogen forms only water or muriatic acid in the new combination, but it is possible that hydrogen may combine with oxygen and chlorine in many different proportions, and that its union with a peculiar basis may modify its power of attraction; so that even allowing the general hypothesis, no confidence can be placed in the numerical expressions of the proportions of hydrogen and basis; they are offered merely as possible circumstances.

6. The probabilities that the metals and inflammable solids may be constituted by different and various proportions of hydrogen and an unknown basis, are, how-

ever, strengthened by the fact, that the metals in which hydrogen is supposed to be attracted by the largest quantity of other matter, are the least disposed to combine with oxygen and chlorine; and those that are supposed to contain the largest quantity of hydrogen to the smallest quantity of other matter, are the most combustible, and likewise those supposed to contain the largest, and consequently the least attracted quantity of hydrogen, have the lowest specific gravity.

7. When the analogy of the oxides to many of the hydrates, and that of the combinations of chlorine to many of the neutral salts, is considered, bodies so much alike that till lately they have been confounded together; the view that the inflammable bodies contain hydrogen becomes still more likely. Water cannot be separated from the hydrates of potassa or soda by heat; and the hydrate of lime is exceedingly analogous to the pure earth; and supposing the oxides to be compounds of unknown bases and water, it might be expected that the water would adhere to them with great energy, and would only be separated in consequence of the bases entering into a new combination.

Common salt is very analogous to sulphate of potassa and other bodies known to consist of acid matter and alkaline matter; and if sodium consist of a basis combined with hydrogen, then common salt may be considered as composed of the same basis united to muriatic acid.

8. Chlorine and oxygen agree in many of their characters; but the weight of chlorine, its colour, its absorbability by water are all in favour of its being a compound. The number representing chlorine is so high that it may include four proportions of oxygen; and if this body be supposed to consist of oxygen united to

an unknown basis, the analogy of the combinations of chlorine, both to the oxides and the salts, might be easily explained. The evidences in favour of such an idea of the constitution of chlorine are, however, much inferior to those which render it probable that the inflammable solids contain hydrogen; and this speculation on the composition of chlorine must not be confounded with the notion that chlorine is a compound of oxygen and muriatic acid free from water; for supposing a basis to exist in chlorine, it does not follow that it will be acid in its nature. The characteristic acid belonging to the combinations of chlorine is formed by the union of that body with hydrogen; and sulphur likewise forms an acid by combining with hydrogen.

9. I have mentioned, page 125, that in the electrization of a globule of mercury in water, oxygen appears to be combined with the metal, and yet no hydrogen evolved. I have made a number of experiments on this subject, and have ascertained that, in the process described, oxide is formed, without any apparent compensation in the production of inflammable matter; nor was I able to detect any combination into which the hydrogen could have entered;* so that these experiments, as they now stand, would induce the belief that water is the ponderable basis of both oxygen and hydrogen, and that these two forms of matter owe their peculiar properties either to the agency of imponderable substances, or to peculiar arrangements of the particles of the same matter; but such a formidable conclusion as

* [From the experiments of Sir J. F. Herschel it appears that in the process above referred, potassium or sodium was reduced, according to the kind of saline matter present, and which, combining with the mercury, imparted peculiar properties.—Vide Bakerian Lecture in Phil. Trans. for 1824, "On certain Motions produced in Fluid Conductors when transmitting the Electric Current."]

this must not be hastily adopted, for in all other cases oxygen and hydrogen appear as perfectly inconvertible substances, and in no other instance can one be procured from water without the correspondent quantity of the other, or without some product in which the other may be supposed to enter. In all cases in which the circuit appears to be interrupted even, this is the case. When the finger is plunged in a glass of water connected with a wire of platina positively electrified from the battery of 2000 double plates of the Royal Institution, oxygen is produced, and there is no appearance of hydrogen; but in this case the body is connected with a floor containing moisture, and at the extreme point of the moist surface, where it is in contact with a metallic body, hydrogen must be disengaged; and the same changes occur if a circuit be made through eight persons, their hands being in contact, the two forming the extremity of the chain, having their fingers plunged in two glasses connected by wires of platinum with the two poles of the battery; hydrogen is produced from one wire, and oxygen from the other. Till I ascertained that even acids and alkalies could be attracted from a central vessel in the Voltaic circuit to the two extremities of the positive and negative metallic surfaces, it appeared very mysterious that oxygen and hydrogen should be separately produced in the Voltaic electrization of water; but if it be possible for lime to be attracted through sulphuric acid to the negative surface, it seems equally possible that hydrogen may be attracted through the moisture in a living body; or a series of decompositions and recompositions may be simultaneously produced throughout the whole extent of the moist surface, by which, whilst a particle of oxy-

gen is produced at one extremity of the chain, a particle of hydrogen is evolved at the other.

10. There is, however, no impossibility in the supposition that the same ponderable matter in different electrical states, or in different arrangements, may constitute substances chemically different: there are parallel cases in the different states in which bodies are found, connected with their different relations to temperature. Thus steam, ice, and water, are the same ponderable matter; and certain quantities of ice and steam mixed together produce ice-cold water. Even if it should be ultimately found that oxygen and hydrogen are the same matter in different states of electricity, or that two or three elements in different proportions constitute all bodies, the great doctrines of chemistry, the theory of definite proportions, and the specific attractions of bodies must remain immutable; the causes of the difference of form of the bodies supposed to be elementary, if such a step were made, must be ascertained, and the only change in the science would be, that those substances now considered as primary elements must be considered as secondary; but the numbers representing them would be the same, and they would probably be all found to be produced by the additions of multiples of some simple numbers or fractional parts.

11. That the forms of natural bodies may depend upon different arrangements of the same particles of matter has been a favourite hypothesis advanced in the earliest era of physical research, and often supported by the reasonings of the ablest philosophers. This sublime chemical speculation sanctioned by the authority of Hooke, Newton, and Boscovich, must not be confounded with the ideas advanced by the alchemists

concerning the convertibility of the elements into each other. The possible transmutation of metals has generally been reasoned upon, not as a philosophical research, but as an empirical process. Those who have asserted the actual production of the precious metals from other elements, or their decomposition, or who have defended the chimera of the philosopher's stone, have been either impostors, or men deluded by impostors. In this age of rational inquiry it will be useless to decry the practices of the adepts, or to caution the public against confounding the hypothetical views respecting the elements founded upon distinct analogies, with the dreams of alchemical visionaries, most of whom, as an author of the last century justly observed, professed an art without principles, the beginning of which was deceit, the progress delusion, and the end poverty.

II. *Of the Analogies between the Properties of the Primary Compounds, and on their Chemical Relations.*

1. In those compounds which contain the same element combined with bases that resemble each other, a very great degree of similarity might be expected ; and it is found that a number of secondary combinations are still more analogous to each other than any of the undecomposed bodies. Ittria and glucina, baryta and strontia, potassa and soda are instances of bodies which, as to many of their properties, might be mistaken for each other ; and a chain of analogies may be traced through all the combinations of inflammable bodies and metals with chlorine, oxygen, and each other. All the acids, alkaline earths, alkalies, and combinations of chlorine in their pure states at common temperatures

are non-conductors of electricity; by far the greater number possess a certain degree of transparency; in their combinations with each other they display analogous results; most of them form hydrats; they render solid a certain quantity of water, and are usually dissolved by a greater quantity; and even acids combine with each other in consequence of the intermedium of water, as is the case with the sulphurous and nitrous acid gases.

Libavius's liquor is a limpid fluid; if mixed with a certain proportion of water it becomes a solid crystalline body. The glacial oil of vitriol, and the hydrophosphorous acid are instances of oxidated bodies forming crystalline solids with water.

2. The earths and the oxides which are insoluble in water still condense a certain quantity of this fluid, and it gives a greater fusibility to those which retain it with sufficient energy to be submitted to a strong heat. All oxides and earths obtained by precipitation from aqueous solution, that I have examined, are hydrats, and such of them, as I have carefully analyzed, I find contain the water in definite proportions. The combination of an earth, an alkali, or an oxide, with water may be considered as amongst its weakest combinations, for the water is expelled by carbonic acid. The expulsion of water from the earths seems to be connected, as I stated in page 50, with the contraction of volume, which many of them undergo by ignition: the particles, when the water is driven off, approach nearer to each other, and a great contraction is the result, and probably sometimes a semi-fusion. This quality on which, as it has been stated, the pyrometer of Wedgwood is founded, is elegantly exemplified in an experiment I have lately made on the hydrate of zircona. When this body is

heated, at the moment of the expulsion of the water, there is so great and rapid a contraction of the particles of the earth, that they become incandescent in the process; and, from being as soft as resin, become sufficiently hard to scratch rock crystal.

3. In general those compounds of oxygen, the bases of which combine with most energy, likewise exert the greatest force of attraction on each other; such, for instance, are the metals of the fixed alkalies in their relations to sulphur, phosphorus, arsenic, and tellurium; and potassa and soda readily combine with the acids of sulphur, phosphorus, and arsenic, and with the oxide of tellurium.

4. No refined experiments have as yet been made on the mutual action of these compounds of chlorine and oxygen, which are capable of co-existing; but the salts called hyperoxymuriates are substances in which chlorine and oxygen exist combined with metals; and the facility with which they are decomposed depends upon the tendency of the metal to unite to chlorine, so as to form a binary compound, a circumstance connected with the expulsion of the oxygen. The hyperoxymuriate of potassa, when it was first formed by Dr. Higgins, was supposed by him to be a species of nitre, from the similarity of its obvious properties: and it is remarkable that its composition is the same as that of nitre, except that in the first salt there is a proportion of chlorine, and in the second one of azote. Hyperoxymuriate of potassa consists of 1 proportion of potassium 75, 6 of oxygen 90, and 1 of chlorine 67. Nitre consists of 1 of potassium 75, 6 of oxygen 90, and 1 of azote 26. The combinations of ammonia with the compounds of chlorine, offer a class of curious bodies to the chemical inquirer, the properties of which have

never been investigated: that formed by phosphorana, and referred to page 215, is a most extraordinary substance, and its elements are combined with a degree of energy which renders it analogous to a primary compound.

5. In the combinations of ammonia with acids and oxides the hydrogen of the ammonia is always in some definite proportion to the oxygen of the acid or oxide, so that water may be formed by the decomposition of the compound; this is obvious from the decomposition of the fulminating ammoniacal metallic compounds. If a solution of ammonia be poured into a solution of gold, a brown powder falls down, which when washed and dried, explodes by a gentle heat. I caused it to detonate in small quantities in exhausted glass retorts, and I found that the products were water, azote, and gold.

Fulminating silver is a compound in which the elements seem to be in similar relations to each other; it was discovered by M. Berthollet, and may be made by dissolving the oxide of silver, procured from the nitrous solution by lime water in solution of ammonia at common temperatures, and exposing the mixture to spontaneous evaporation; black crystals form, which must be examined with great caution, and only in small quantities, as they explode by the mere contact of a soft body.

6. The extensive class of bodies called neutral salts are formed by the mutual action of acids and oxides, alkalies and earths; and in general those oxidated bodies that contain least oxygen, are such as most readily enter into combination with acids; thus the peroxides generally are either insoluble in acids, or require the abstraction of a portion of oxygen to become soluble; and in general two inflammable bodies in com-

binning with oxygen, unite to less than the added sums of the quantity they would separately combine with to saturation. Many of the neutral salts may be considered either as combinations of peroxides with inflammable bases or as alkalies united to acids, or as peroxides united to oxides; for instance, the compound formed from sulphurous acid gas and potassa consists of potassium and sulphur, with three proportions of oxygen, and may be regarded as a compound of peroxide of potassium and sulphur. Sulphate of potassa contains four proportions of oxygen, and might be regarded as a compound of peroxide of potassium, and oxide of sulphur. They are in fact all compounds of oxygen with double bases; and when one fixed alkali, or earth, or oxide, separates another, it may be supposed that the basis only is changed: thus, where hydrate of potassa separates lime from its nitric solution, it may be conceived that the potassium only takes the place of calcium. And that the oxygen and water of the hydrate of potassium unite to this metal, and that the potassium unites to the oxygen, nitrous acid and water of the solution.

7. It is very easy to estimate the composition of any of the combinations of alkalies, earths, or oxides with acids, by adding together the numbers representing their elements; thus sulphate of soda is composed of 60 sulphur, 90 oxygen, which make two proportions of sulphuric acid; and 88 of sodium, and 30 of oxygen, which make one proportion of soda. Carbonate of lead is composed of two proportions of carbonic acid, equal to 82·8, two proportions of oxygen 30, and one of lead 398. Sulphate of lead is composed of two proportions of sulphuric acid 150, two of oxygen 30, and one of lead 398: sulphate of nickel of two proportions of sulphuric acid 150, and one of oxide of nickel 141; and

these proportions agree almost precisely with the best analysis.

8. It appears that in the neutrosaline compounds in which there is a perfect harmony between the proportions of the elements, the result is neutralization; and that in this case a crystalline compound, or an insoluble compound is usually formed. Thus in the instances above mentioned, in the sulphates of soda and lead; the sulphur is a binary proportion, and the oxygen a binary proportion, or a multiple of a binary proportion; and in the carbonate of lead, the carbon is a binary proportion, and the oxygen a multiple of a binary proportion; and to give another instance, in the sulphate of barytes the sulphur is a single proportion, and the oxygen a single proportion, or a multiple.

When, on the contrary, there is a want of harmony in the proportions, the excess either of acid or basis seems to be shewn in the properties of the result; and it is seldom a crystallized body. Thus in the soluble red sulphate of iron, the number of proportions of oxygen in the oxide are three, and those of the sulphur in the acid are four: and this body is strongly acid and uncrystallizable.

III. *On the relative Attractions of the undecomposed Substances for each other.*

1. The attractions of the undecomposed substances vary with the temperature, probably, chiefly in consequence of their different degrees of volatility; for although freedom of motion in the parts of bodies wonderfully promotes combination, yet the disposition in bodies to assume the aëriform state at high temperatures, enables decompositions to take place in an order which

would not be expected from the known agencies of the substances under common circumstances.

2. The bodies that follow are arranged in the order of their attractions for oxygen, at the lowest temperature of visible ignition, after the results of my own observation. Potassium, sodium, barium, boron, carbon, manganese, zinc, iron, tin, phosphorus, antimony, bismuth, lead, sulphur, arsenic, tungstenum, azote, palladium, mercury, silver, gold, platinum.

3. The attractions of bodies for chlorine follow an order not very different, though with some exceptions; potassium, sodium, zinc, iron, lead, silver, antimony, bismuth, phosphorus, copper, sulphur, mercury, platinum, gold.

4. The attractions of the undecomposed bodies for sulphur have not been determined to any extent. Potassium and sodium seem to have the highest attraction of any substances: then iron, copper, antimony, palladium, lead, and silver.

5. No bodies combine with phosphorus with more energy than the metals of the fixed alkalies: and after them platinum, zinc, antimony, and sulphur, appear to have the strongest attractions; but no very definite knowledge has been as yet obtained on the relations of the phosphurets.

6. The general phenomena of the decomposition of the binary compounds, by undecomposed bodies, can require no illustration. Potassium separates chlorine and oxygen from all known bodies; usually it produces potassa, but sometimes by acting on compounds containing abundance of oxygen, it forms the peroxide of potassium. Carbon, in reducing metallic oxides, forms either carbonic acid, or gaseous oxide of carbon, according as the oxygen is more or less strongly

attracted by the basis. When oxides are decomposed by sulphur, sulphurous gas, and sulphurets are almost always formed.

7. Some of the instances which were formerly supposed instances of single attractions are now known to be connected with double attractions. This is remarkably the case in the production of potassium by iron. The water in the hydrate of potassa, and the potassa seem to be decomposed at the same time; the iron unites to the oxygen of both; the hydrogen and potassa combine; and their gaseous compound deposits potassium on cooling.

IV. *On the Methods of separating the Undecomposed Bodies from each other.*

1. General methods of separating the undecomposed bodies from each other may be learnt from a consideration of the processes by which they are procured; but there are other modes which apply to many of their compounds, and which are still more simple.

2. As all the undecomposed bodies differ in the manner in which they are affected by heat, many of them may be separated from compounds, by exposing them to different temperatures. Thus oxygen, chlorine, mercury, phosphorus, and sulphur may be detached from many bodies by the process of ignition.

3. In most cases, however, complicated methods are necessary, particularly in cases when the bodies are united to oxygen and acids or to chlorine; the compounds of chlorine differ very much in volatility, and in cases when they are mixed together, they appear to act upon each other with very little energy only: hence, if

it is possible to combine all the elements of a compound with chlorine, by the action of the gas, or of muriatic acid, or nitro-muriatic acid, they may be easily separated by the application of a heat gradually increased. Amongst the metallic combinations, that of tin when saturated with chlorine rises first, then those of arsenic, antimony, tellurium, iron, zinc, bismuth, in the order in which they have been named.

Silver is easily separated from solutions in which it exists by muriatic acid, with the chlorine of which it forms an insoluble compound; and in the same way chlorine is separated, and its quantity in any substance ascertained by means of solution of silver.

Oxide of iron is separated from solutions by succinate of ammonia, with which it forms an insoluble salt. The oxides of copper, nickel, and cobalt, are all soluble in ammonia. Those of zinc, tellurium, tin, and platinum, in solution of potassa. Acids are separated by alkalies; and alumina, silica, zircona, ittria, glucina, and the alkaline earths, may with facility be detached from their combinations by the action of acids, alkalies, and carbonates. Oxide of lead and baryta may be easily separated from other bodies, in consequence of their forming insoluble compounds with sulphuric acid.

4. The order in which metals precipitate each other from solutions is nearly in the ratio of their attraction for oxygen; and in all cases of neutral compounds, the precipitating metal takes the oxygen and acid of the metal thrown down. Iron readily precipitates copper; zinc readily throws down tin, lead, tellurium, bismuth, &c. and in general the metallic substances, as has been stated page 107, attract oxygen, and precipitate each other in a ratio connected with their electrical relations;

those that are positive with respect to others having the highest attractive powers for oxygen and acids.

By Voltaic electricity all substances are separated from their compounds with oxygen and chlorine; or alkalies, earths, and oxides are separated from acids; as has been mentioned page 116, and that in an uniform order and in definite proportions, so that Voltaic electricity offers general methods of decomposing all compounds soluble in water; and for most experiments of this kind very small combinations only are necessary: if small quantities of the materials are employed, two or three double plates are sufficient for decomposing most metallic solutions. The energies of small powers in acting upon bodies by diminishing the quantities exposed to their agency, has been happily shown by Dr. Wollaston, in the decomposition of water by a common small electrical machine, by passing the electricity from surfaces of about the $\frac{1}{1000}$ of an inch square; and the same philosopher has produced the ignition of platinum in leaf of $\frac{1}{1000}$ of an inch in thickness, by a single series of double metals of a few inches square: the zinc is circular, forming a small hollow tube, and surrounded by copper opposed to each side of it, and bent so as to correspond to the form of the zinc; when the two metals are exposed to the action of an acid, and connected by the leaf of platinum, the effect is produced.

V. *General Observations, and Conclusion.*

1. Few of the undecomposed bodies, or even of the primary compounds, as is evident from what has been said, are found in an uncombined state in our globe; their tendency to unite with each other is constantly exerted; and a series of decompositions and

recombinations are constantly occurring in the phenomena of nature, and in the operations of art. The compounds containing more than two elements, will form the subjects of consideration for the second part of this work, and their arrangements in the mineral, vegetable, and animal kingdoms; when the principles that have been advanced in the preceding pages, will be applied to the elucidation of an important series of changes belonging to inorganic and to organic matter. As far as our investigations have extended, the same elements belong to the same parts of the system. The composition of the atmosphere and the ocean are analogous, as far as the heights of one, and the depths of the other have been examined. The matters thrown out by volcanoes are earthy or stony aggregates, and they may owe their origin to the action of air and water upon the metallic bases of the earths and alkalies; an action which may be supposed to be connected with the production of subterraneous fires. Even the substances that fall from meteors, though differing in their form and appearance from any of the bodies belonging to our earth, yet contain well known elements, silica, magnesia, sulphur, and the two magnetic metals, iron and nickel.

2. A few undecomposed bodies, which may, perhaps, ultimately be resolved into still fewer elements, or which may be different forms of the same material, constitute the whole of our tangible universe of things. By experiment they are discovered, even in the most complicated arrangements; and experiment is as it were, the chain that binds down the Proteus of nature, and obliges it to confess its real form and divine origin.

The laws which govern the phenomena of chemistry, produce invariable results; which may be made the

guide of operations in the arts ; and which insure the uniformity of the system of nature, the arrangements of which are marked by creative intelligence, and made constantly subservient to the production of life, and the increase of happiness.

ERRATA.

- Page 64, line 21, last word, *for* solids *read* gases.
158, — 21, *for* octohedrous *read* octohedrons.
290, — 13, *for* Morven *read* Morveau.
300, last line, *for* tubes *read* cubes.
324, foot note, *after* 99 *add* chlorine and oxygen being
represented by 35·4 and 8; or by 198, according to
the author's mode of estimation.
217, last line, *for* phosphoretted *read* phosphuretted.
243, line 1, *for* phosphoret *read* phosphuret.

DESCRIPTION OF PLATES.

PLATE I.

- Fig. 1.—Apparatus for shewing the expansion of air by heat, referred to page 48.
- Fig. 2.—Sir John Leslie's differential thermometer.
- Fig. 3.—An instrument described by Van Helmont, resembling in form the differential thermometer; referred to page 14.
- Fig. 4.—Apparatus for shewing that fluids are very bad conductors of heat, described page 55.
- Fig. 5.—Apparatus for shewing the conversion of a fluid into an elastic fluid by heat, referred to page 56.
- Fig. 6.—Apparatus for exploding a mixture of oxygen and hydrogen, (Volta's eudrometer) referred to page 73.

PLATE II.

- Fig. 7.—Apparatus for shewing the combustion and solution of charcoal in oxygen gas, and the formation of carbonic acid gas, described page 74.
- Fig. 8.—Electrical machine, described page 91.
- Fig. 9.—The gold leaf differential electrometer, noticed in p. 93.
- Fig. 10.—Coulomb's electrical balance, described page 94.

DESCRIPTION OF PLATES.

Fig. 11.—The Leyden vial, referred to page 96, where its principle is explained.

PLATE III.

Fig. 12.—Apparatus for shewing electricity by induction, described page 97.

Fig. 13.—Apparatus referred to page 97, explanatory of the common electrical battery.

Fig. 14.—Condensing electrometer, described page 98.

Fig. 15.—The electrical pile of Volta, described page 104.

Fig. 16.—Form of Voltaic battery, referred to page 106, similar to that of the great battery of 2000 double plates, the effects of which are described in page 110.

Fig. 17.—Apparatus for shewing the discharge of Voltaic electricity through air, noticed page 111.

PLATE IV.

Fig. 18.—Apparatus for shewing that the chemical effect of Voltaic electricity is in a certain ratio to the number of plates composing it, described in page 112.

Fig. 19.—Apparatus for shewing the transfer of acid and alkaline matter, in electro-chemical decompositions, described in page 116.

PLATE V.

Fig. 20.—Apparatus for shewing what has been called the radiation of cold, explained in page 149.

PLATE VI.

Fig. 21.—Apparatus for shewing the production of heat in vacuo by Voltaic electricity, described page 150.

Fig. 22, 23.—Apparatus for obtaining oxygen gas from black oxide of manganese, described page 167.

DESCRIPTION OF PLATES.

PLATE VII.

Fig. 24.—Apparatus for burning oxygen in hydrogen, noticed in page 182,—in illustration of the author's doctrine, that combustion, or the disengagement of heat and light, is the effect of combination,—and that, strictly speaking, hydrogen is no more entitled to be considered a combustible than oxygen.

PLATE VIII.

Fig. 25.—Gun-barrel apparatus for procuring potassium, described page 237.

PLATE IX.

Fig. 26.—Represents a furnace for the general purposes of experiments. The upper part of the furnace is a sand bath. The lower part may be employed for fusion or distillation, or for igniting tubes. It may be used in any room where there is a flue, and it serves the purpose of a stove.

PLATE X.

Fig. 27.—Represents the mercurial apparatus. The tube in the frame represents the apparatus for detonating mixtures of gases: it is connected with a spiral spring.

Fig. 28.—Represents another apparatus for detonation.

PLATE XI.

Represents an apparatus for minute experiments; the instruments are delineated of their real size. The cups should be of platinum, the tubes and small retorts of glass; the blow-pipe may be of silver, with an extremity of platinum. By the help of these instruments, and a little wooden trough holding a few pounds of mer-

DESCRIPTION OF PLATES.

cury, an electrophorus for giving a spark to act upon mixtures of gases, or compound gases; and a few bottles containing acids, alkalies, and precipitants, a number of useful experiments may be made.

PLATE XII.

Represents a gasometer, by which a stream of oxygen gas may be thrown upon ignited charcoal, for the purpose of fusing or burning bodies, &c.

Plate I.




Fig. 1.




Fig. 2.




Fig. 3.




Fig. 4.




Fig. 5.



Fig. 6.

Fig. 9.



Fig. 7.




Fig. 8.




Fig. 11.



Fig. 10.




Fig. 17.




Fig. 16




Fig. 13.




Fig. 15.




Fig. 14.







Fig. 12.








Fig. 20.

Fig. 21.




Fig. 23.




Fig. 22.




Fig. 24.





Fig. 25.








Fig. 27.




Fig. 28.

